

First author	Year	Pleural cases	Oncological treatments		
			Type(s)	Number of cases	Details
Boddeart	2015	80	Radiofrequency ablation only (1) at the time of relapse		
Bylicki	2015	68	Chemotherapy	First intent pre-operative chemotherapy (3) (2 of 3 patients had disease progression during treatment)	
				First intent post-operative chemotherapy (7) (Based on De-Perrot staging)	
				First intent exclusive chemotherapy (1)	
				Chemotherapy at recurrence (14) (13 patients received further line of chemotherapy)	
Cardillo	2009	110	Chemotherapy and radiotherapy	Chemotherapy (2) Radiotherapy (4)	Only used in patients who developed recurrence
Chu	2012	40	None		
DeVito	2015	28	Chemotherapy	Chemotherapy (1)	
England	1989	223	Chemotherapy and radiotherapy	Chemotherapy (3) Radiotherapy (7) Chemoradiotherapy (4)	All mSFTP. 8 cases were in combination with surgery, 6 had oncological therapy alone
Guo	2011	39	None		
Gupta	2017	28	None		
Harrison–Phipps	2009	84	Chemotherapy and radiotherapy	Chemotherapy (1) Chemoradiotherapy (1)	Post-operative
Kim	2017	40	Chemotherapy and radiotherapy	Chemotherapy (2) Radiotherapy (1)	Post-operative
Lahon	2012	157	Chemotherapy and radiotherapy	Chemotherapy (1), Chemoradiotherapy (4)	
				Chemotherapy (1) Radiotherapy (3) Chemoradiotherapy (1)	
Liu	2010	21	None		
McGuire	2016	26	Chemotherapy and radiotherapy	Chemotherapy (1) Radiotherapy (1)	At recurrence
Santos	2005	34	Chemotherapy and brachytherapy	Chemotherapy (1) Iodine brachytherapy (3)	Chemotherapy after recurrence of mSFTP
Tan	2018	82	Chemotherapy	Chemotherapy (2)	Post-operative, both mSFTP
Vejvodova	2017	22	Chemotherapy	Chemotherapy (1)	For high malignant potential

Key: mSFTP = malignant solitary fibrous tumour of the pleura

No reference to chemotherapy or radiotherapy in 8 articles (Enon, Franzen, Helage, O'Neil, Rena, Schmid, Tapias, Yeom)

Pleural specific information not available (Demicco 2012, Demicco 2017 and Gholami)