

**Supplementary material 2: Tables showing survey data covered in the report**

<b>How respondents rated the standard of care received for their lung condition while they were pregnant</b>						
	N	Very Poor (%)	Poor (%)	Satisfactory (%)	Good (%)	Excellent (%)
All	126	4.0	11.1	23.0	28.6	33.3
<i>Asthma</i>	50	6.0	18.0	26.0	24.0	26.0
<i>CF</i>	48	–	4.2	18.6	27.1	50.0
<i>LAM</i>	10	-	10.0	-	60.0	30.0
<i>Sarcoidosis</i>	18	11.1	11.1	33.3	33.3	11.1

<b>The extent to which participants said their lung condition influenced the decision of where they gave birth (e.g. hospital, home)</b>							
	N	Not at all (%)	A little (%)	Moderately (%)	A lot (%)	It was the biggest factor (%)	Not relevant to me
All	129	34.1	5.4	5.4	22.5	25.6	7.0
<i>Asthma</i>	48	50	10.4	4.2	16.7	14.6	4.2
<i>CF</i>	47	20.5	2.6	5.1	25.6	38.5	7.7
<i>LAM</i>	11	-	-	9.1	9.1	63.6	18.2
<i>Sarcoidosis</i>	23	52.2	4.3	4.3	26.1	8.7	4.3

<b>The extent to which participants said their lung condition influenced the decision of how they gave birth (e.g. water birth, pain management, caesarean section)</b>							
	N	Not at all (%)	A little (%)	Moderately (%)	A lot (%)	It was the biggest factor (%)	Not relevant to me (%)
All	129	41.9	7.0	10.9	11.6	24.8	3.9
<i>Asthma</i>	48	54.2	8.3	10.4	14.6	12.5	-
<i>CF</i>	47	21.3	4.3	14.9	12.8	40.4	6.4
<i>LAM</i>	11	9.1	9.1	9.1	9.1	45.5	18.2
<i>Sarcoidosis</i>	23	73.9	8.7	4.3	4.3	8.7	-

<b>Whether after giving birth respondents felt they had received support and advice about breastfeeding with a lung condition</b>				
	N	Yes (%)	No (%)	Not relevant to me
All	121	19.8	66.9	13.2
<i>Asthma</i>	45	11.1	68.9	20.0
<i>CF</i>	45	37.8	55.6	6.7
<i>LAM</i>	10	-	90.0	10.0
<i>Sarcoidosis</i>	21	9.5	76.2	14.3

<b>Whether after giving birth respondents felt they had received support and advice about how breastfeeding can increase or reduce the chances of their baby developing a lung condition in future</b>				
	N	Yes (%)	No (%)	Not relevant to me
All	117	17.9	66.7	15.4
<i>Asthma</i>	42	42.9	50.0	7.1
<i>CF</i>	45	4.4	73.3	22.2
<i>LAM</i>	10	-	90.0	10.0
<i>Sarcoidosis</i>	20	5.0	75.0	20.0

<b>How important respondents considered the safety of changing or stopping their lung treatments when pregnant</b>							
	N	Not applicable to me %	Not at all important %	Not very important %	Moderately important %	Very important %	Extremely important %
<b>For your health</b>	127	15.0	9.4	5.5	16.5	22.0	31.6
<i>Asthma</i>	49	2.0	16.3	8.2	26.5	16.3	30.6
<i>CF</i>	48	4.2	4.2	2.1	12.5	31.3	45.8
<i>LAM</i>	10	80.0	10.0	-	-	-	10.0
<i>Sarcoidosis</i>	20	40.0	5.0	10.0	10.0	25.0	10.0
<b>For the baby</b>	127	14.2	4.7	2.4	11.0	16.5	51.2
<i>Asthma</i>	49	-	10.2	4.1	20.4	14.3	51.0
<i>CF</i>	48	4.2	-	-	6.3	25.0	64.6
<i>LAM</i>	10	80.0	-	-	-	-	20.0
<i>Sarcoidosis</i>	20	40.0	5.0	5.0	5.0	10.0	35.0

<b>How important respondents considered the impact of pregnancy and labour on their body and lungs when pregnant</b>							
	N	Not applicable to me %	Not at all important %	Not very important %	Moderately important %	Very important %	Extremely important %
All	127	1.6	5.5	8.7	15.0	33.9	35.4
<i>Asthma</i>	47	-	10.6	8.5	17.0	31.9	31.9
<i>CF</i>	49	-	2.0	6.1	6.1	44.9	8.2
<i>LAM</i>	11	9.1	-	-	9.1	27.3	54.5
<i>Sarcoidosis</i>	20	5.0	5.0	20.0	35.0	15.0	20.0

<b>How important respondents considered the possibility of increased risks during pregnancy because of their lung condition when pregnant</b>							
	N	Not applicable to me %	Not at all important %	Not very important %	Moderately important %	Very important %	Extremely important %
All	128	3.9	6.3	13.3	17.2	28.9	30.5
<i>Asthma</i>	48	-	14.6	16.7	22.9	18.8	27.1
<i>CF</i>	49	-	-	12.2	10.2	44.9	32.7
<i>LAM</i>	11	9.1	-	-	9.1	27.3	54.5
<i>Sarcoidosis</i>	20	15.0	5.0	15.0	30.0	15.0	20.0

<b>Importance given by respondents to the possibility of increased risks during pregnancy because of their lung condition before becoming pregnant</b>							
	N	Not at all important %	Not very important %	Moderately important %	Very important %	Extremely important %	
All	181	18.2	14.9	24.9	21.5	20.4	
<i>Asthma</i>	67	25.4	23.9	26.9	10.4	13.4	
<i>CF</i>	66	9.1	7.6	28.8	28.8	25.6	
<i>LAM</i>	13	23.1	7.7	15.4	23.1	30.8	
<i>Sarcoidosis</i>	35	20.0	14.3	17.1	28.6	20.0	

<b>Extent to which respondents who had not had a pregnancy said that their lung condition influenced their decision and options to have children</b>							
	N	Not at all %	A little %	Moderately %	A lot %	It was the biggest factor %	Not relevant to me %
All	83	4.8	6.0	9.6	18.1	53.0	8.4
<i>Asthma</i>	14	21.4	7.1	21.4	21.4	14.3	14.3
<i>CF</i>	51	-	7.8	5.9	15.7	62.7	7.8

<i>LAM</i>	7	-	-	-	14.3	85.7	-
<i>Sarcoidosis</i>	11	9.1	-	18.2	27.3	36.4	9.1

**How respondents rated the information received from different sources about pregnancy and their lung condition before they were pregnant and during the pregnancy**

		N	Not relevant to me %	Very poor %	Poor %	Satisfactory %	Good %	Excellent %
Primary care professionals	<b>Before pregnancy</b>	181	20.4	8.8	18.2	24.9	18.2	9.4
	<b>During pregnancy</b>	128	12.5	8.6	16.4	23.4	28.1	10.9
Respiratory professionals	<b>Before pregnancy</b>	179	19.6	6.1	8.4	17.9	24.6	23.5
	<b>During pregnancy</b>	128	18.8	7.0	9.3	12.5	24.2	28.1

**Actions taken by respondents before becoming pregnant**

	N	Not applicable (%)	Yes (%)	No (%)
<b>Speak to a primary care professional (e.g. nurse, GP or family doctor) about becoming pregnant with your lung condition</b>	183	7.7	48.1	44.3
<b>Speak to a respiratory professional (e.g. respiratory consultant, respiratory nurse) about becoming pregnant with your lung condition</b>	183	7.7	60.7	31.7