



Early View

Original article

High-sensitivity troponin I for risk stratification in normotensive pulmonary embolism

Matthias Ebner, Niklas Guddat, Karsten Keller, Marie Christine Merten, Markus H. Lerchbaumer, Gerd Hasenfuß, Stavros V. Konstantinides, Mareike Lankeit

Please cite this article as: Ebner M, Guddat N, Keller K, *et al.* High-sensitivity troponin I for risk stratification in normotensive pulmonary embolism. *ERJ Open Res* 2020; in press (<https://doi.org/10.1183/23120541.00625-2020>).

This manuscript has recently been accepted for publication in the *ERJ Open Research*. It is published here in its accepted form prior to copyediting and typesetting by our production team. After these production processes are complete and the authors have approved the resulting proofs, the article will move to the latest issue of the ERJOR online.

Copyright ©ERS 2020. This article is open access and distributed under the terms of the Creative Commons Attribution Non-Commercial Licence 4.0.

High-sensitivity troponin I for risk stratification in normotensive pulmonary embolism

Authors:

Matthias Ebner, MD^{1,2}; Niklas Guddat³; Karsten Keller, MD^{4,5}; Marie Christine Merten, MD³; Markus H. Lerchbaumer, MD⁶, Gerd Hasenfuß, MD^{3,7}, Stavros V. Konstantinides, MD^{4,8}; Mareike Lankeit, MD^{2,3,4,9}

Affiliations:

¹ Department of Cardiology and Angiology, Campus Charité Mitte (CCM), Charité – University Medicine Berlin, Germany

² German Center for Cardiovascular Research (DZHK), partner site Berlin, Germany

³ Clinic of Cardiology and Pneumology, University Medical Center Goettingen, Germany

⁴ Center for Thrombosis and Hemostasis (CTH), University Medical Center Mainz, Johannes Gutenberg-University Mainz, Germany

⁵ Center for Cardiology, Cardiology I, University Medical Center Mainz, Johannes Gutenberg-University Mainz, Germany

⁶ Department of Radiology, Campus Charité Mitte (CCM), Charité – University Medicine Berlin, Germany

⁷ German Cardiovascular Research Centre (DZHK), partner site Goettingen, Germany

⁸ Department of Cardiology, Democritus University of Thrace, Alexandroupolis, Greece

⁹ Department of Internal Medicine and Cardiology, Campus Virchow Klinikum (CVK), Charité
– University Medicine Berlin, Germany

Corresponding author:

Mareike Lankeit, Department of Internal Medicine and Cardiology, Campus Virchow Klinikum (CVK), Charité – University Medicine Berlin, Augustenburger Platz 1, 13353 Berlin, Germany. Phone: +49 (0)30 / 450 665 381, fax: +49 (0)30 / 450 7 565 381, email: mareike.lankeit@charite.de

Take home message:

The study confirms the prognostic relevance of high-sensitivity troponin I in normotensive pulmonary embolism. A cut-off value of 16 pg/ml can be used for risk-stratification in male and female patients; sex-specific adjustments do not appear necessary.

Word count:

Take home message:	253 characters (max. 256 characters)
Abstract	250 words (max. 250 words)
Manuscript	2,939 words (max. 3,000 words)
Tables / Figures	4 / 1
References	26
Supplemental Material:	1 Table / 1 Figure

Abstract

While numerous studies confirmed the prognostic role of high-sensitivity troponin T (hsTnT) in pulmonary embolism (PE), high-sensitivity troponin I (hsTnI) is inappropriately studied. This study aimed to investigate the prognostic relevance of hsTnI in normotensive PE, establish the optimal cut-off value for risk stratification and compare the prognostic performances of hsTnI and hsTnT.

Based on data from 459 consecutive PE patients enrolled in a single-centre registry, receiver operating characteristic analysis was used to identify an optimal hsTnI cut-off value for prediction of in-hospital adverse outcomes (PE-related death, cardiopulmonary resuscitation or vasopressor treatment) and all-cause mortality.

Patients who suffered an in-hospital adverse outcome (4.8%) had higher hsTnI concentrations compared to those with a favourable clinical course (57 [IQR 22-197] vs. 15 [IQR 10-86] pg/ml, $p=0.03$). A hsTnI cut-off value of 16 ng/ml provided optimal prognostic performance and predicted in-hospital adverse outcomes (Odds ratio [OR] 6.5, 95% confidence interval [CI] 1.9-22.4) and all-cause mortality (OR 3.7, 95% CI 1.0-13.3). Between female and male patients, no relevant differences in hsTnI concentrations (17 [IQR 10-97] vs. 17 [IQR 10-92] pg/ml, $p=0.79$) or optimised cut-off values were observed. Risk stratification according to the 2019 European Society of Cardiology algorithm revealed no differences if calculated based on either hsTnI or hsTnT ($p=0.68$).

Our findings confirm the prognostic role of hsTnI in normotensive PE. HsTnI concentrations >16 pg/ml predicted in-hospital adverse outcome and all-cause mortality; sex-specific cut-off values do not appear necessary. Importantly, our results suggest that hsTnI and hsTnT can be used interchangeably for risk stratification.

Introduction

Pulmonary embolism (PE) is a life-threatening disease with high morbidity and mortality. [1, 2] Current guideline recommendations emphasize the importance of early risk stratification in the heterogeneous group of normotensive patients with acute PE to guide therapeutic decision making. [3]

Consensus exists that right ventricular (RV) dysfunction is the critical determinant of outcome in PE. [3] In the absence of chronic pulmonary hypertension, the RV has a narrow range to handle an acute increase in afterload caused by the occlusion of pulmonary vasculature during acute PE. [4] If the adaptive potential of the RV is exceeded, RV dilation and elevated wall tension occurs. [5, 6] Consequently, coronary perfusion of the RV myocardium is impaired, leading to ischemia that further aggravates RV dysfunction and ultimately cardiac failure. [6] Thus, detection of myocardial injury indicated by cardiac troponin elevation is a key element for identification of normotensive patients at higher risk for short-term adverse outcomes. [3, 7-9]

High-sensitivity troponin assays have largely replaced conventional assays due to their increased ability to detect myocardial injury. [10] While a large number of studies confirmed the prognostic relevance of high-sensitivity troponin T (hsTnT) in acute PE and established cut-off values defining elevated hsTnT concentrations [8, 11], only a single study investigated high-sensitivity troponin I (hsTnI) for risk stratification in a small cohort of 65 unselected PE patients. [12] Therefore, the prognostic relevance of hsTnI elevation in patients with acute PE is inappropriately studied and disease specific cut-off values remain undefined.

In the present study we aimed to investigate the prognostic relevance of hsTnI, establish the optimal hsTnI cut-off value and compare the prognostic performances of hsTnI and hsTnT for risk stratification of normotensive patients with acute PE.

Methods

Study design and definition of outcomes

The Pulmonary Embolism Registry of Göttingen (PERGO) prospectively includes consecutive patients with objectively confirmed PE ≥ 18 years of age admitted to the University Medical Center Göttingen, Germany. The study protocol has been described in detail previously. [13, 14] Briefly, patient recruitment is performed by daily screening of new admissions to the emergency department and of reports of conducted computed tomography pulmonary angiographies (CTPA). After obtaining informed consent for participation in PERGO, complete data on comorbidities, previous medication, symptoms, results from diagnostic tests (including laboratory parameters, electrocardiogram [ECG], transthoracic echocardiography [TTE] and CTPA), treatment and clinical course is obtained using a standardised case report form. In addition, adjudication of parameters defining RV dysfunction is performed by an independent investigator by reviewing the electronically stored TTE and CTPA images. The present analysis included patients enrolled in PERGO between September 2008 and April 2018. We excluded patients with (1) missing hsTnI measurements at presentation, (2) cardiac arrest or haemodynamic instability at presentation and (3) another acute cardio-respiratory illness such as acute myocardial infarction, left heart decompensation or respiratory decompensation responsible for clinical presentation and symptoms. All patients were followed for the in-hospital stay.

Diagnostic and therapeutic management was in accordance with the ESC 2008 (09/2008-08/2014) and 2014 (09/2014-04/2018) guidelines [15, 16] and local standard operating procedures. All related decisions were left to the discretion of the treating physicians and were not influenced by the study protocol. Treating physicians were not informed about study results, thus any influence of the study on patient management or

monitoring of treatment effects can be excluded. The study was conducted in accordance with the amended Declaration of Helsinki and approved by the local independent Ethic Committee of the University Medical Center Göttingen, Germany. All patients gave informed written consent for participation in the study.

Tachycardia was defined as heart rate of ≥ 100 beats per minute (bpm) and hypoxaemia as peripheral oxygen saturation $< 90\%$. Renal insufficiency was defined as glomerular filtration rate (GFR) < 60 ml/min/1.73 m² body surface area. Altered mental status was defined as disorientation, somnolence, sopor or coma. Active cancer was defined as known malignant disease, treatment with antitumour therapy within the last 6 months, metastatic state or haematologic cancer not in complete remission. [17] RV dysfunction on CTPA was defined as right-to-left ventricular (RV/LV) diameter ratio ≥ 1.0 . [3] Patients were stratified to risk classes post-hoc according to the algorithm proposed by the ESC 2019 guidelines and the modified FAST score. [3, 9] The modified FAST score includes elevated troponin (1.5 points), tachycardia (2 points) and syncope (1.5 points); a patient is classified as high-risk if ≥ 3 points are scored. For calculation of all algorithms and scores, missing values were considered to be normal. [18]

An in-hospital adverse outcome was defined as PE-related death or haemodynamic collapse (cardiopulmonary resuscitation or administration of catecholamines). The second study outcome was in-hospital all-cause mortality. Death was determined to be PE-related if either confirmed by autopsy or following a clinically severe episode of acute PE in absence of an alternative diagnosis. All events and causes of death were independently adjudicated by two of the authors (M.E. and N.G.) and disagreement was resolved by a third author (M.L.).

Biomarker measurements

Venous blood samples were collected on admission or at the moment of PE diagnosis, processed using standard operating procedures and immediately stored at -80°C . Plasma concentrations of hsTnI (ARCHITECT stat hsTnI assay, Abbott Laboratories, Chicago, IL, USA), hsTnT (Roche Diagnostics, Mannheim, Germany) and N-terminal pro-brain natriuretic peptide (NT-proBNP; Roche Diagnostics, Mannheim, Germany) were measured in batches after a single thaw by the *amedes MVZ wagnerstibbe* laboratory in Göttingen, Germany. For hsTnI, the following predefined cut-off values were investigated: (1) >10 pg/ml (lower detection limit of the assay), (2) >27 pg/ml (99th percentile in healthy individuals) [19], (3) >16 pg/ml in women and >34 pg/ml in men (sex-specific 99th percentile in healthy individuals) [20] and (4) >100 pg/ml (previously evaluated for risk stratification in acute PE) [12]. For other biomarkers, cut-off values indicating elevated concentrations were prospectively defined as hsTnT ≥ 14 pg/ml [18] and NT-proBNP ≥ 600 pg/ml [21]. Serial biomarker measurements were not performed.

Statistical analysis

Categorical variables are presented as total numbers and percentages and continuous variables are presented as medians with interquartile ranges (IQR). Associations between binary and categorical variables were analysed using Fisher's exact test, Chi-square test or the Mantel-Haenszel test of trend, as appropriate. Spearman's rank correlation coefficient was used to test for statistical dependence of hsTnI from continuous variables. For comparison of continuous variables, the Mann-Whitney U-test was employed.

To evaluate the overall prognostic performance of hsTnI with regard to study outcomes, receiver operating characteristic (ROC) curve analyses were performed to determine the area under the curve (AUC) with corresponding 95% confidence intervals (CI)

in the whole study cohort and in patients stratified according to sex. The optimal cut-off value for prediction of study outcomes was identified based on Youden index quantification and compared to the established hsTnI and hsTnT cut-off values.

Further, the prognostic value of selected cut-off values, patient characteristics, comorbidities and biomarkers with regard to study outcomes was tested using univariable logistic regression analyses and results are presented as odds ratios (OR) with the corresponding 95% CI. To confirm the independent prognostic value of hsTnI, hsTnI levels dichotomised according to the optimal cut-off value were entered in three multivariable logistic regression models correcting for (I) presence of coronary artery disease, (II) renal insufficiency and (III) age ≥ 75 years.

To test for differences in the prognostic performance of the ESC 2019 algorithm and the modified FAST score if calculated based on either hsTnT or hsTnI levels, the net reclassification improvement (NRI) with corresponding standard error (SE) was calculated. [22] Kaplan-Meier analysis was used to compare the probability of one-year survival in subgroups stratified according to hsTnI levels at presentation; the log-rank test was used for comparison between groups.

A two-sided significance level of $\alpha < 0.05$ was defined as appropriate to indicate statistical significance. As this was explorative testing, no adjustments for multiple testing were carried out. P-values were provided for descriptive reasons only and should be interpreted with caution and in connection with effect estimates. Statistical analysis was performed through Statistics Package for Social Sciences (IBM SPSS Statistics, version 26, IBM Corp. Armonk, NY, USA).

Results

Between September 2008 and April 2018, 854 patients were enrolled in PERGO. Exclusion criteria applied to (1) 283 patients with missing hsTnI measurements, (2) 88 patients with cardiogenic shock or haemodynamic instability at presentation and (3) 24 patients with significant acute cardio-respiratory illness responsible for clinical presentation and symptoms. Hence, 459 patients (53.7%) were included in the present analysis. A comparison of the study cohort with patients excluded due to missing hsTnI measurements is provided in the Supplementary Material (**Table s1**).

At presentation, 58 (12.6%) patients were classified as low-, 265 (57.7%) as intermediate-low- and 136 (29.6%) as intermediate-high-risk according to the ESC 2019 risk stratification algorithm. Reperfusion treatment was performed in 10 (2.2%) patients and seven (1.5%) were enrolled in the PEITHO trial. [23] An in-hospital adverse outcome occurred in 22 (4.8%) patients. Overall, 14 (3.1%) patients died during the in-hospital stay; of those, nine (64.3%) due to PE. Information on baseline characteristics, comorbidities, initial presentation and outcomes in are shown in **Table 1**, left column.

The median biomarker concentrations on admission were 17.0 (IQR 9.9-96.0) pg/ml for hsTnI, 20.0 (IQR 8.7-56.7) pg/ml for hsTnT and 499 (IQR 108-2,386) pg/ml for NT-proBNP. More specifically, 41.6% of patients had hsTnI concentrations >27 pg/ml and 23.5% of patients >100 pg/ml, while elevated hsTnT concentrations ≥ 14 pg/ml were observed in 61.0% of patients. A strong correlation between hsTnI and hsTnT concentrations ($r=0.82$, $p<0.001$) was observed. Furthermore, hsTnI concentrations showed a positive correlation with age ($r=0.31$, $p<0.001$) and heart rate ($r=0.20$, $p<0.001$) and an inverse correlation with GFR ($r=-0.32$, $p<0.001$).

Identification of the optimal hsTnI cut-off value for risk stratification

Patients who developed an in-hospital adverse outcome had higher hsTnI concentrations compared to patients with a favourable clinical course (57 [IQR 22-197] pg/ml vs. 15 [IQR 10-86] pg/ml, $p=0.030$). Further, the rate of in-hospital adverse outcomes increased with higher hsTnI concentrations at presentation ($p=0.002$ for trend; **Figure s1** of the Supplementary Material).

ROC analyses indicated an association between hsTnI and in-hospital adverse outcome (AUC 0.68 [95% CI 0.59-0.77]; **Figure 1**) and in-hospital all-cause mortality (AUC 0.64 [95% CI 0.51-0.77]). Using Youden index quantification, an optimal cut-off value of 17 pg/ml for the identification of patients with an in-hospital adverse outcomes was identified. As this value is numerically almost identical to the 99th percentile of hsTnI concentrations in healthy women (16 pg/ml) and Youden's indices for the two cut-off values differed only minimally (0.38 and 0.37, respectively), we selected a cut-off value of 16 pg/ml for all further analysis.

Patients with hsTnI levels >16 pg/ml were older, had higher rates of chronic heart failure, coronary artery disease and renal insufficiency as well as higher rates of dyspnoea, syncope, tachycardia, hypoxaemia and signs of RV dysfunction on diagnostic imaging compared to patients with hsTnI levels ≤16 pg/ml (**Table 1**, right columns).

Evaluation of sex-specific differences

No difference in the median hsTnI concentrations between female and male PE patients (17.0 [IQR 9.9-97.2] vs. 16.5 [IQR 9.9-92.3] pg/ml, $p=0.79$) was observed and Youden index

quantification indicated similar optimal hsTnI cut-off values for both sexes (17 pg/ml for female and 19 pg/ml for male patients). However, a larger AUC with regard to an in-hospital adverse outcome was calculated for female (0.74 [95% CI 0.63-0.85]) than for male patients (0.62 [95% CI 0.47-0.76]).

Prognostic performance of the optimal hsTnI cut-off value

Using univariable logistic regression analysis, hsTnI >16 pg/ml predicted an in-hospital adverse outcome (OR 6.5 [95% CI 1.9-22.4]; **Table 2**, middle columns) as well as all-cause mortality (OR 3.7 [95% CI 1.0-13.3]; **Table 2**, right columns). The independent prognostic value of hsTnI with regard to the primary outcome was confirmed in three separate multivariable models, correcting for the presence of (I) coronary artery disease (OR 6.5 [95% CI 1.9-22.6]), (II) renal insufficiency (OR 4.7 [95% CI 1.3-16.6]) or (III) age \geq 75 years (OR 6.0 [95% CI 1.7-20.9]).

A comparison of test characteristics for the newly identified PE-specific cut-off value of 16 pg/ml and all predefined hsTnI and hsTnT cut-off values is provided in **Table 3**.

Prognostic performance of risk stratification based on hsTnT or hsTnI measurements

A comparison of the prognostic performance of the modified FAST score and the ESC 2019 algorithm calculated based on (I) hsTnI >16 pg/ml and (II) hsTnT \geq 14 pg/ml is provided in **Table 4**. Calculations of sensitivity, specificity and odds ratios for outcome prediction did not reveal relevant differences in test performance. Further, no difference in the net reclassification improvement was observed.

Discussion

In the present study we evaluated the prognostic value of hsTnI in normotensive PE patients and established a disease-specific hsTnI cut-off value for risk stratification. Our findings obtained in 459 patients can be summarised as follows: I) hsTnI concentrations >16 pg/ml were predictive of both, an in-hospital adverse outcome and all-cause mortality; II) sex-specific cut-off values do not appear necessary for risk stratification of acute PE; III) if hsTnI and hsTnT were used to risk-stratify patients according to the modified FAST score and ESC 2019 algorithm, no difference in the prognostic performance was observed.

High-sensitivity troponin assays are recommended for the diagnosis of myocardial infarction in current European and American guidelines and have seen high adoption rates in clinical practice. [10, 24] Although these contemporary assays improve the diagnostic accuracy at low analyte concentrations, their increase in sensitivity comes at the disadvantage of reduced specificity. [25] This emphasizes the importance of cut-off values tailored to specific disease states and applications. In patients with suspected acute coronary syndrome (ACS), troponin concentrations below the 99th percentile upper reference limit (URL) are used to rule-out non-ST-elevation ACS. [10, 24] In contrast, troponin is not measured for diagnostic purposes in patients with acute PE but contributes to risk stratification *after* the diagnosis has been established. Hence, the diagnostic troponin threshold for myocardial infarction might not be identical to the optimal cut-off value for risk stratification in acute PE.

Data on disease-specific troponin I cut-off values for risk stratification of acute PE are limited, as most previous reports relied on the 99th percentile URL. [7, 26] In addition, the majority of investigations relied on conventional troponin I assays rather than highly sensitive ones. [7, 26] Until now, the only evidence supporting the use of hsTnI for risk

stratification of acute PE comes from a single study including 65 patients only. [12] Importantly, this study investigated the cut-off value suggested for evaluation of ACS at the time of publication (100 pg/ml) [12], that is much higher than contemporary thresholds [20].

Our study adds substantially to these previous investigations. Based on our results, a hsTnI cut-off value of 16 pg/ml provides high sensitivity (86.4%) and a high negative predictive value (NPV; 98.7%) for the prediction of adverse outcomes and thus offers better prognostic performance than cut-off values corresponding to the 99th percentile URL [19], the sex-specific 99th percentile URL [20] and the cut-off value suggested by Walter et al. [12]. Of note, the identified optimal hsTnI cut-off value is lower than the 99th percentile URL reported by the manufacturer [19] but identical to the 99th percentile URL reported for females and higher than cut-off values recommended to rule-out non-ST-elevation ACS [10] and for risk stratification of asymptomatic patients with cardiovascular disease [19]. Thus, potentially more male patients with PE will be classified as intermediate-high risk requiring hospital admission and monitoring, which might increase safety and is very unlikely to cause harm. However, as a lower cut-off value may lead to more false positive results (and hence potentially overtreatment), the costs and effectiveness associated with a higher number of PE patients classified as intermediate-high-risk require additional investigation in further studies.

Interestingly, although sex-specific differences of hsTnI values have been described for patients with myocardial infarction [20], we did not observe such differences in our cohort of PE patients. Optimal cut-off values for female and male patients were similar (17 pg/ml and 19 pg/ml, respectively) and hence, do not indicate the use of sex-specific cut-off values for risk stratification of acute PE.

The recently published ESC 2019 guidelines emphasize the central role of troponin elevation for risk stratification in acute PE and consider the troponin subunits I and T as equally suitable for this purpose. [3] This is based on the results of two meta-analyses concluding that the prognostic value of increased troponin concentrations in patients with acute PE is independent regardless of whether troponin T or I subunits are measured. [7, 26] The findings of our study are consistent with these previous reports and add to the evidence that hsTnI and hsTnT have comparable prognostic performance. Hence, clinicians are free to choose either assay and troponin subunit for risk stratification of normotensive PE in clinical practice.

Limitations

The present study is a post-hoc analysis of a prospective single-centre cohort, thus the generalisability of the study findings may be limited. In particular, the high prevalence of comorbidities such as cancer (24.0%) and renal insufficiency (29.2%) should be taken into account when interpreting the present study findings. A considerable number of patients were excluded from the study due to missing hsTnI measurements at presentation. However, the comparison of patients with and without hsTnI measurements provided in **Table s1** in the Supplementary Material did not reveal relevant differences between the two groups. Finally, due to the small number of events, the ability to perform robust multivariable analyses was limited.

Conclusion

Our findings confirm the prognostic value of hsTnI for risk stratification of patients with normotensive PE. Elevation of hsTnI concentrations exceeding a cut-off value of 16 pg/ml predict in-hospital adverse outcome and all-cause mortality in both male and female patients. Furthermore, our results suggest that hsTnI and hsTnT have similar prognostic value and can be used interchangeably for risk stratification.

Acknowledgements

The authors thank Dr. Christian Thode and Daniela Brehm (amedes MVZ wagnerstibbe, Göttingen, Germany) for performing the biomarker measurements.

This research was supported by the UMG Biobank as a core facility of the University Medical Center Göttingen, Germany by provision of quality assured biomaterial.

This publication is part of the medical doctoral thesis of Niklas Guddat.

Funding

The work of Karsten Keller, Stavros V. Konstantinides and Mareike Lankeit was supported by the German Federal Ministry of Education and Research (BMBF 01EO1503). The authors are responsible for the contents of this publication.

BRAHMS GmbH, part of Thermo Fisher Scientific, Hennigsdorf/Berlin, Germany provided financial support for biomarker measurements. The sponsor was neither involved in biomarker measurements, statistical analyses, writing of the manuscript nor had any influence on the scientific contents of this publication.

Conflicts of interest

None of the authors reports a relationship with industry and other relevant entities – financial or otherwise – that might pose a conflict of interest in connection with the submitted article. The following authors report financial activities outside the submitted work:

Matthias Ebner reports no conflicts of interest.

Niklas Guddat reports no conflict of interest.

Karsten Keller reports no conflicts of interest.

Marie Christine Merten reports no conflict of interest.

Markus H. Lerchbaumer reports having received consultancy honoraria from Siemens Healthineers.

Gerd Hasenfuß reports having received consultancy and lecture honoraria from AstraZeneca, Berlin Chemie, Corvia, Impulse Dynamics, Novartis, Servier and Vifor Pharma; and editor honoraria from Springer International Publishing AG.

Stavros V. Konstantinides reports having received consultancy and lecture honoraria from Bayer, Boehringer Ingelheim, Daiichi-Sankyo, MSD and Pfizer – Bristol-Myers Squibb; and institutional grants from Actelion, Bayer, Boehringer Ingelheim, Daiichi-Sankyo and Pfizer – Bristol-Myers Squibb.

Mareike Lankeit reports having received consultancy and lecture honoraria from Actelion, Bayer, BRAHMS – Thermo Fisher Scientific, Daiichi-Sankyo, MSD and Pfizer – Bristol-Myers Squibb and project funding from BRAHMS – Thermo Fisher Scientific.

References

1. Wendelboe AM, Raskob GE. Global Burden of Thrombosis: Epidemiologic Aspects. *Circulation research* 2016; 118: 1340-1347.
2. Keller K, Hobohm L, Ebner M, Kresoja KP, Münzel T, Konstantinides S, Lankeit M. Trends in thrombolytic treatment and outcomes of acute pulmonary embolism in Germany. *Eur Heart J* 2019.
3. Konstantinides SV, Meyer G, Becattini C, Bueno H, Geersing GJ, Harjola VP, Huisman MV, Humbert M, Jennings CS, Jimenez D, Kucher N, Lang IM, Lankeit M, Lorusso R, Mazzolai L, Meneveau N, Ainle FN, Prandoni P, Pruszczyk P, Righini M, Torbicki A, Van Belle E, Zamorano JL. 2019 ESC Guidelines for the diagnosis and management of acute pulmonary embolism developed in collaboration with the European Respiratory Society (ERS): The Task Force for the diagnosis and management of acute pulmonary embolism of the European Society of Cardiology (ESC). *Eur Respir J* 2019; 54.
4. Ribeiro A, Lindmarker P, Juhlin-Dannfelt A, Johnsson H, Jorfeldt L. Echocardiography Doppler in pulmonary embolism: right ventricular dysfunction as a predictor of mortality rate. *American heart journal* 1997; 134: 479-487.
5. Vonk Noordegraaf A, Westerhof BE, Westerhof N. The Relationship Between the Right Ventricle and its Load in Pulmonary Hypertension. *J Am Coll Cardiol* 2017; 69: 236-243.
6. Harjola VP, Mebazaa A, Celutkiene J, Bettex D, Bueno H, Chioncel O, Crespo-Leiro MG, Falk V, Filippatos G, Gibbs S, Leite-Moreira A, Lassus J, Masip J, Mueller C, Mullens W, Naeije R, Nordegraaf AV, Parissis J, Riley JP, Ristic A, Rosano G, Rudiger A, Ruschitzka F, Seferovic P, Sztrymf B, Vieillard-Baron A, Yilmaz MB, Konstantinides S. Contemporary management of acute right ventricular failure: a statement from the Heart Failure Association and the Working Group on Pulmonary Circulation and Right Ventricular Function of the European Society of Cardiology. *Eur J Heart Fail* 2016; 18: 226-241.
7. Becattini C, Vedovati MC, Agnelli G. Prognostic value of troponins in acute pulmonary embolism: a meta-analysis. *Circulation* 2007; 116: 427-433.
8. Kaeberich A, Seeber V, Jimenez D, Kostrubiec M, Dellas C, Hasenfuss G, Giannitsis E, Pruszczyk P, Konstantinides S, Lankeit M. Age-adjusted high-sensitivity troponin T cut-off value for risk stratification of pulmonary embolism. *The European respiratory journal* :

- official journal of the European Society for Clinical Respiratory Physiology 2015: 45: 1323-1331.
9. Hobohm L, Hellenkamp K, Hasenfuss G, Munzel T, Konstantinides S, Lankeit M. Comparison of risk assessment strategies for not-high-risk pulmonary embolism. *Eur Respir J* 2016; 47: 1170-1178.
 10. Roffi M, Patrono C, Collet JP, Mueller C, Valgimigli M, Andreotti F, Bax JJ, Borger MA, Brotons C, Chew DP, Gencer B, Hasenfuss G, Kjeldsen K, Lancellotti P, Landmesser U, Mehilli J, Mukherjee D, Storey RF, Windecker S, Group ESCSD. 2015 ESC Guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation: Task Force for the Management of Acute Coronary Syndromes in Patients Presenting without Persistent ST-Segment Elevation of the European Society of Cardiology (ESC). *Eur Heart J* 2016; 37: 267-315.
 11. Lankeit M, Friesen D, Aschoff J, Dellas C, Hasenfuss G, Katus H, Konstantinides S, Giannitsis E. Highly sensitive troponin T assay in normotensive patients with acute pulmonary embolism. *European heart journal* 2010; 31: 1836-1844.
 12. Walter T, Apfaltrer P, Weilbacher F, Meyer M, Schoenberg SO, Fink C, Gruettner J. Predictive value of high-sensitivity troponin I and D-dimer assays for adverse outcome in patients with acute pulmonary embolism. *Exp Ther Med* 2013; 5: 586-590.
 13. Ebner M, Kresoja KP, Keller K, Hobohm L, Rogge NIJ, Hasenfuss G, Pieske B, Konstantinides SV, Lankeit M. Temporal trends in management and outcome of pulmonary embolism: a single-centre experience. *Clin Res Cardiol* 2019.
 14. Lankeit M, Friesen D, Schafer K, Hasenfuss G, Konstantinides S, Dellas C. A simple score for rapid risk assessment of non-high-risk pulmonary embolism. *Clin Res Cardiol* 2013; 102: 73-80.
 15. Torbicki A, Perrier A, Konstantinides S, Agnelli G, Galie N, Pruszczyk P, Bengel F, Brady AJ, Ferreira D, Janssens U, Klepetko W, Mayer E, Remy-Jardin M, Bassand JP, Guidelines ESCCfP. Guidelines on the diagnosis and management of acute pulmonary embolism: the Task Force for the Diagnosis and Management of Acute Pulmonary Embolism of the European Society of Cardiology (ESC). *Eur Heart J* 2008; 29: 2276-2315.
 16. Konstantinides SV, Torbicki A, Agnelli G, Danchin N, Fitzmaurice D, Galie N, Gibbs JS, Huisman MV, Humbert M, Kucher N, Lang I, Lankeit M, Lekakis J, Maack C, Mayer E, Meneveau N, Perrier A, Pruszczyk P, Rasmussen LH, Schindler TH, Svitil P, Vonk

- Noordegraaf A, Zamorano JL, Zompatori M, Task Force for the D, Management of Acute Pulmonary Embolism of the European Society of C. 2014 ESC guidelines on the diagnosis and management of acute pulmonary embolism. *Eur Heart J* 2014; 35: 3033-3069, 3069a-3069k.
17. Raskob GE, van Es N, Segers A, Angchaisuksiri P, Oh D, Boda Z, Lyons RM, Meijer K, Gudz I, Weitz JI, Zhang G, Lanz H, Mercuri MF, Buller HR, Hokusai VTEi. Edoxaban for venous thromboembolism in patients with cancer: results from a non-inferiority subgroup analysis of the Hokusai-VTE randomised, double-blind, double-dummy trial. *Lancet Haematol* 2016; 3: e379-387.
 18. Lankeit M, Jimenez D, Kostrubiec M, Dellas C, Hasenfuss G, Pruszczyk P, Konstantinides S. Predictive value of the high-sensitivity troponin T assay and the simplified Pulmonary Embolism Severity Index in hemodynamically stable patients with acute pulmonary embolism: a prospective validation study. *Circulation* 2011; 124: 2716-2724.
 19. Beshiri A, Ravalico T. A New Paradigm for Acute Coronary Syndrome: High Sensitive Troponin. www.corelaboratory.abbott/sal/whitePaper/CR_ADD-00001076_ARCHITECT_hsTnI_WhitePaper_072613.pdf. Date last updated: 2013. Date last accessed: Nov 14 2019.
 20. Shah ASV, Anand A, Strachan FE, Ferry AV, Lee KK, Chapman AR, Sandeman D, Stables CL, Adamson PD, Andrews JPM, Anwar MS, Hung J, Moss AJ, O'Brien R, Berry C, Findlay I, Walker S, Cruickshank A, Reid A, Gray A, Collinson PO, Apple FS, McAllister DA, Maguire D, Fox KAA, Newby DE, Tuck C, Harkess R, Parker RA, Keerie C, Weir CJ, Mills NL, High SI. High-sensitivity troponin in the evaluation of patients with suspected acute coronary syndrome: a stepped-wedge, cluster-randomised controlled trial. *Lancet* 2018; 392: 919-928.
 21. Lankeit M, Jimenez D, Kostrubiec M, Dellas C, Kuhnert K, Hasenfuss G, Pruszczyk P, Konstantinides S. Validation of N-terminal pro-brain natriuretic peptide cut-off values for risk stratification of pulmonary embolism. *Eur Respir J* 2014; 43: 1669-1677.
 22. Pencina MJ, D'Agostino RB, Sr., D'Agostino RB, Jr., Vasan RS. Evaluating the added predictive ability of a new marker: from area under the ROC curve to reclassification and beyond. *Stat Med* 2008; 27: 157-172; discussion 207-112.
 23. Meyer G, Vicaut E, Danays T, Agnelli G, Becattini C, Beyer-Westendorf J, Bluhmki E, Bouvaist H, Brenner B, Couturaud F, Dellas C, Empen K, Franca A, Galie N, Geibel A,

- Goldhaber SZ, Jimenez D, Kozak M, Kupatt C, Kucher N, Lang IM, Lankeit M, Meneveau N, Pacouret G, Palazzini M, Petris A, Pruszczyk P, Rugolotto M, Salvi A, Schellong S, Sebbane M, Sobkowicz B, Stefanovic BS, Thiele H, Torbicki A, Verschuren F, Konstantinides SV, Investigators P. Fibrinolysis for patients with intermediate-risk pulmonary embolism. *N Engl J Med* 2014; 370: 1402-1411.
24. Amsterdam EA, Wenger NK, Brindis RG, Casey DE, Jr., Ganiats TG, Holmes DR, Jr., Jaffe AS, Jneid H, Kelly RF, Kontos MC, Levine GN, Liebson PR, Mukherjee D, Peterson ED, Sabatine MS, Smalling RW, Zieman SJ. 2014 AHA/ACC Guideline for the Management of Patients with Non-ST-Elevation Acute Coronary Syndromes: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol* 2014; 64: e139-e228.
25. Januzzi JL, Jr. What to expect when measuring high-sensitivity troponin: practical advice for clinicians. *J Am Coll Cardiol* 2015; 65: 1665-1667.
26. Bajaj A, Saleeb M, Rathor P, Sehgal V, Kabak B, Hosur S. Prognostic value of troponins in acute nonmassive pulmonary embolism: A meta-analysis. *Heart Lung* 2015; 44: 327-334.

Tables

Table 1. Comorbidities, results from risk stratification and outcomes of study patients stratified according to the optimal hsTnI cut-off value

	All patients (n=459)	hsTnI >16 pg/ml (n=234)	hsTnI ≤16 pg/ml (n=225)	p-value
Age ≥75 years	155 (33.8%)	102 (43.6%)	53 (23.6%)	<0.001
Sex (female)	241 (52.5%)	125 (53.4%)	116 (51.6%)	0.60
Comorbidities				
Chronic heart failure	59 (12.9%)	43 (18.4%)	16 (7.1%)	<0.001
Coronary artery disease	67 (14.6%)	46 (19.7%)	21 (9.3%)	0.002
Chronic pulmonary disease	60 (13.1%), n=458	27 (11.6%), n=233	33 (14.7%)	0.33
Diabetes mellitus	65 (14.2%)	39 (16.7%)	26 (11.6%)	0.12
Renal insufficiency	134 (29.2%)	98 (41.9%)	36 (16.0%)	<0.001
Active cancer	110 (24.0%)	45 (19.2%)	65 (28.9%)	0.015
Symptoms at presentation				
Dyspnoea	359 (78.6%), n=457	190 (81.9%), n=232	169 (75.1%)	0.08
Syncope	53 (11.6%), n=457	39 (16.8%), n=232	14 (6.2%)	<0.001
Altered mental status	19 (4.1%), n=458	11 (4.7%), n=233	8 (3.6%)	0.53
Clinical findings at presentation				
Tachycardia	135 (30.4%), n=444	92 (40.7%), n=226	43 (19.7%), n=218	<0.001
Hypoxaemia	74 (19.8%), n=374	64 (32.3%), n=198	10 (5.7%), n=176	<0.001
Laboratory and imaging markers				
hsTnT ≥14 pg/ml	259 (60.2%), n=430	196 (92.9%), n=211	63 (28.8%), n=219	<0.001
NT-proBNP ≥600 pg/ml	212 (47.5%), n=446	168 (75.3%), n=223	44 (19.7%), n=223	<0.001
RV/LV diameter ratio ≥1.0 on CTPA	270 (65.4%), n=413	163 (79.5%), n=205	107 (51.4%), n=208	<0.001
Risk stratification				
modified FAST score ≥3 points ^a	115 (25.1%)	113 (48.3%)	2 (0.9%)	<0.001
ESC 2019 algorithm ^a				<0.001
Low risk	58 (12.6%)	0 (0.0%)	58 (25.8%)	
Intermediate-low risk	265 (57.7%)	98 (41.9%)	167 (74.2%)	
Intermediate-high risk	136 (29.6%)	136 (58.1%)	0 (0.0%)	
Outcome				
In-hospital adverse outcome	22 (4.8%)	19 (8.1%)	3 (1.3%)	<0.001
Catecholamine administration	15 (3.3%)	13 (5.6%)	2 (0.9%)	0.005
Cardiopulmonary resuscitation	5 (1.1%)	5 (2.1%)	0 (0.0%)	0.027
PE-related death	9 (2.0%)	8 (3.4%)	1 (0.4%)	0.022
In-hospital all-cause mortality	14 (3.1%)	11 (4.7%)	3 (1.3%)	0.036

^a calculation of scores based on hsTnI measurements

Abbreviations: HsTnI denotes high-sensitivity troponin I; hsTnT, high-sensitivity troponin T; NT-proBNP, N-terminal pro-brain natriuretic peptide; RV/LV, right/left ventricle; CTPA, computed tomography pulmonary angiography; ESC, European society of cardiology; PE, pulmonary embolism.

Table 2. Predictors of in-hospital adverse outcome and all-cause mortality

	All patients	In-hospital adverse outcome			In-hospital all-cause mortality		
	n=459	Yes (n=22)	No (n=437)	OR (95% CI)	Yes (n=14)	No (n=445)	OR (95% CI)
Age ≥75 years	155 (33.8%)	11 (50.0%)	144 (33.0%)	2.03 (0.86-4.8)	7 (50.0%)	148 (33.3%)	2.01 (0.69-5.83)
Sex (female)	241 (52.5%)	11 (50.0%)	230 (52.6%)	0.90 (0.38-2.12)	9 (64.3%)	232 (52.1%)	1.65 (0.55-5.01)
Comorbidities							
Chronic heart failure	59 (12.9%)	4 (18.2%)	55 (12.6%)	1.54 (0.50-4.73)	3 (21.4%)	56 (12.6%)	1.89 (0.51-7.00)
Coronary artery disease	67 (14.6%)	4 (18.2%)	63 (14.4%)	1.32 (0.43-4.03)	3 (21.4%)	64 (14.4%)	1.62 (0.44-5.98)
Chronic pulmonary disease	60 (13.1%), n=458	3 (14.3%), n=21	57 (13.0%)	1.11 (0.32-3.89)	4 (30.8%), n=13	56 (12.6%)	3.09 (0.92-10.36)
Diabetes mellitus	65 (14.2%)	4 (18.2%)	61 (14.0%)	1.37 (0.45-4.18)	3 (21.4%)	62 (13.9%)	1.68 (0.46-6.21)
Renal insufficiency	134 (29.2%)	14 (63.6%)	120 (27.5%)	4.62 (1.89-11.30)	8 (57.1%)	126 (28.3%)	3.38 (1.15-9.92)
Active cancer	110 (24.0%)	5 (22.7%)	105 (24.0%)	0.93 (0.34-2.58)	7 (50.0%)	103 (23.1%)	3.32 (1.14-9.69)
Symptoms at presentation							
Dyspnoea	359 (78.6%), n=457	17 (81.0%), n=21	342 (78.4%), n=436	1.17 (0.38-3.55)	11 (84.6%), n=13	348 (78.4%), n=444	1.52 (0.33-6.96)
Syncope	53 (11.6%), n=457	6 (28.6%), n=21	47 (10.8%), n=436	3.31 (1.23-8.95)	3 (23.1%), n=13	50 (11.3%), n=444	2.36 (0.63-8.88)
Altered mental status	19 (4.1%), n=45	5 (23.8%), n=21	14 (3.2%)	9.44 (3.03-29.42)	3 (23.1%), n=13	16 (3.6%)	8.04 (2.02-32.08)
Clinical findings at presentation							
Tachycardia	135 (30.4%), n=444	12 (57.1%), n=21	123 (29.1%), n=423	3.25 (1.34-7.91)	6 (46.2%), n=13	129 (29.9%), n=431	2.01 (0.66-6.09)
Hypoxaemia	74 (19.8%), n=374	7 (33.3%), n=21	67 (19.0%), n=353	2.13 (0.83-5.49)	3 (23.1%), n=13	71 (19.7%), n=361	1.23 (0.33-4.57)
Laboratory and imaging markers							
hsTnI >10 pg/ml	279 (60.8%)	20 (90.9%)	259 (59.3%)	6.87 (1.59-29.77)	12 (85.7%)	267 (60.0%)	4.00 (0.89-18.09)
hsTnI >16 pg/ml	234 (51.0%)	19 (86.4%)	215 (49.2%)	6.54 (1.91-22.42)	11 (78.6%)	223 (50.1%)	3.65 (1.00-13.26)
hsTnI >27 pg/ml	191 (41.6%)	15 (68.2%)	176 (40.3%)	3.18 (1.27-7.95)	9 (64.3%)	182 (40.9%)	2.60 (0.86-7.89)
hsTnI >16 pg/ml (women) / >34 pg/ml (men)	209 (45.5%)	17 (77.3%)	192 (43.9%)	4.34 (1.57-11.97)	11 (78.6%)	198 (44.5%)	4.57 (1.26-16.62)
hsTnI >100 pg/ml	108 (23.5%)	10 (45.5%) ⁹	98 (22.4%)	2.89 (1.21-6.87)	6 (42.9%)	102 (22.9%)	2.52 (0.86-7.44)
hsTnT ≥14 pg/ml	271 (61.0%), n=444	20 (95.2%), n=21	251 (59.3%), n=423	13.71 (1.82-103.08)	12 (85.7%)	259 (60.2%), n=430	3.96 (0.88-17.92)
NT-proBNP ≥600 pg/ml	212 (47.5%), n=446	18 (85.7%), n=21	194 (45.6%), n=425	7.14 (2.07-24.62)	12 (85.7%)	200 (46.3%), n=432	6.96 (1.54-31.47)
RV/LV diameter ratio ≥1.0 on CTPA	270 (65.4%), n=413	11 (64.7%), n=17	259 (65.4%), n=396	0.97 (0.35-2.68)	8 (80.0%), n=10	262 (65.0%), n=403	2.15 (0.45-10.27)

Abbreviations: OR denotes odds ratio; CI, confidence interval; hsTnI, high-sensitivity troponin I; hsTnT, high-sensitivity troponin T; NT-proBNP, N-terminal pro-brain natriuretic peptide; RV/LV, right/left ventricle; CTPA, computed tomography pulmonary angiography.

Table 3. Prognostic performance of different troponin cut-off values

Cut-off value	Prevalence	In-hospital adverse outcome	Sensitivity (95% CI)	Specificity (95% CI)	PPV (95% CI)	NPV (95% CI)	LR+	OR (95% CI)	In-hospital all-cause mortality	Sensitivity (95% CI)	Specificity (95% CI)	PPV (95% CI)	NPV (95% CI)	LR+	OR (95% CI)
hsTnI >10 pg/ml (lower limit of detection)	60.8%	7.2%	90.9% (72.2-97.5)	40.7% (36.2-45.4)	7.2% (4.7-10.8)	98.9% (96.0-99.7)	1.5	6.87 (1.59-29.77)	4.3%	85.7% (60.1-96)	40.0% (35.6-44.6)	4.3% (2.5-7.4)	98.9% (96.0-99.7)	1.4	4.00 (0.89-18.09)
hsTnI >16 pg/ml (optimal cut-off value)	51.0%	8.1%	86.4% (64.0-96.4)	50.8% (46.0-55.6)	8.1% (5.3-12.3)	98.7% (96.2-99.5)	1.8	6.54 (1.91-22.42)	4.7%	78.6 (48.9-94.3)	49.9% (45.2-54.6)	4.7% (2.6-8.2)	98.7% (96.2-99.5)	1.6	3.65 (1.01-13.26)
hsTnI >27 pg/ml (99 th percentile URL)	41.6%	7.9%	68.2% (47.3-83.6)	59.7% (55.1-64.2)	7.9% (4.8-12.6)	97.4% (94.7-98.7)	1.7	3.18 (1.27-7.95)	4.7%	64.3% (38.8-83.7)	59.1% (54.5-63.6)	4.7% (2.5-8.7)	98.1% (95.7-99.2)	1.6	2.60 (0.86-7.89)
hsTnI >16 pg/ml (women), >34 pg/ml (men) (sex-specific 99 th percentile URL)	45.5%	8.1%	77.3% (54.2-91.3)	56.1% (51.3-60.8)	8.1% (5.1-12.6)	98.0% (95.4-99.1)	1.8	4.34 (1.57-11.97)	5.3%	78.6 (48.8-94.3)	55.5% (50.7-60.2)	5.3% (3.0-9.2)	98.8% (96.5-99.6)	1.8	4.57 (1.26-16.62)
hsTnI >100 pg/ml (cut-off value proposed by [12])	23.5%	9.3%	45.5% (26.9-65.3)	77.6% (73.4-81.2)	9.3% (5.1-16.2)	96.6% (94.1-98)	2.0	2.89 (1.21-6.87)	5.6%	42.9% (21.4-67.4)	77.1% (73-80.7)	5.6% (2.6-11.6)	97.7% (95.6-98.8)	1.9	2.52 (0.86-7.44)
hsTnT ≥14 pg/ml	61.0%	7.4%	95.2% (74.1-99.8)	40.7% (36.0-45.6)	7.4% (4.8-11.1)	99.4% (96.8-99.9)	1.6	13.71 (1.82-103.08)	4.4%	85.7% (56.2-97.5)	39.8% (35.1-44.6)	4.4% (2.6-7.6)	98.8% (95.9-99.7)	1.4	3.96 (0.88-17.92)

Abbreviations: CI denotes confidence interval; PPV, positive predictive value; NPV, negative predictive value; LR+, positive likelihood ratio; OR, odds ratio. hsTnI, high-sensitivity troponin I; URL, upper reference limit; hsTnT, high-sensitivity troponin T.


Table 4. Comparison of risk stratification algorithms calculated based on hsTnI and hsTnT

Risk stratification algorithm	Troponin assay and cut-off value	Prevalence	In-hospital adverse outcome	Sensitivity (95% CI)	Specificity (95% CI)	PPV (95% CI)	NPV (95% CI)	LR+	OR (95% CI)	NRI
modified FAST score ≥ 3 points	hsTnT ≥ 14 pg/ml	21.8%	12.0%	54.5% (32.7-74.9)	79.9% (75.7-83.5)	12.0% (7.0-19.8)	97.2% (94.9-98.5)	2.7	4.76 (1.99-11.37)	0.06 (SE 0.09), p=0.51
	hsTnI >16 pg/ml	25.1%	12.2%	63.6% (40.8-82.0)	76.9% (72.6-80.7)	12.2% (7.4-19.4)	97.7% (95.5-98.8)	2.8	5.82 (2.38-14.27)	
ESC 2019 intermediate-high-risk*	hsTnT ≥ 14 pg/ml	32.2%	8.8%	59.1% (36.7-78.5)	69.1% (64.5-73.4)	8.8% (5.2-14.4)	97.1% (94.6-98.5)	1.9	3.23 (1.35-7.74)	0.03 (SE 0.07), p=0.68
	hsTnI >16 pg/ml	29.6%	9.6%	59.1% (36.7-78.5)	71.9% (67.3-76.0)	9.6% (5.7-15.7)	97.2% (94.8-98.5)	2.1	3.69 (1.54-8.85)	

* compared to ESC 2019 low / intermediate-low risk

Abbreviations: HsTnI denotes high-sensitivity troponin I; hsTnT, high-sensitivity troponin T; CI, confidence interval; PPV, positive predictive value; NPV, negative predictive value; LR+, positive likelihood ratio; OR, odds ratio; NRI, net reclassification improvement; SE, standard error; ESC, European Society of Cardiology.

Figure 1. Receiver operating characteristics curve of hsTnI concentrations with regard to an adverse in-hospital outcome



Bullet points indicate different investigated hsTnI cut-off values.

Abbreviations: HsTnI denotes high-sensitivity troponin I.

Supplementary Material

Supplementary Tables

Table s1. Comorbidities, results from risk stratification and outcomes of study patients stratified according to the availability of hsTnI measurements

	All patients (n=742)	Patients with hsTnI measurements (n=459)	Patients without hsTnI measurements (n=283)	p-value
Age ≥75 years	249 (33.6%)	155 (33.8%)	94 (33.2%)	0.88
Sex (female)	388 (52.3%)	241 (52.5%)	147 (51.9%)	0.88
Comorbidities				
Chronic heart failure	105 (14.2%)	59 (12.9%)	46 (16.3%)	0.20
Coronary artery disease	121 (16.3%)	67 (14.6%)	54 (19.1%)	0.11
Chronic pulmonary disease	110 (14.8%), n=741	60 (13.1%), n=458	50 (17.7%)	0.09
Diabetes mellitus	115 (15.5%)	65 (14.2%)	50 (17.7%)	0.20
Renal insufficiency	222 (30.5%), n=729	134 (29.2%)	88 (32.6%), n=270	0.34
Active cancer	155 (20.9%)	110 (24.0%)	45 (15.9%)	0.009
Symptoms at presentation				
Dyspnoea	589 (79.7%), n=739	359 (78.6%), n=457	230 (81.6%), n=282	0.32
Syncope	90 (12.2%), n=740	53 (11.6%), n=457	37 (13.1%)	0.55
Altered mental status	34 (4.6%), n=741	19 (4.1%), n=458	15 (5.3%)	0.47
Clinical findings at presentation				
Tachycardia	236 (32.8%), n=719	135 (30.4%), n=444	101 (36.7%), n=275	0.08
Hypoxemia	134 (21.9%), n=612	74 (19.8%), n=374	60 (25.2%), n=238	0.11
Laboratory and imaging markers				
hsTnT ≥14 pg/ml	433 (63.2%), n=685	271 (61.0%), n=444	162 (67.2%), n=241	0.11
NT-proBNP ≥600 pg/ml	326 (49.9%), n=653	212 (47.5%), n=446	114 (55.1%), n=207	0.07
RV/LV diameter ratio ≥1.0 on CTPA	443 (67.4%), n=657	270 (65.4%), n=413	173 (70.9%), n=244	0.14
Risk stratification				
modified FAST score ≥3 points ^a	182 (24.5%)	100 (21.8%)	82 (29.0%)	0.027
ESC 2019 algorithm ^a				0.24
Low risk	90 (12.1%)	56 (12.2%)	34 (12.0%)	
Intermediate-low risk	396 (53.4%)	255 (55.6%)	141 (49.8%)	
Intermediate-high risk	256 (34.5%)	148 (32.2%)	108 (38.2%)	
Outcome				
In-hospital adverse outcome	42 (5.7%)	22 (4.8%)	20 (7.1%)	0.19
Catecholamine administration	31 (4.2%)	15 (3.3%)	16 (5.7%)	0.11
Cardiopulmonary resuscitation	14 (1.9%)	5 (1.1%)	9 (3.2%)	0.042
PE-related death	16 (2.2%)	9 (2.0%)	7 (2.5%)	0.64
In-hospital all-cause mortality	29 (3.9%)	14 (3.1%)	15 (5.3%)	0.12

^a calculation of scores based on hsTnI measurements

Abbreviations: HsTnI denotes high-sensitivity troponin I; hsTnT, high-sensitivity troponin T; NT-proBNP, N-terminal pro-brain natriuretic peptide; RV/LV, right/left ventricle; CTPA, computed tomography pulmonary angiography; ESC, European Society of Cardiology; PE, pulmonary embolism.

Supplementary Figures

Figure s1: Rate of an in-hospital adverse outcome stratified according to hsTnl levels at presentation

