



Early View

Original article

Anxiety and depression in Dutch patients with Primary Ciliary Dyskinesia and their caregivers: associations with health-related quality of life

Marieke Verkleij, Iris Appelman, Josje Altenburg, Jos Twisk, Alexandra L. Quittner, Eric Haarman

Please cite this article as: Verkleij M, Appelman I, Altenburg J, *et al.* Anxiety and depression in Dutch patients with Primary Ciliary Dyskinesia and their caregivers: associations with health-related quality of life. *ERJ Open Res* 2021; in press (<https://doi.org/10.1183/23120541.00274-2021>).

This manuscript has recently been accepted for publication in the *ERJ Open Research*. It is published here in its accepted form prior to copyediting and typesetting by our production team. After these production processes are complete and the authors have approved the resulting proofs, the article will move to the latest issue of the ERJOR online.

Anxiety and depression in Dutch patients with Primary Ciliary Dyskinesia and their caregivers: Associations with health-related quality of life

Marieke Verkleij¹, PhD, Iris Appelman¹, MSc, Josje Altenburg², MD, PhD, Jos Twisk³, PhD,
Alexandra L. Quittner⁴, PhD, Eric Haarman⁵, MD, PhD

¹ Department of Pediatric Psychology, Amsterdam UMC, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands

² Department of Pulmonology, Amsterdam UMC, Amsterdam, the Netherlands

³ Department of Epidemiology and Data Science, Amsterdam UMC, Amsterdam, The Netherlands

⁴ Behavioral Health Systems Research, Miami, United States

⁵ Department of Pediatric Pulmonology, Emma Children's Hospital, Amsterdam UMC, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands

Corresponding author: Marieke Verkleij, PhD. Amsterdam UMC, location VU University Medical Center, Department of Pediatric Psychology, Reception L, PO Box 7057, 1007 MB Amsterdam, the Netherlands. E-mail: m.verkleij@amsterdamumc.nl

Take home message:

Individuals with PCD and their caregivers are at risk for developing symptoms of anxiety and depression, which are associated with worse health-related quality of life. There is an urgent need for psychological care in the PCD populations, especially in adults with PCD and caregivers.

ABSTRACT

Background: Primary Ciliary Dyskinesia (PCD) might be a risk factor for the development of anxiety and depression. This study investigated the associations between anxiety, depression and health-related quality of life (HRQoL) in individuals with PCD and their caregivers.

Methods: Children, adolescents and adults with PCD and their caregivers were invited to participate in a mental health screening program. During regular yearly outpatient visits, measures of anxiety (GAD-7), depression (PHQ-9), HRQoL (QOL-PCD), lung function (FEV₁) and Body Mass Index (BMI) were collected and associations of anxiety, depression and HRQL were estimated.

Results: 103 individuals participated in the mental health screening program. Elevated levels of anxiety (scores ≥ 10 on GAD-7) were found in 6% of adults (n=33), 14% of children (n=7), 6% of adolescents (n=17) and 20% of caregivers (n=46, 52% mothers). Elevated depression levels (scores ≥ 10 on PHQ-9) were found in 18% of adults, 14% of children, 6% of adolescents and 11% of caregivers. Anxiety and depression were associated with scales on the QOL-PCD. Mothers reported higher anxiety scores than fathers (30% vs 9%, p=0.03). A strong negative relationship was found between depression in caregivers and Physical Functioning (QOL-PCD) of the child. Anxiety and depression were not significantly associated with anxiety/depression in their child.

Conclusion: This is the first study investigating anxiety and depression in individuals with PCD and their caregivers. Our results revealed elevated levels of anxiety and depression, which were associated with worse HRQoL. These results suggest the need for psychological support in PCD.

INTRODUCTION

Primary Ciliary Dyskinesia (PCD) is a chronic, genetic disease with a worldwide prevalence of 1 in 10000-20000 people (1). This is likely an underestimate given that making the diagnosis is highly complex and many individuals are incorrectly diagnosed (2). However, in some more or less genetically isolated and consanguineous populations like Volendam in the Netherlands, incidence can be as high as 1:400, with a carrier frequency of 1:10 (3). PCD is associated with upper and lower respiratory tract infections due to structural and functional abnormalities of respiratory cilia (4). Other clinical symptoms include situs inversus, neonatal respiratory distress and infertility (4, 5).

Individuals with a chronic, respiratory disease are at increased risk for developing anxiety and depression symptoms (6-8). These symptoms are associated with negative health outcomes, including poor treatment adherence, increased mortality, higher health-care utilization and costs, lower body mass index (BMI), decreased lung function (FEV₁) and impaired health-related quality of life (HRQoL) (9-14). In cystic fibrosis (CF) for example, a large-scale epidemiological study in 9 countries indicated that rates of elevated anxiety and depression were 2 to 3 times higher than those found in community samples (14). This led to the development of international mental health guidelines for CF, co-sponsored by the CF Foundation and the European Cystic Fibrosis Society (15, 16), which recommended annual mental health screening and treatment for people with CF and their caregivers. Moreover, caregivers of patients with chronic respiratory diseases also report higher rates of anxiety and depression than caregivers of healthy controls (14, 17-18).

Until now, research and clinical care in PCD has mainly focused on the genetics, the complexities of the diagnostic process and physical manifestations of the disease, with limited attention to psychosocial problems (19, 20). The few studies that have examined the psychosocial impact of PCD, have shown that these individuals report feelings of

embarrassment, isolation and anxiety about the future (21, 22). Children with PCD exhibited more behavioral problems than healthy controls, specifically internalizing problems (23), and mothers of children with PCD reported higher levels of parenting stress and general stress, compared to mothers of healthy children (23). Moreover, HRQoL in individuals with PCD was worse than healthy controls (24). Studies have also shown that both physical functioning and respiratory symptoms in adults with PCD decrease with age (21). Recently, a disease-specific instrument for measuring HRQoL in PCD patients was developed and validated, 'Quality of Life for Primary Ciliary Dyskinesia' (QOL-PCD) (19, 21, 25-26).

Given the limited literature on psychosocial adjustment in individuals with PCD and their caregivers, including the prevalence of psychological symptoms, such as anxiety and depression, and their associations with HRQoL, a biopsychosocial approach was taken in this study. Thus, anxiety and depression may affect disease severity via mechanisms associated with poor treatment adherence and self-care (e.g. airway clearance, unhealthy diet, smoking) (10, 27). Biological stress systems in the brain (eg., sympathetic nervous system) are often dysregulated in individuals experiencing anxiety and depression, which might lead to increases in physical symptoms (28). Evidence also supports this model in CF (29, 30).

Depression is associated with a chronic, low-grade inflammatory response and activation of cell-mediated immunity, as well as activation of the compensatory anti-inflammatory reflex system (31). Studies in several chronic conditions have shown that depression, independent of disease, causes an increase in inflammation (e.g., coronary artery disease, diabetes). In patients with chronic respiratory conditions, inflammation is associated with pulmonary exacerbations and worsening disease (31-34). Combined, these processes were expected to negatively impact HRQoL in people with PCD (35).

The psychological wellbeing of caregivers may also affect the psychological and physical health of children with PCD. Caregivers of children with chronic conditions report higher levels of anxiety and depression than caregivers of healthy children, which, in turn, has been associated with a higher risk of those children developing symptoms of anxiety and depression (14, 36, 37). Psychological well-being of the caregiver has also been associated with disease severity. Parents experiencing depression, for example, may have more difficulty organizing and assisting with treatments (29), decreasing adherence which could potentially lead to more severe disease (29, 38). On the other hand, more severe disease in the child has also been associated with higher rates of anxiety and depression in their caregivers (18).

To better inform the inclusion of psychological assessment and treatment into the multidisciplinary PCD care-team, this study aimed to evaluate the complex associations between symptoms of anxiety and depression and HRQoL in individuals with PCD and their caregivers. We hypothesized that anxiety and depression in children and adults with PCD and their caregivers is more prevalent than reported for community samples, and that elevated symptoms of anxiety and/or depression will be associated with worse HRQoL. Moreover, we hypothesized that children with PCD who report elevated anxiety and/or depression scores and worse HRQoL will be more likely to have caregivers who also reported elevated psychological symptomatology.

METHODS

Study Design

This was a cross-sectional study conducted at a single PCD center in the Netherlands. The medical ethics committee of the VU University Medical Center (VUmc), Amsterdam, the Netherlands, approved the study.

Participants

From April 2019 to June 2020, all children (6-11 yrs.), adolescents (12-17 yrs.) and adults (aged ≥ 18) with PCD, and caregivers of children with PCD (0-17 yrs.) from Amsterdam University Medical Centers (Amsterdam UMC) were invited to participate in the study. PCD-diagnosis was based on a combination of tests including: high-speed-videomicroscopy before and after cell culture, nasal NO, transmission electron-microscopy and genetics. Similar to the iPCD cohort study (39), patients in this study can be divided into three subgroups based on diagnostic tests:

1. Definite PCD: patients with a clinical phenotype consistent with PCD and a hallmark transmission electron microscopy (TEM) finding, and/or pathogenic biallelic PCD genetic mutations (40);
2. Probable PCD: clinical phenotype consistent with PCD and abnormal high-speed video microscopy findings and/or low nasal nitric oxide;
3. Clinical phenotypes consistent with PCD diagnosis in whom other possible diagnoses such as CF were excluded (none included in this study).

Adolescents and adults with PCD, as well as caregivers of children with PCD, provided written, informed consent. Participants were excluded if they were unable to read and complete the questionnaires.

Instruments

Demographic and clinical variables

Demographic variables, such as age, gender, education level, marital status, ethnic background, comorbid diseases and previous and current psychological support were obtained from patients and caregivers during their annual clinic visit (see Table 1) by a demographic questionnaire. Height and weight of the adults and children/adolescents with PCD were obtained from chart review to calculate their BMI. Spirometry values (lung function) were assessed using the Masterscreen (Jaeger®, CareFusion Corporation). Before assessing lung function, short or long acting β_2 -adrenergic agonists were stopped for 12 hours. Forced expiratory volume in one second (FEV₁) was obtained for all adults with PCD and for children aged ≥ 6 yrs. Normal values of FEV₁ range between 80% and 120%.

Measures

Generalized Anxiety Disorder 7-item (GAD-7, Dutch version (41)). The GAD-7 assesses anxiety symptoms. It consists of 7 items, each scored as 0 ('not at all'), 1 ('several days'), 2 ('more than half of the days') or 3 ('nearly every day'). Total scores range from 0 to 21; scores of 5, 10 and 15 points are used as cut-off points from a general population for mild, moderate, and severe anxiety, respectively. The GAD-7 has excellent psychometric properties, including strong internal consistency (Cronbach's $\alpha = .85$) and convergent validity with DSM-V diagnostic criteria. Using a cut-off score of 10 or higher, sensitivity was 0.89 and specificity 0.82 (41, 42). The minimal important difference score of 4 points has been

established for the GAD-7 (43). In this study, the moderate score of the GAD-7 (≥ 10) was used as the clinical cut-off.

Patient Health Questionnaire-9 (PHQ-9, Dutch version (44)). The PHQ-9 assesses depressive symptoms and consists of 9 items corresponding to DSM-V diagnostic criteria for major depression. It is one of the most widely used depression screeners, is free and available in all major world languages. It has demonstrated strong psychometric properties. Items are scored on a 4-point Likert scale: 0 ('not at all'), 1 ('several days'), 2 ('more than half of the days') or 3 ('nearly every day'). Scores range from 0 to 27, with higher scores indicating more severe depressive symptoms. Scores can be divided according to level of severity: 'minimal' (scores 0-4), 'mild' (scores 5-9), 'moderate' (scores 10-14), 'moderately severe' (scores 15-19) and 'severe' (scores 20-27) with cut-off scores determined from a general population. The PHQ-9 also assesses suicidal ideation (question #9) which can accompany more severe depressive symptoms. Studies have shown that using a threshold score of 10 or higher has a sensitivity of .85 and specificity of 0.89 (45). The minimal clinically important difference score of 5 points has been established (46). The moderate score of the PHQ-9 (≥ 10) was used as the clinical cut-off.

Scores on the GAD-7 and PHQ-9 were categorized twice; first, scores were categorized on the basis of severity as follows: 0 (scores ≤ 4 ; 'minimal'), 1 (scores 5-9; 'mild'), 2 (scores 10-14; 'moderate') or 3 (scores ≥ 15 'severe'). Then, all scores of ≥ 10 on the GAD-7 and PHQ-9 were considered clinically elevated (in accordance with the ICMH recommendations) and manually scored as '1' ('elevated'). All scores <10 were manually scored as 0 ('not elevated').

In children from 6-11 yrs., the GAD-7 and PHQ-9 were filled in by the parent caregiver assessing their child's behavior. Adolescents (12-17 yrs.) filled in the GAD-7 and PHQ-9 by themselves.

Quality of Life for Primary Ciliary Dyskinesia (QOL-PCD; Dutch version (25, 26)).

The QOL-PCD is a profile measure of HRQoL for people with PCD, assessing quality of life across several domains. Different versions of the QOL-PCD were developed for children (6-11 yrs.; 34 items, 7 scales), adolescents (12-17 yrs.; 35 items, 9 scales) and adults (aged ≥ 18 yrs.; 40 items, 10 scales). The 7 scales that were included in all three versions of the QOL-PCD were: Physical Functioning, Emotional Functional, Social Functioning, Treatment Burden, Upper Respiratory Symptoms, Lower Respiratory Symptoms, and Ear and Hearing Symptoms. Other scales for adolescents and adults included: Role Functioning and Vitality (adolescents and adult version) and Health Perceptions (adult version). The different versions of the QOL-PCD have all demonstrated good reliability and validity (19, 21).

Procedure

Following the CF mental health guidelines (15), the procedure was as follows:

1. Adults, children and adolescents with PCD and caregivers of children with PCD received the questionnaires (GAD-7, PHQ-9 and QOL-PCD) from the PCD nurse at their annual medical check-up. The questionnaires were either completed immediately and returned to the PCD nurse, or filled in at home and returned by mail.
2. Upon receipt of the questionnaires, the psychologist immediately checked the PHQ-9 for suicidal ideation. If this item was endorsed, the psychologist determined the severity by using the Columbia Suicide Severity Rating Scale (C-

SSRS). If indicated, a referral to the psychiatrist in attendance at the hospital was made.

3. The other measures were scored within a week.
4. Psychologists called the participants to discuss the results of the GAD-7, PHQ-9 and QOL-PCD, and a referral plan to a healthcare provider was developed if needed.

Statistical analysis

All statistical analyses were performed with IBM SPSS Statistics, version 26 for Windows. P-values of $<.05$ (two-sided) were considered statistically significant. For missing values on the QOL-PCD, the median value of the corresponding scale was entered.

Descriptive statistics were used to evaluate and summarize the demographic and clinical characteristics of the participants as well as the mean anxiety, depression and HRQoL-scores. One-way ANOVA's were used to examine differences between the three groups of adults, adolescents and children with PCD and independent t-tests were used to examine differences in caregivers' in scores of anxiety, depression, and HRQoL.

Regression analyses were used to analyze the univariable associations between anxiety, depression, and HRQoL across the three groups. Mixed model analyses were used to estimate the associations between anxiety, depression and HRQoL in children with anxiety and depression in their caregivers. Mixed model analysis was used to take into account the dependency of the observations for two caregivers with one child. All regression analyses were adjusted for age, gender, BMI and lung function.

Results

Participant characteristics

In total, 43 adults and 45 children/adolescents with a confirmed diagnosis of PCD were invited to participate in the study with at least one primary caregiver. Total inclusion rate of participants with PCD was 66%. Ten adults and 19 children/adolescents with PCD did not respond or did not return the questionnaires; 2 children were unable to read in Dutch (2%). This resulted in a total of 103 participants: 33 (of 43) adults with PCD, 77%, 24 (of 43) children/adolescents with PCD (7 children 6-11 yrs. and 17 adolescents 12-17 yrs.), 56%, and 46 caregivers (24 mothers and 22 fathers of children/adolescents with PCD). Of the 24 children/adolescents with PCD, all mothers and fathers participated (2 adolescents had no father). Based on diagnostic tests performed 26 out of 33 adults (79%) and 24 out of 27 children/adolescents (89%) were classified as definite PCD. The remaining patients were classified as probable PCD.

Table 1 presents the demographic characteristics of the adults and children with PCD and their caregivers. Mean lung function of adults and children with PCD (6-11 yrs.) was below average (< 80%); mean BMI of adults with PCD was considered healthy (between 18.5 – 24.9 kg/m²).

In the demographic questionnaire, 21% of adults with PCD filled in that they are currently receiving or have received psychological support in the past year, and 15% need psychological support. In children with PCD (6-17 yrs.), 46% are currently receiving or have received psychological support in the past year, 15% need psychological support. Thirty-four % mothers and 28% fathers are currently receiving or have received psychological support in

the past year, 17% of mothers emphasized the need for psychological support and 5% of fathers need psychological support.

Table 1. Demographic characteristics of adults and children with PCD and parent caregivers (n = 103)

	Adults with PCD (n = 33)	Children with PCD (age 6-17 yrs.; n = 24)	Mothers (n = 24)	Fathers (n = 22)
Female total group, n (%)	20 (61)	14 (58)		
Mean age, (SD), range (yrs.)	38.3 (18.0), 18-77 yrs.	13.3 (3.3) 7-17 yrs.	42.8 (6.1), 32-55 yrs.	44.7 (5.2), 34-54 yrs.
Lung function (FEV ₁ %pred)	72.8 (21.2)	78.7 (13.4)		
BMI	24.2 (4.2)	20.1 (4.4)		
Received psychological support in past year, n (%)	5 (15)	7 (29)	4 (17)	5 (23)
Current receiving psychological support, n (%)	2 (6)	4 (17)	4 (17)	1 (5)
Need for psychological support ^a , n (%)	5 (15)	3 (15)	4 (17)	1 (5)
Relational status, n				
Alone	9		0	0
Married/living together	19		23	20
Divorced	0		1	2
Widowed	1		0	0
Other	1			
Ethnic background, n				
Dutch	27		22	21

Turkish	1	1	1
Other	1	1	0
Education level ^b , n			
Low	10	3	1
Middle	3	7	12
High	14	13	9
Currently working, n	22	22	22
Other chronic illness, n	9	6	5

Note. FEV₁ = Forced Expiratory Volume in one second, expressed as percentage of the predicted value; BMI = Body Mass Index.

^a This group also includes people who had already received psychological support and still need it.

^b 'Low' educational level: primary school, primary vocational education; 'Middle': intermediate general secondary education, secondary vocational education; 'High': higher general secondary education, higher vocational education, university education.

Prevalence of anxiety and depression

We hypothesized that anxiety and depression in individuals with PCD and their caregivers would be more prevalent than among community samples. Table 2 shows the classification of anxiety and depression scores in adults and children/adolescents with PCD and caregivers. Clinically elevated symptoms of anxiety were reported by 6% of adults, 14 % of children and 6% of adolescents with PCD and 30% of mothers and 9% of fathers. Clinically elevated symptoms of depression were reported by 18% of adults, 14% of children and 6% of adolescents with PCD, 12% of mothers and 9% of fathers.

Additional analyses computed for adolescents with PCD and their caregivers showed that 6% of adolescents scored in the elevated range (on either GAD-7 or PHQ-9), as did 29% of mothers and 14% of fathers.

Table 2. Classification of anxiety and depression scores in adults, children and adolescents with PCD and caregivers (n = 103)

	Adults with PCD (n = 33)	Children with PCD (age 6- 11 yrs., n = 7)	Adolescents with PCD (age 12-17 yrs., n = 17)	Mothers (n = 24)	Fathers (n = 22)
<i>Anxiety (GAD-7), n (%)</i>					
Normal range (score ≤ 4)	18 (55)	6 (86)	5 (29)	12 (50)	17 (77)
Mild range (score 5-9)	13 (39)	0	11 (65)	5 (21)	3 (14)
Moderate range (score 10-14)	2 (6)	1 (14)	1 (6)	3 (13)	2 (9)
Severe range (score ≥ 15)	0	0	0	4 (17)	0
<i>Depression (PHQ-9), n (%)</i>					
Normal range (score ≤ 4)	16 (49)	4 (57)	7 (41)	12 (50)	15 (68)
Mild range (score 5-9)	11 (33)	2 (29)	9 (53)	9 (38)	5 (23)
Moderate range (score 10-14)	5 (15)	1 (14)	0	1 (4)	2 (9)
Severe range (score ≥ 15)	1 (3)	0	1 (6)	2 (8)	0

Note. GAD-7 = Generalized Anxiety Disorder 7-Item; PHQ-9 = Patient Health Questionnaire-

Anxiety, depression and HRQoL

Table 3 shows mean anxiety, depression and HRQoL scores of adults, adolescents and children with PCD and caregivers. No significant differences were found in mean anxiety and depression scores between adults, adolescents and children with PCD. Mothers reported more symptoms of anxiety than fathers ($p = .03$).

The HRQoL domains Physical Functioning, Social Functioning, Treatment Burden, Role Functioning and Symptoms are worse with increasing age.

Table 3. Mean scores (SD) of anxiety (GAD-7), depression (PHQ-9) and health-related quality of life (QOL-PCD), in adults, adolescents and children with PCD and caregivers (n = 103)

	Adults with PCD (n = 33)		Adolescents with PCD (age 12-17 yrs., n = 17)		Children with PCD (age 6-11 yrs., n = 7)		Mothers (n = 24)		Fathers (n = 22)	
	M	(SD)	M	(SD)	M	(SD)	M	(SD)	M	(SD)
Anxiety (GAD-7)^a	4.3	(3.1)	5.4	(2.9)	3.0	(4.4)	6.2	(6.2)	3.0	(3.3)
Depression (PHQ-9)^b	5.6	(4.2)	5.5	(3.5)	3.6	(4.2)	5.4	(4.7)	3.1	(3.8)
Health-related quality of life (QOL-PCD)										
<i>Domains</i>										
Physical Functioning	65.3	(29.2)	87.5	(15.0)	84.8	(14.8)				
Emotional Functioning	81.9	(13.1)	73.4	(7.6)	76.2	(13.1)				
Social Functioning	34.7	(27.0)	62.5	(19.7)	91.4	(10.7)				
Treatment Burden	60.9	(23.2)	74.8	(22.4)	80.0	(26.1)				

Role Functioning	66.2	(25.4)	86.8	(15.3)		
Vitality	55.9	(19.3)	60.4	(14.6)		
Health Perceptions	50.8	(22.6)				
Upper Respiratory Symptoms	55.1	(21.6)	69.6	(19.3)	75.2	(17.5)
Lower Respiratory Symptoms	51.3	(18.7)	58.9	(19.1)	73.0	(10.4)
Hearing Symptoms	68.2	(32.9)	83.9	(18.1)	86.9	(10.6)

Note. GAD-7 = Generalized Anxiety Disorder 7-Item; PHQ-9 = Patient Health Questionnaire-9; QOL-PCD = Quality of Life Instrument for Primary Ciliary Dyskinesia.

^aScores (0-21), scores ≥ 10 are considered clinically elevated. ^bScores (0-27), scores ≥ 10 are considered clinically elevated.

As hypothesized, elevated symptoms of anxiety and/or depression were associated with worse HRQoL. In adults, anxiety was significantly associated with a worse score on the QOL-PCD domain Emotional Functioning. Depression was significantly associated with worse scores on the QOL-PCD domains Role Functioning and Vitality (Table 4). In adolescents, higher depression scores were significantly associated with worse scores on the QOL-PCD domains Physical Functioning and Vitality. In children (6-11 yrs.), anxiety and depression were not significantly associated with their scores on any of the QOL-PCD domains (Table 5).

Table 4. Unstandardized regression coefficients, confidence interval and p-values regarding the associations between anxiety, depression and health-related quality of life in adults (n=33) with PCD

	Adults with PCD (n = 33)	
	Anxiety (GAD-7)	Depression (PHQ-9)
Health-related quality of life		
(QOL-PCD) Domains		
Physical Functioning	-2.03 (-6.83 to 2.77) p=0.39	-1.74 (-5.18 to 1.71) p=0.31
Emotional Functioning	-2.95 (-5.21 to -0.70) p=0.01	-0.84 (-1.70 to 1.54) p=0.92
Social Functioning	3.93 (-2.56 to 10.43) p=0.22	-3.93 (-8.60 to 0.73) p=0.10
Treatment Burden	4.37 (-2.89 to 11.62) p=0.22	-4.12 (-8.97 to 0.73) p=0.09
Role Functioning	-0.96 (-5.14 to 3.24) p=0.64	-3.55 (-6.55 to -0.54) p=0.02
Vitality	-1.18 (-4.42 to 2.05) p=0.46	-2.35 (-4.67 to -0.02) p=0.05
Health Perceptions	-3.11 (-6.93 to 0.71) p=0.11	-1.53 (-4.10 to 1.04) p=0.23
Upper Respiratory Symptoms	0.59 (-3.74 to 4.91) p=0.78	-2.67(-5.78 to 0.43) p=0.09
Lower Respiratory Symptoms	-1.01 (-4.95 to 2.92) p=0.60	-0.75 (-3.57 to 2.08) p=0.59
Hearing Symptoms	2.05 (-4.76 to 8.86) p=0.54	-3.29 (-8.17 to 1.61) p=0.18

Note GAD-7 = Generalized Anxiety Disorder 7-Item; PHQ-9 = Patient Health Questionnaire-9; QOL-PCD = Quality of Life Instrument for Primary Ciliary Dyskinesia.

Table 5. Unstandardized regression coefficients, confidence interval and p-values regarding the associations between anxiety, depression and health-related quality of life in adolescents and children with PCD

	Adolescents with PCD (12-17 yrs.), n = 17		Children with PCD (6-11 yrs.), n = 7	
	Anxiety (GAD-7)	Depression (PHQ-9)	Anxiety (GAD-7)	Depression (PHQ-9)
Health related				
quality of life (QOL-				
PCD) Domains				
Physical Functioning	2.55 (-1.47 to 6.57) p=0.19	-4.79 (-8.16 to -1.42) p=0.01	-2.36 (-9.80 to 5.08) p=0.31	0.11 (-8.92 to 9.15) p=0.96
Emotional Functioning	-0.93 (-4.63 to 4.45) p=0.96	-0.23 (-4.04 to 3.58) p=0.90	-2.91 (-6.69 to 0.87) p=0.08	0.71 (-3.88 to 5.30) p=0.57
Social Functioning	-2.95 (-14.62 to 8.73) p=0.58	2.59 (-7.21 to 12.38) p=0.57	0.30 (-7.98 to 8.58) p=0.89	0.76 (-9.30 to 10.82) p=0.78
Treatment Burden	2.45 (-21.66 to 26.55) p=0.82	-3.27 (-23.07 to 16.53) p=0.71	-0.01 (-16.16 to 16.13) p=0.99	-3.58 (-23.18 to 16.03) p=0.52
Role Functioning	2.75 (-4.73 to 10.22) p=0.43	-3.78 (-10.05 to 2.49) p=0.21		
Vitality	3.21 (-2.29 to 8.71) p=0.22	-5.82 (-10.43 to -1.21) p=0.02		
Upper Respiratory Symptoms	3.28 (-6.04 to 12.61) p=0.45	-2.23 (-10.05 to 5.59) p=0.54	-2.30 (-15.92 to 11.32) p=0.54	-0.30 (-16.85 to 16.24) p=0.95
Lower Respiratory Symptoms	0.40 (-6.63 to 7.42) p=0.90	0.42 (-5.48 to 6.32) p=0.88	1.78 (-3.67 to 7.22) p=0.30	-3.40 (-10.02 to 3.22) p=0.16
Hearing Symptoms	-2.13 (-11.88 to 7.62) p=0.63	1.80 (-6.38 to 9.98) p=0.63	-1.43 (-10.51 to 7.64) p=0.57	0.54 (-10.48 to 11.56) p=0.85

Note GAD-7 = Generalized Anxiety Disorder 7-Item; PHQ-9 = Patient Health Questionnaire-9; QOL-PCD = Quality of Life Instrument for Primary Ciliary Dyskinesia.

Associations between caregivers and children

Thirdly, we hypothesized that children/adolescents with PCD who report elevated anxiety and/or depression scores and worse HRQoL will be more likely to have caregivers who also reported elevated symptoms of anxiety and/or depression. Table 6 shows the results of the mixed model analyses regarding the association between anxiety and depression in caregivers and HRQoL in their children with PCD. A strong, negative relationship was found between depression in caregivers and the QOL-PCD domain Physical Functioning in their child (Table 6). Parents who reported higher depressive symptoms had children /adolescents who reported worse physical functioning. All other relationships between anxiety and depression in caregivers of children/adolescents with PCD and HRQoL (domains of the QOL-PCD) of their children/adolescents were not statistically significant.

Table 6. Standardized regression coefficients and 95% confidence intervals (between brackets) regarding the associations between anxiety and depression in caregivers and health-related quality of life in their child/adolescent with PCD

Health related quality of life (QOL-PCD)		Anxiety (GAD-7)	Depression (PHQ-9)
Domains			
Physical Functioning	Crude	-0.12 (-0.46 to 0.22) p=0.48	-0.39 (-0.69 to -0.09) p=0.01
	Adjusted	-0.21 (-0.67 to 0.26) p=0.38	-0.50 (-0.82 to -0.18) p=0.002
Emotional Functioning	Crude	0.27 (-0.11 to 0.66) p=0.17	-0.13 (-0.49 to 0.23) p=0.47
	Adjusted	0.32 (-0.10 to 0.74) p=0.14	-0.15 (-0.51 to 0.20) p=0.40
Treatment Burden	Crude	0.24 (-0.13 to 0.61) p=0.21	-0.17 (-0.49 to 0.16) p=0.32
	Adjusted	0.25 (-0.13 to 0.63) p=0.20	-0.18 (-0.50 to 0.14) p=0.27
Social Functioning	Crude	0.05 (-0.31 to 0.41) p=0.80	0.10 (-0.21 to 0.41) p=0.53

	Adjusted	0.07 (-0.45 to 0.59) p=0.79	0.09 (-0.38 to 0.56) p=0.72
Upper Respiratory Symptoms	Crude	0.10 (-0.27 to 0.46) p=0.61	0.02 (-0.31 to 0.35) p=0.91
	Adjusted	0.21 (-0.37 to 0.78) p=0.49	0.15 (-0.29 to 0.60) p=0.49
Lower Respiratory Symptoms	Crude	0.07 (-0.30 to 0.44) p=0.71	0.08 (-0.25 to 0.42) p=0.63
	Adjusted	0.28 (-0.56 to 1.11) p=0.52	0.27 (-0.27 to 0.81) p=0.33
Hearing Symptoms	Crude	-0.12 (-0.44 to 0.19) p=0.45	-0.03 (-0.38 to 0.32) p=0.85
	Adjusted	-0.13 (-0.44 to 0.19) p=0.44	-0.08 (-0.43 to 0.28) p=0.68

Note Adjusted = adjusted for gender, age, FEV₁ and BMI.

GAD-7 = Generalized Anxiety Disorder 7-Item; PHQ-9 = Patient Health Questionnaire-9;

FEV₁ = Forced Expiratory Volume in one second, expressed as percentage of the predicted

value; BMI = Body Mass Index; QOL-PCD = Quality of Life Instrument for Primary Ciliary

Dyskinesia.

The results of the mixed model analyses regarding the association between anxiety and depression in children with PCD and anxiety and depression in their caregivers do not indicate significant associations (Table 7), although there was a positive relationship for depression, which trended toward significance (p=0.052 for the crude relationship). When adjusted for age, gender, BMI and lung function, this association did not reach statistical significance.

Table 7. Standardized regression coefficients and 95% confidence intervals (between brackets) regarding the association between anxiety and depression in children (n=24; independent variables) with PCD and anxiety and depression in their caregivers (n=46; outcome variables)

		Anxiety (GAD-7)	Depression (PHQ-9)
Anxiety (GAD-7)	Crude	.04 (-.26 to .34) p=0.79	
	Adjusted	.14 (-.22 to .49) p=0.45	
Depression (PHQ-9)	Crude	.28 (-.03 to .58) p=0.052	
	Adjusted	.24 (-.09 to .57) p=0.16	

Note Adjusted = adjusted for gender, age, FEV₁ and BMI

GAD-7 = Generalized Anxiety Disorder Scale 7-Item; PHQ-9 = Patient Health Questionnaire-9; FEV₁ = Forced Expiratory Volume in one second, expressed as percentage of the predicted value; BMI = Body Mass Index.

DISCUSSION

To our knowledge, this is the first study examining the prevalence of anxiety and depression, and its associations with HRQoL in people with PCD and parent caregivers. Results showed that elevated symptoms of anxiety and depression were prevalent in people with PCD and their caregivers. In particular, nearly half of the adults with PCD and half of the mothers of children/adolescents with PCD reported psychological symptoms in the mild to severe range. Anxiety and depression scores in children/adolescents and adults with PCD were associated with worse HRQoL, and depression in caregivers was associated with worse Physical Functioning (HRQoL) in their child/adolescent.

In accordance with our hypothesis, elevated symptoms of anxiety and depression were prevalent in children/adolescents and adults with PCD and caregivers. However, these scores differ from the CF population in a prior study of a sample of Dutch CF patients who are cared for at same academic medical hospital (VUmc) (16). Our study showed that 6% of adolescents with PCD reported elevated scores in comparison to 12% of adolescents with CF. The lower rates in PCD might be associated with the high burden of disease severity in adolescents with CF. Even though both conditions are chronic and complex, treatment of CF is more complex, time-consuming and burdensome than treatment of PCD (48). In addition, life expectancy is more severely reduced in CF. In contrast, social functioning appears to be negatively affected in both PCD and CF, possibly since individuals with both diseases report feeling stigmatized and/or socially isolated (22, 49).

Importantly, symptoms of anxiety and depression in parents of adolescents with PCD were higher than those reported for parents of adolescents with CF, using the same screening measures (16). Nearly 30% of mothers caring for adolescents with PCD reported elevated depression/anxiety compared to 13% of mothers of adolescents with CF and 14% of fathers caring for an adolescent with PCD compared to 9% of fathers of adolescents with CF.

The high rates of anxiety and depression documented in this study are in line with previous studies, suggesting that caregivers of chronically ill children experience more anxiety and depression symptoms than parents of healthy children, especially mothers (37, 49). In accordance with these studies, mothers of children/adolescents with PCD reported more symptoms of anxiety than fathers; 30% of mothers reported anxiety scores in the moderate to severe range, indicating that these mothers are at high risk. This may be due to the lack of psychological support embedded into the multidisciplinary PCD-team, the lack of well-organized patient organizations and foundations, and support among patients and families.

Anxiety and/or depression symptoms in children/adolescents and adults with PCD were negatively associated with worse HRQoL thereby confirming our hypothesis. The association of anxiety and depression with worse HRQoL is supported by previous studies examining other, chronic respiratory diseases (10, 12). Our results indicated that this is true of not just Physical Functioning and Vitality in those with PCD, but also affects Emotional and Role Functioning. Given that anxiety and depression were negatively related to HRQoL, measuring and treating these symptoms might serve as a preventive treatment. Future research should investigate whether identifying and treating psychological symptoms might reduce the negative impact on HRQoL.

Studies of CF and asthma (14, 38) have shown that anxiety and/or depression in caregivers were associated with similar elevation in their children/adolescents. In this study, results showed that depression in caregivers was associated with Physical Functioning of the child on the QOL-PCD. Due to our study design, causality cannot be established. However, there might be a bidirectional relationship (50), indicating that a “family systems” approach, involving both children/adolescents with PCD and their caregivers in clinical care. The importance of this approach is highlighted by previous studies indicating that worse emotional well-being in caregivers is negatively related to treatment adherence and disease severity in their children (38, 51). More adults reported elevated symptoms of depression (18%) than anxiety (6%). Depression is seen as far more harmful for disease management and health outcomes. In CF, depression has been associated with earlier mortality, more frequent exacerbations and lower lung function (30). Therefore, it is important to screen and treat depression at early stage.

Strengths and limitations should be considered. Screening measures such as the GAD-7 and PHQ-9 are not a full diagnostic clinical interview, and these screening measures are not sufficient for a diagnosis. However, both the GAD-7 and PHQ-9 have proven to be effective

and valid in screening anxiety and depression symptoms and are the recommended screening instruments according to the ICMH guidelines and most major international guidelines (e.g., NICE guidelines, USPTF) (15, 42, 45). A limitation of the study is that parents completed the GAD-7 and PHQ-9 for children under age 12, since there are very few well-established screeners for this age group.

Importantly, the results of our study may represent a first step towards more multidisciplinary clinical care of PCD. The use of a disease-specific instrument (QOL-PCD) provided a great deal of insight into the impact of PCD on daily functioning. Given the rare nature of this disease, a relatively large number of people with PCD and parent caregivers were included in this study. However, a larger sample size would increase statistical power of our analyses and improve their generalizability. Due to the cross-sectional design of the study, without multiple measurements over time, no assumptions can be made about the causal directions of these relationships. In the future, a longitudinal repeated measures design should investigate whether anxiety and depression are associated with other important health variables and outcomes, such as treatment adherence, mortality, health-care utilization and costs (10, 13).

Clinical implications

Our results showed that there is an urgent need for psychological care in the PCD populations, especially in adults with PCD and parents of children/adolescents with PCD. Psychological care is already embedded in CF care at many CF centers in the Netherlands, in which a psychologist/social worker is part of the multidisciplinary team. This model might work well in PCD too. In the CF mental health guidelines, annual mental health screening and treatment is recommended beginning at age 12 for adolescents with CF and for parents of children 0 through 17 years (15). National implementation of these guidelines has been highly successful (52), partly because of the development of a mental health “toolbox” and

additional mental health training sessions at national conferences. The CF ICMH guidelines can be implemented in clinical PCD care as well, if the appropriate mental health experts are integrated into the care team. Evaluating the possibilities of implementing these screening recommendations in clinical practice should be a focus of future research. Moreover, additional studies are needed to identify and address factors that might influence mental health, disease severity and HRQoL in individuals with PCD and their caregivers.

Acknowledgements

We thank the children, adolescents and adults with PCD and their caregivers for their cooperation, and the personnel for their help with recruitment.

REFERENCES

1. Mirra, V., Werner, C., & Santamaria, F. (2017). Primary ciliary dyskinesia: an update on clinical aspects, genetics, diagnosis, and future treatment strategies. *Front Pediatr.* 5: 135. doi: 10.3389/fped.2017.00135.
2. Behan, L. et al. (2016). Diagnosing primary ciliary dyskinesia: an international patient perspective. *Eur. Respir. J.* 48:1096–107. doi: 10.1183/13993003.02018-2015.
3. Onoufriadis, A., Paff, T., Antony, D., Shoemark, A., Micha, D., Kuyt, B., ...& Chung, E.M. (2013). Splice-site mutations in the axonemal outer dynein arm docking complex gene *CCDC114* cause primary ciliary dyskinesia. *Am J Hum Genet.* 92:88-98. doi: 10.1016/j.ajhg.2012.11.002.
4. Goutaki, M., Meier, A. B., Halbeisen, F. S., Lucas, J. S., Dell, S. D., Maurer, E., ... & Kuehni, C. E. (2016). Clinical manifestations in primary ciliary dyskinesia: systematic review and meta-analysis. *Eur Respir J.* 48:1081-1095. doi: 10.1183/13993003.00736-2016.
5. Behan, L., Dimitrov, B. D., Kuehni, C. E., Hogg, C., Carroll, M., Evans, H. J., ...& Lucas, J.S. (2016). PICADAR: a diagnostic predictive tool for primary ciliary dyskinesia. *Eur Respir J.* 47:1103-12.
6. Ryu, Y. J., Chun, E. M., Lee, J. H., & Chang, J. H. (2010). Prevalence of depression and anxiety in outpatients with chronic airway lung disease. *Korean J Intern Med.* 25:51-7. doi: 10.3904/kjim.2010.25.1.51.
7. Stoop, C. H., Nefs, G., Pommer, A. M., Pop, V. J. M., & Pouwer, F. (2015). Effectiveness of a stepped care intervention for anxiety and depression in people with diabetes, asthma or COPD in primary care: a randomized controlled trial. *J Affect Disord.* 184:269-76. doi: 10.1016/j.jad.2015.05.063.

8. Verkleij M, Beelen A, van Ewijk BE, Geenen R (2017). Multidisciplinary treatment in children with problematic severe asthma: A prospective evaluation. *Pediatr Pulmonol.* 52:588-97. doi: 10.1002/ppul.23623.
9. Janssen, D. J., Spruit, M. A., Leue, C., Gijzen, C., Hameleers, H., Schols, J. M., ... & Ciro network. (2010). Symptoms of anxiety and depression in COPD patients entering pulmonary rehabilitation. *Chron Respir Dis.* 7:147-57. doi: 10.1177/1479972310369285.
10. Knudsen, K. B., Pressler, T., Mortensen, L. H., Jarden, M., Skov, M., Quittner, A. L., ... & Boisen, K. A. (2016). Associations between adherence, depressive symptoms and health-related quality of life in young adults with cystic fibrosis. *Springerplus.* 5:1-8. doi: 10.1186/s40064-016-2862-5.
11. Lu, Y., Mak, K. K., Van Bever, H. P., Ng, T. P., Mak, A., & Ho, R. C. M. (2012). Prevalence of anxiety and depressive symptoms in adolescents with asthma: A meta-analysis and meta- regression. *Pediatr Allergy Immunol.* 23:707-15. doi: 10.1111/pai.12000.
12. Riekert, K. A., Bartlett, S. J., Boyle, M. P., Krishnan, J. A., & Rand, C. S. (2007). The association between depression, lung function, and health-related quality of life among adults with cystic fibrosis. *Chest.* 132:231-7. doi: 10.1378/chest.06-2474.
13. Panagioti, M., Scott, C., Blakemore, A., & Coventry, P. A. (2014). Overview of the prevalence, impact, and management of depression and anxiety in chronic obstructive pulmonary disease. *Int J Chron Obstruc Pulmon Dis.* 9:1289-306. doi: 10.2147/COPD.S72073.
14. Quittner, A. L., Goldbeck, L., Abbott, J., Duff, A., Lambrecht, P., Solé, A., ... & Blackwell, L. (2014). Prevalence of depression and anxiety in patients with cystic

- fibrosis and parent caregivers: results of The International Depression Epidemiological Study across nine countries. *Thorax*. 69:1090-7. doi: 10.1136/thoraxjnl-2014-205983.
15. Quittner, A.L., Abbott, J., Georgiopoulos, A.M., Goldbeck, L., Smith, B., Hempstead, S.E... & Elborn, S. (2016). International Committee on Mental Health in Cystic Fibrosis: Cystic Fibrosis Foundation and European Cystic Fibrosis Society consensus statements for screening and treating depression and anxiety. *Thorax*.71:26-34. <https://doi.org/10.1136/thoraxjnl-2015-207488>.
16. Verkleij, M., de Winter, D., Hurley, M. A., & Abbott, J. (2018). Implementing the International Committee on Mental Health in Cystic Fibrosis (ICMH) guidelines: Screening accuracy and referral-treatment pathways. *J Cyst Fibros*. 17: 821-27. doi: 10.1016/j.jcf.2018.02.005.
17. Besier, T., & Goldbeck, L. (2011). Anxiety and depression in adolescents with CF and their caregivers. *J Cyst Fibros*. 10:435-42. doi: 10.1016/j.jcf.2011.06.012.
18. Easter, G., Sharpe, L., & J. Hunt, C. (2015). Systematic review and meta-analysis of anxious and depressive symptoms in caregivers of children with asthma. *J Pediatr Psychol*. 40:623-32. doi: 10.1093/jpepsy/jsv012.
19. Behan, L., Leigh, M. W., Dell, S. D., Quittner, A. L., Hogg, C., & Lucas, J. S. (2019). Validation of pediatric health- related quality of life instruments for primary ciliary dyskinesia (QOL- PCD). *Pediatr Pulmonol*. 54:2011-20. doi: 10.1002/ppul.24507.
20. Daniels, M. L. A., & Noone, P. G. (2015). Genetics, diagnosis and future treatment strategies for primary ciliary dyskinesia. *Expert Opin Orphan Drugs*. 3:31-44. doi: 10.1517/21678707.2015.989212.
21. Behan, L., Leigh, M. W., Dell, S. D., Galvin, A. D., Quittner, A. L., & Lucas, J. S. (2017). Validation of a health-related quality of life instrument for primary ciliary dyskinesia (QOL-PCD). *Thorax*. 72:832-9. doi: 10.1136/thoraxjnl-2016-209356.

22. Whalley, S., & McManus, I. C. (2006). Living with primary ciliary dyskinesia: a prospective qualitative study of knowledge sharing, symptom concealment, embarrassment, mistrust, and stigma. *BMC Pulm Med.* 6: 25. doi: 10.1186/1471-2466-6-25.
23. Carotenuto, M., Esposito, M., Di Pasquale, F., De Stefano, S., & Santamaria, F. (2013). Psychological, cognitive and maternal stress assessment in children with primary ciliary dyskinesia. *World J Pediatr.* 9:312-7. doi: 10.1007/s12519-013-0441-1.
24. Pifferi, M., Bush, A., Di Cicco, M., Pradal, U., Ragazzo, V., Macchia, P., & Boner, A. L. (2010). Health-related quality of life and unmet needs in patients with primary ciliary dyskinesia. *Eur Resp J.* 35:787-94. doi: 10.1183/09031936.00051509.
25. Lucas, J. S., Behan, L., Galvin, A. D., Alpern, A., Morris, A. M., Carroll, M. P., ... & Quittner, A. L. (2015). A quality-of-life measure for adults with primary ciliary dyskinesia: QOL-PCD. *Eur Respir J.* 46:375-83. doi: 10.1183/09031936.00216214.
26. Dell, S. D., Leigh, M. W., Lucas, J. S., Ferkol, T. W., Knowles, M. R., Alpern, A., ... & Quittner, A. L. (2016). Primary ciliary dyskinesia: first health-related quality-of-life measures for pediatric patients. *Ann Am Thorac Soc.* 13:1726-35. doi: 10.1513/AnnalsATS.201603-198OC.
27. Lin, E. H., Katon, W., Von Korff, M., Rutter, C., Simon, G. E., Oliver, M., ... & Young, B. (2004). Relationship of depression and diabetes self-care, medication adherence, and preventive care. *Diabetes Care.* 27: 2154-60. doi: 10.2337/diacare.27.9.2154.
28. Katon, W., Lin, E. H., & Kroenke, K. (2007). The association of depression and anxiety with medical symptom burden in patients with chronic medical illness. *Gen Hosp Psychiatry.* 29:147-55. doi: 10.1016/j.genhosppsy.2006.11.005.

29. Barker, D.H., & Quittner, A.L. (2016). Parental depression and pancreatic enzymes adherences in children with cystic fibrosis. *Pediatrics*. 137:e20152296. doi: 0.1542/peds.2015-2296.
30. Schechter, M.S., Ostrenga, J.S., Fink, A.K., Barker, D.H., Sawicki, G.S., & Quittner, A.L. (2021). Decreased survival in cystic fibrosis patients with a positive screen for depression. *J Cyst Fibros*. 20:120-6. doi: 10.1016/j.jcf.2020.07.020.
31. Berk, M., Williams, L. J., Jacka, F.N., O'Neil, A., Pasco, J. A., Moylan, S....& Maes, M. (2013). So depression is an inflammatory disease, but where does the inflammation come from? *BMC Med*. 11:200. doi: 10.1186/1741-7015-11-200.
32. Amodeo, G., Trusso, M.A., & Fagiolini, A. (2017). Disentangling a clear yet complex and multi-faceted link. *Neuropsychiatry*. 7:448-57.
33. Felger, J.C. & Lotrich, F.E. (2013). Inflammatory cytokines in depression: neurobiological mechanisms and therapeutic implications. *Neuroscience*. 246:199–229. doi: 10.1016/j.neuroscience.2013.04.060.
34. Hoen, P., Kupper, N., & de Jonge P.(2012). Depression and cardiovascular disease progression: epidemiology, mechanisms and treatment. *Stress and cardiovascular disease*. Springer 211-233 London, United Kingdom.
35. Pumar, M.I., Gray, C.R., Walsh, J.R., Yang, I.A., Rolls, T.A., Ward, D.L. (2014). Anxiety and depression-Important psychological comorbidities of COPD. *J Thorac Dis*. 6:1615-31. doi: 10.3978/j.issn.2072-1439.2014.09.28.
36. Graziano, S., Spanò, B., Majo, F., Righelli, D., Vincenzina, L., Quittner A., & Tabarini P. (2020). Rates of depression and anxiety in Italian patients with cystic fibrosis and parent caregivers: Implementation of the Mental Health Guidelines. *Respir Med*. 172:106147. doi: 10.1016/j.rmed.2020.106147.

37. van Oers, H. A., Haverman, L., Limperg, P. F., van Dijk-Lokkart, E. M., Maurice-Stam, H., & Grootenhuis, M. A. (2014). Anxiety and depression in mothers and fathers of a chronically ill child. *Matern Child Health J.* 18:1993-2002.
38. Verkleij, M., Van de Griendt, E. J., Colland, V., Van Loey, N., Beelen, A., & Geenen, R. (2015). Parenting stress related to behavioral problems and disease severity in children with problematic severe asthma. *J Clin Psychol Med Settings.* 22:179-93. doi: 10.1007/s10880-015-9423-x.
39. Halbeisen, F.S., Goutaki, M., Spycher, B.D., Amirav, I., Behan, L., Boon, M.,...& Kuehni, C.E. (2018). Lung function in patients with primary ciliary dyskinesia: an iPCD Cohort study. *Eur Respir J.* 52:1801040. doi: 10.1183/13993003.01040-2018.
40. Lucas, J.S., Barbato, A., Collins, S.A., et al. (2017). European Respiratory Society guidelines for the diagnosis of primary ciliary dyskinesia. *Eur Respir J.* 49:1601090. doi: 10.1183/13993003.01090-2016.
41. Spitzer, R. L., Kroenke, K., Williams, J. B., & Löwe, B. (2006). A brief measure for assessing generalized anxiety disorder: the GAD-7. *Arch Intern Med.* 166:1092-7. doi: 10.1001/archinte.166.10.1092.
42. Hinz, A., Klein, A. M., Brähler, E., Glaesmer, H., Luck, T., Riedel-Heller, S. G., ... & Hilbert, A. (2017). Psychometric evaluation of the Generalized Anxiety Disorder Screener GAD-7, based on a large German general population sample. *J Affect Disord.* 210:338-44. doi: 10.1016/j.jad.2016.12.012.
43. Toussaint, A., Hüsing, P., Gumz, A., Wingenfeld, K., Härter, M., Schramm, E., & Löwe B. (2020). Sensitivity to change and minimal clinically important difference of the 7-item Generalized Anxiety Disorder Questionnaire (GAD-7). *J Affect Disord.* 265:395-401. doi:10.1016/j.jad.2020.01.032.
44. Kroenke, K., Spitzer, R. L., & Williams, J. B. (2001). The PHQ- 9: validity of a brief

depression severity measure. *J Gen Intern Med.* 1:606-13. doi: 10.1046/j.1525-1497.2001.016009606.x.

45. Löwe, B., Unützer, J., Callahan, C.M., Perkins, A.J., & Kroenke, K. (2004). Monitoring depression treatment outcomes with the patient health questionnaire-9. *Med Care.* 42:1194-201. doi: 10.1097/00005650-200412000-00006.
46. Manea, L., Gilbody, S., & McMillan, D. (2012). Optimal cut-off score for diagnosing depression with the Patient Health Questionnaire (PHQ-9): a meta-analysis. *CMAJ.* 184:E191-6. doi: 10.1503/cmaj.110829.
47. Keniş Coşkun, Ö., Gençer Atalay, K., Erdem, E., Karadağ- Saygi, E., Gökdemir, Y., & Karadağ, B. (2019). Caregiver burden in children with cystic fibrosis and primary ciliary dyskinesia. *Pediatr Pulmonol.* 54:1936-40. doi: 10.1002/ppul.24503.
48. Sawicki, G. S., Heller, K. S., Demars, N., & Robinson, W. M. (2015). Motivating adherence among adolescents with cystic fibrosis: youth and parent perspectives. *Pediatr Pulmonol.* 50:127-36. doi: 10.1002/ppul.23017.
49. Besier, T., Born, A., Henrich, G., Hinz, A., Quittner, A. L., Goldbeck, L., & TIDES Study Group. (2011). Anxiety, depression, and life satisfaction in parents caring for children with cystic fibrosis. *Pediatr Pulmonol.* 46:672-82. doi: 10.1002/ppul.21423.
50. Lavigne, J. V., Hopkins, J., Gouze, K. R., & Bryant, F. B. (2015). Bidirectional influences of anxiety and depression in young children. *J Abnorm Child Psychol.* 43:163-76. doi: 10.1007/s10802-014-9884-7.
51. Smith, B. A., Modi, A. C., Quittner, A. L., & Wood, B. L. (2010). Depressive symptoms in children with cystic fibrosis and parents and its effects on adherence to airway clearance. *Pediatr Pulmonol.* 45:756-63. doi: 10.1002/ppul.21238.

52. Quittner, A.L., Abbott, J., Hussain, S., Ong, T., Uluer, A., Hempstead, S., ...& Smith, B. (2020). Integration of mental health screening and treatment into cystic fibrosis clinics: Evaluation of initial implementation in 84 programs across the United States. *Pediatr Pulmonol.* 5:2995-3004. doi: 10.1002/ppul.24949.