



Early View

Research letter

Observational study of inhaled corticosteroid treatment for improved expiratory variability index in steroid-naive asthmatic children

Janne Burman, Pekka Malmberg, Ville-Pekka Seppä, Tuomas Jartti, Sami Remes, Ove Mickelsson, Anna S. Pelkonen, Mika J Mäkelä

Please cite this article as: Burman J, Malmberg P, Seppä V-P, *et al.* Observational study of inhaled corticosteroid treatment for improved expiratory variability index in steroid-naive asthmatic children. *ERJ Open Res* 2021; in press (<https://doi.org/10.1183/23120541.00499-2021>).

This manuscript has recently been accepted for publication in the *ERJ Open Research*. It is published here in its accepted form prior to copyediting and typesetting by our production team. After these production processes are complete and the authors have approved the resulting proofs, the article will move to the latest issue of the ERJOR online.

Copyright ©The authors 2021. This version is distributed under the terms of the Creative Commons Attribution Non-Commercial Licence 4.0. For commercial reproduction rights and permissions contact permissions@ersnet.org

Research letter: Observational study of inhaled corticosteroid treatment for improved expiratory variability index in steroid-naive asthmatic children

Janne Burman¹, MD; Pekka Malmberg¹, MD, PhD; Ville-Pekka Seppä², PhD, Tuomas Jartti^{3,4}, MD, PhD; Sami Remes⁵, MD, PhD, Ove Mickelsson⁶, MD, Anna S Pelkonen¹, MD, PhD, Mika J Mäkelä¹, MD, PhD

¹ Helsinki University and University Hospital of Helsinki, Skin and Allergy Hospital, Department of Allergology, Helsinki, Finland

² Revenio Research Ltd., Vantaa, Finland

³ Department of Paediatrics and Adolescent Medicine, Turku University Hospital and University of Turku, Finland

⁴ Department of Paediatrics, Oulu University Hospital, Finland

⁵ Department of Paediatrics, Kuopio University Hospital, Finland

⁶ Mehiläinen Töölö, Helsinki, Finland

Running title: ICS improved EVI in asthmatic children

Correspondence details

Janne Burman

E-mail: janne.burman@hus.fi

Address: Helsinki University, University Hospital of Helsinki, Skin and Allergy Hospital,

P.O. Box 160, FIN 00029 HUS

E-mail: janne.burman@hus.fi

Word count: 1582 words

Figures and Tables: 1 figure

Number of references: 9

Online supplementary material: No

Conflict of interest: VPS holds patents relating to impedance pneumography (IP) and is an employee of Revenio Group corporation that commercializes IP technology. The other authors declare no conflicts of interest.

Funding Acknowledgements: The founders played no role in the analysis, study design, or data collection. The corresponding author had final responsibility for the decision to submit for publication.

Financial support: This work was supported by Revenio Research Ltd. and research grants from the Allergy Research Foundation, Jalmari and Rauha Ahokas Foundation (Helsinki), the Helsinki Region Allergy Foundation (Helsinki), the Culture Foundation, the Sigrid Juselius Foundation (Helsinki), all in Finland.

Keywords: asthma, children's health, impedance pneumography, inhaled corticosteroid treatment, lung function test

Abbreviations:

C-ACT: Childhood Asthma Control Test; EVI: expiratory variability index; ICS; inhaled corticosteroid; IP: impedance pneumography

Research letter: Inhaled corticosteroid treatment for improved expiratory variability index in steroid-naive asthmatic children

To the Editor:

The Lancet Asthma Commission calls for objective measurements to support asthma diagnosis, especially for children[1]. The expiratory variability index (EVI) is a novel method of testing lung function testing that can be measured passively at night with an impedance pneumography (IP) technique using four skin electrodes and a small body-worn recorder. The aim of EVI is to measure variation in flow-volume curve shapes within 15-45% of the expired volume. It has been demonstrated that normal variations in the curve shapes are mainly caused by comparing different sleep cycles [2], which are reduced in the presence of bronchial obstructions [3].

IP is feasible for young children, and EVI values have previously been associated with the severity of wheezing episodes in children aged 1-5 years [4,5]. In previous studies, up to 94% of recordings were successfully made, and EVI values in young children exhibiting wheezing decreased after inhaled steroids (ICS) were withdrawn [5]. In infants, EVI is associated with lung function and the risk of developing asthma [6]. Our aim was to determine whether ICS treatment improves EVI in children aged 4-7 years with doctor-diagnosed asthma.

This study was conducted in a tertiary care hospital (Skin and Allergy Hospital) and a private clinic (Töölö, Mehiläinen), with children monitored for six months after starting ICS-treatment. Inclusion criteria were age (4-7 years), doctor-diagnosed asthma and a clinical need to start ICS-treatment. Exclusion criteria included the use of ICS in the past 30 days, acute respiratory infection in the previous 14 days, preterm birth, another chronic lung disease, or the use of other

active medical equipment. In total, 53 children participated in the study, and 21 were excluded from the analysis due to drop out (n=4) or missing EVI values at any visit (n=17). The feasibility analysis was performed on 49 children who took part during every study visit.

After Visit I, every child started ICS treatment. Visit II occurred within 43 days (interquartile range 40–48 days) after Visit I, and Visit III was within 68 days (interquartile range 61–71) of visit II. Impulse oscillometry (Jaeger GmbH, Würzburg, Germany) and bronchodilatation and/or free running tests and the Childhood Asthma Control Test (C-ACT) were performed during every visit. Exercise-induced bronchoconstriction was defined as having a $\geq 40\%$ increase in resistance in 5 Hz after exercise. After each visit, IP registration (Ventica, Revenio Research Ltd., Finland) was performed for two nights and average values were used in the analysis. The EVI was measured by software (Ventica Analytics v2.1.0) that calculates Pearson's correlations among the tidal breathing flow-volume curves of 15%–45% of exhaled volume during sleep. The EVI value is represented by the formula $[\log_{10}[\text{IQR}(r)] + 2] \times 10$, in which $\text{IQR}(r)$ is the interquartile range of the correlations. A low EVI indicates reduced variability in tidal breathing flow-volume curves. Any EVI values below 14 were considered abnormal, and the reference values are based on children aged 1–5 years [4,5].

The Helsinki University Hospital of Children and Adolescents Ethics Committee approved the study and each participant (including parents) provided written informed consent. The study was registered at Clinicaltrials.gov (NCT03377192).

SPSS version 22 (IBM Corp, Armonk, NY, USA) was used for statistical analysis, and Friedman two-way analysis and the Wilcoxon signed rank test adopted for analysing continuous parametric data at different time points. For categorical data, the chi-square and Fisher exact tests

(when counts were <5) were employed. In the correlation analysis, Spearman's correlations were used. The statistical significance was established at $p < 0.05$.

The median age of the 32 children in the final analysis was 5.1 years (interquartile range 4.6–5.8 years), including 21 (66%) boys. In total, 19 (59%) had doctor-diagnosed wheezing episodes, 12 (38%) had histories of hospital ward treatment, 20 (63%) had atopic eczema, 23 (72%) were sensitized to aeroallergens, and 24 (75%) had positive Asthma Predictive Index. Thirteen children had no history of doctor-diagnosed wheezing episodes but long-term asthmatic symptoms, and 11 of them were confirmed to have EIB through a free running test

After their baseline visit, most children 27/32 (84%) started twice daily use of Flixotide evohaler (Glaxo Wellcome Production; Evreux, France), which provides 125 ug fluticasone per dose. The remaining five children started twice a day a budesonide dry powder inhalator (Budesonid Easyhaler, Orion Pharma; Espoo, Finland)—two at 100 ug and three at 200 ug per dose.

After a second visit, one child was switched from the Flixotide Evohaler to a formoterol and budesonide aerosol (Symbicort 4,5/160 ug 1x2, AstraZeneca AB; Södertelje, Sweden), and one was switched from budesonide to a formoterol and budesonide powder inhaler (Bufomix Easyhaler, Orion Pharma; Espoo, Finland). Montelukast was added for treatment of two children at a dose of 4–5 mg/day.

Five (16%) of the children had EVI values <14 at baseline; two (6.3%) had these values at Visit II, and one (3.1%) at Visit III. After ICS, the median EVI-values improved significantly from the baseline visit (Median 17.5 [15.2–18.9] vs 18.4 [17.3–20.0] vs 17.9 [17.1–19.6],

p=0.028). Further, significant improvements were observed between Visit I and II and between Visits I and III. The individual results are presented in Figure 1.

During follow-up, one child required emergency evaluation due to wheezing the day after the baseline visit. This child exhibited an abnormal EVI-value (10.1) that night. At baseline visit 12 hours before the emergency visit, lung auscultation was normal, and no obstruction was observed in impulse oscillometry. However, the child experienced significant exercise-induced bronchoconstriction in the free running test and measured 18 points in the C-ACT. At Visit II (48 days after baseline) the child exhibited a normal 23 points for the C-ACT, significant bronchoconstriction was not observed in the free running test, and the EVI value improved significantly to 20.7.

Overnight IP registration was successful 219/294 (74%) times. Further, there were no differences between centres, and success was often achieved at Visit II (70/98 [71%] vs 84/98 [86%] vs 66/98 [67%], p=0.008). The primary reason for failed registrations was a batch of defective electrodes. Minor problems included a lack of registration time (due to short sleeping times) and electrodes that disconnected during sleep. Children who failed both registrations were more often boys (16/17 [94%] vs 21/32 [66%], p=0.037) and older children (5.9 [4.8–7.1] vs 5.1 [4.6–5.8] years, p=0.045) compared to 32 children who successfully registered at least once every visit.

After ICS treatment, the C-ACT results improved progressively from Visit I to visit II and III (Median 21 vs 23.5 vs 24, p<0.001) and acute physician visits due to respiratory symptoms decreased compared to six months before and after ICS began (Median (range) 0 (0-3) vs 0 (0-1), p=0.017). Further, exercise-induced bronchoconstriction alleviated (Median 61% vs 16.5% vs

13.%, $p < 0.001$), but there were no significant changes in bronchodilatation test (Median -21% vs -21% vs -16%, $p = 0.12$) or baseline lung function defined as resistance at 5 Hz (Median 0.06SD vs -0.55SD vs -0.69SD, $p = 0.054$).

The correlation analysis revealed no significant correlations between EVI values at baseline and oscillometry results at 5 Hz ($r_s = -0.038$, $p = 0.835$) or reactance at 5 Hz ($r_s = 0.217$, $p = 0.267$). On one hand, no significant correlation was found between EVI and C-ACT score ($r_s = -0.085$, $p = 0.645$). On the other hand, there were no correlations between C-ACT scores at the baseline visit and impulse oscillometry resistance at 5 Hz ($r_s = 0.033$, $p = 0.949$) or reactance at 5 Hz ($r_s = -0.054$, $p = 0.783$).

In previous studies, EVI decreased among wheezy hospitalized children aged 1-5 years after ICS ceased, and improved significantly immediately after exacerbation during treatment in hospital [4,5]. In our study of children with relative mild asthma, concordant with C-ACT scores, significant improvements were observed after starting ICS treatment in older preschool children without recent acute exacerbation. According to our findings, EVI also appears to be associated with asthma control in older children, up to seven years of age.

Previously, 77%-94% of registrations were successful [4,5,7]. However, a defective batch of electrodes was the primary reason for the lower rate in our study. Accordingly, 17 out of 49 (35%) were excluded from the final analysis. To ensure successful results, we recommend that registration should be performed for two nights.

No significant correlations were observed between EVI and impulse oscillometry, baseline parameters or C-ACT scores. However, two other studies have found only weak correlations

between asthma test scores and both fractional exhaled nitric oxide and forced expiratory volume in the first second of a spirometry test [8,9]

The strength of our study was that patients were monitored for 6 months. The relatively mild patient population could also be considered a strength, because for those children the diagnosis of asthma is clinically challenging and lung function tests would be required especially for this patient group. A major shortcoming of the study was the significant proportion of excluded children. Moreover, this was an observational study that did not include a healthy control group, and children older than 5 years have no reference values for EVI. In this study, most children already had EVI values >14 at Visit I and the proportion of participants with an EVI value below 14 did not differ significantly from that among healthy children aged under 5 years, which is considered normal in children aged 1-5 years. This high proportion of normal test results could be related to a mild disease, or inappropriate reference values for older children.

In conclusion, for children aged 4-7 years with mild asthma, EVI improved significantly after starting ICS. Future studies are needed to explore optimal EVI cut-off values and clinical utility in children in within this age group.

REFERENCES

1. Bush A, Pavord ID. The Lancet Asthma Commission: Towards the Abolition of Asthma? *EMJ*. 2018;3:10-15.
2. Hult A, Juraski RG, Gracia-Tabuenca J, Partinen M, et al. Sources of variability in expiratory flow profiles during sleep in healthy young children. *Respir Physiol Neurobiol*. 2020;274:103352. doi: 10.1016/j.resp.2019.103352.
3. Seppä VP, Hult A, Gracia-Tabuenca J, Paasilta M, et al. Airway obstruction is associated with reduced variability in specific parts of the tidal breathing flow–volume curve in young children. *ERJ Open Res*. 2019;17;5(2):00028-2019. doi: 10.1183/23120541.00028-2019.
4. Seppä VP, Turkalj M, Hult A, et al. Expiratory variability index (EVI) is associated with the severity of acute bronchial obstruction in small children: A proof-of-concept study. *Pediatr Allergy Immunol*, 2020;31:636-642.
5. Seppä VP, Paasilta M, Kivistö J, et al. Reduced expiratory variability index (EVI) is associated controller medication withdrawal and symptoms in wheezy aged 1-5 years. *Pediatr Allergy Immunol*. 2020;31:489-495.
6. Seppä VP, Gracia-Tabuenca J, Kotaniemi-Syrjänen, Malmström K, et al. Expiratory variability index is associated with asthma risk, wheeze and lung function in infants with recurrent respiratory symptoms. *ERJ Open Research* 2020;6:00167-2020. doi: 10.1183/23120541.00167-2020
7. Seppä VP, Pelkonen AS, Kotaniemi-Syrjänen A, et al. Tidal flow variability measured by impedance pneumography relates to childhood asthma risk. *Eur Respir J*. 2016;47:1687-1696.
8. Nguyen VN, Chavannes NH. Correlation between fractional exhaled nitric oxide and Asthma Control Test score and spirometry parameters in on-treatment-asthmatics in Ho Chi Minh City. *J Thorax Dis*. 2020;12:2197-2209.
9. Melosini L, Dente FL, Bacci E, et al. Asthma control test (ACT): comparison with clinical functional, and biological markers of asthma control. *J Asthma*. 2012;49:317-323.

Figure Legend:

FIGURE 1. Expiratory variability index at every visits for 32 children. In total 19 of 32 children lowest value were observed at Visit I before starting inhaled corticosteroid.

