



Early View

REVIEW

ERS International Congress 2021: Highlights from the Allied Respiratory Professionals assembly

Lucy Robertson, Filipa Machado, Sebastian Rutkowski, Liliana Silva, Sabina Miranda, Ingeborg Farver-Vestergaard, Thomas Janssens, Karl P. Sylvester, Chris Burtin, Andreja Šajnić, Joana Cruz

Please cite this article as: Robertson L, Machado F, Rutkowski S, *et al.* ERS International Congress 2021: Highlights from the Allied Respiratory Professionals assembly. *ERJ Open Res* 2022; in press (<https://doi.org/10.1183/23120541.00674-2021>).

This manuscript has recently been accepted for publication in the *ERJ Open Research*. It is published here in its accepted form prior to copyediting and typesetting by our production team. After these production processes are complete and the authors have approved the resulting proofs, the article will move to the latest issue of the ERJOR online.

Copyright ©The authors 2022. This version is distributed under the terms of the Creative Commons Attribution Non-Commercial Licence 4.0. For commercial reproduction rights and permissions contact permissions@ersnet.org

ERS International Congress 2021: Highlights from the Allied Respiratory Professionals assembly

Lucy Robertson^{1,*}, Filipa Machado^{2,3*}, Sebastian Rutkowski^{4,*}, Liliana Silva^{5,6,*}, Sabina Miranda^{7,*}, Ingeborg Farver-Vestergaard^{8,*}, Thomas Janssens^{9,*}, Karl P. Sylvester¹⁰, Chris Burtin³, Andreja Šajnić¹¹, Joana Cruz¹²

1 Respiratory Physiology, Royal Papworth Hospitals NHS Foundation Trusts, Cambridge, United Kingdom

2 Respiratory Research and Rehabilitation Laboratory (Lab3R), School of Health Sciences (ESSUA), University of Aveiro, Aveiro, Portugal

3 REVAL – Rehabilitation Research Center, BIOMED – Biomedical Research Institute, Faculty of Rehabilitation Sciences, Hasselt University, Diepenbeek, Belgium

4 Department of Physical Education and Physiotherapy, Opole University of Technology, Opole, Poland

5 Matosinhos Local Health Unit, Porto, Portugal;

6 CINTESIS – Centre for Health Technology and Services Research, Porto, Portugal

7 Pneumology Department, HUNSC Hospital, Tenerife, Spain

8 Department of Medicine, Vejle Hospital, Vejle, Denmark

9 Health Psychology, KU Leuven, Belgium

10 Respiratory Physiology, Royal Papworth & Cambridge University Hospitals NHS Foundation Trusts, Cambridge, United Kingdom

11 Department for Respiratory Diseases Jordanovac, University Hospital Center, Zagreb, Croatia

12 Center for Innovative Care and Health Technology (ciTechCare), School of Health Sciences (ESSLei), Polytechnic of Leiria, Leiria, Portugal

*The authors contributed equally

Correspondence:

Joana Cruz, joana.cruz@ipleiria.pt

Center for Innovative Care and Health Technology (ciTechCare), School of Health Sciences (ESSLei), Polytechnic of Leiria

Campus 2, Morro do Lena – Alto do Vieiro, Apartado 4163, 2411-901 Leiria, Portugal

Abstract

This paper provides an overview of some of the most memorable sessions that were (co)organised by the Allied Respiratory Professionals Assembly during the 2021 European Respiratory Society (ERS) International Congress, which was held online for the second consecutive year due to the COVID-19 pandemic. Early Career Members from Assembly 9 summarised the content of the sessions (three oral communication sessions, two symposia, and one experts view), with the support of the chairs from the four Assembly groups: Respiratory Function Technologists and Scientists (Group 9.01); Physiotherapists (Group 9.02); Nurses (Group 9.03); Psychologists and Behavioural Scientists (Group 9.04). The sessions covered the following topics: impact of COVID-19 on lung function and healthcare services and the importance of quality assurance and technology in lung function assessment; diagnosis and management of sarcopenia in patients with chronic respiratory disease; maintenance of the effects of pulmonary rehabilitation; solutions outside the hospital for the management of patients with COVID-19 in need of health care; the nursing perspective during the COVID-19 pandemic; and psychological and behavioural issues in respiratory care. This highlights paper provides valuable insight into the latest scientific data and emerging areas affecting clinical practice of Allied Respiratory Professionals.

Introduction

The Virtual 2021 European Respiratory Society (ERS) International Congress provided the Allied Respiratory Professionals with a wide range of outstanding sessions (co)organised by Assembly 9 including Symposia, Experts view, Skills lab, and 189 abstracts spread over oral communication and e-poster sessions.

In this paper, the four groups of Assembly 9 highlight some of the most memorable sessions of the ERS Congress, targeting delegates who were present in the virtual sessions as well as those unable to attend. Early Career Members of the Assembly (including respiratory function technologists and scientists, physiotherapists, nurses, and psychologists and behavioural scientists) were invited to summarise the latest scientific and clinical insights presented in the following sessions: three oral presentation sessions on the impact of COVID-19 on lung function and health services and quality assurance in lung function assessment, the nursing perspective during the COVID-19 pandemic, and psychological and behavioural issues in respiratory care; two symposia on the diagnosis and management of sarcopenia in patients with chronic respiratory disease and the maintenance of the pulmonary rehabilitation effects; and an experts view on the implementation of solutions outside the hospital for the management of patients with COVID-19 in need of health care. The next pages provide a summary of these sessions, along with important take-home messages, hoping to inspire readers to keep up to date in their areas of interest.

Group 9.01: Respiratory Function Technologists and Scientists

Oral Presentation: Impact of COVID-19, quality assurance and technology on pulmonary function

This oral presentation session (session ID 295) included communications on the impact of COVID-19 on lung function and healthcare services, and on the importance of quality assurance and technology to assess lung function.

Ventilation heterogeneity after COVID-19 (O. Savushkina, Russia)

O. Savushkina [1] presented findings from a cross-sectional study investigating the influence of COVID-19 on ventilation heterogeneity. The study utilised the multiple-breath nitrogen washout test in post-COVID subjects with no chronic respiratory disease and examined the relationship between the lung clearance index (LCI) and pulmonary function parameters. The study demonstrated that ventilation heterogeneity was detected in 85% cases 63 days after the onset

of COVID-19. Additionally, there were significant correlations between LCI, pulmonary function and impulse oscillometry parameters.

Evaluation of patient attendance to respiratory physiology in the midst of the COVID-19 Pandemic (S. Yearsley, United Kingdom)

S. Yearsley [2] presented, on behalf of C. Hurst, retrospective collated attendance data from two Respiratory Physiology Laboratories (North Midlands, UK) during the COVID-19 pandemic (1st January 2019 - 31st December 2020). The data demonstrated that the 'did not attend' (DNA) rate of 8.1% exceeded the 5.2% DNA rate from the Hospital Episode Statistics rate for outpatient appointments (April–October 2020). This equates to an approximate cost to the NHS of £261,171 [3]. Post-service resumption DNA rates increased (7.8% vs. 9.2%), but it was also observed that, with targeted communication, the DNA rate decreased 2.7-fold, which demonstrates the importance of verbal communication prior to appointments to improve outpatient attendance during the pandemic.

Implementation of the 2017 ERS Methacholine Challenge Standard (I. Steenbruggen, The Netherlands)

I. Steenbruggen [4] presented their study determining the percentage of patients with positive and negative methacholine challenge tests (MCT). The classification category for the positive tests additionally determined if there was a difference in provocative dose (PD20) between males and females. In total, 253 results were retrospectively reviewed from two laboratories. The results demonstrated the utility of MCT in either confirming or excluding airway hyperresponsiveness (AHR), with results presented in figure 1 showing the AHR classification according to the 2017 ERS Methacholine challenge standard [5]. Findings from this study also highlighted that men achieved significantly higher PD20 compared with women.

(insert Figure 1 about here)

Evaluation of the Global Lung function Initiative reference values for spirometry and diffusion capacity (K.D. Soomer, Belgium)

K.D. Soomer [6] presented results from a retrospective analysis evaluating the quality of the global lung initiative (GLI) reference values [7, 8] in an Antwerpen population with no known cardiac and respiratory disease history. Overall, it was demonstrated that the GLI reference values suitably predicted lung function. However, it was shown that the values of the Forced expiratory volume in one second/Forced vital capacity (FEV₁/FVC) ratio were lower than

predicted in tall subjects and D_{LCO} values in older female populations. These findings could result in an increase in suspected prevalence of obstructive and diffusion disorders in these groups.

Pulmonary airway resistance variability in normal biologic quality control subjects among pulmonary function testing systems across British Columbia (C.D. Mottram, United States of America)

C.D. Mottram [9] presented a study which investigated the variability of the panting and tidal breathing techniques when measuring airway resistance (R_{aw}) and/or specific conductance ($sGaw$) in biological quality control subjects (BioQC) amongst different pulmonary testing systems across British Columbia. R_{aw} and $sGaw$ data were analysed from 168 sessions over a 6-month period submitted to British Columbia's Diagnostic Accreditation Programme. Analysis of the data suggested that, in BioQC subjects, a $CV \leq 10\%$ would be an acceptable range for R_{aw} and $sGaw$ measurements. Findings also suggested that the panting technique showed the least degree of CV variability. This knowledge may be useful for laboratories implementing BioQC checks.

Do bacterial/viral filters impact quality assurance verification of ultrasonic spirometers? (L. Robertson, United Kingdom)

L. Robertson [10] presented results from a quality control study determining whether application of bacterial viral filters (BVF) impacted expiratory and inspiratory volume verification accuracy of Easy-on PC ultra-sonic spirometers at different flow rates. Findings highlighted that the proximal position, which is the usual site for BVF placement, significantly impacted the accuracy of expiratory volume verification at low flow rates, whereas with the BVF placed distally, the expiratory volume verification was achieved at all flow rates, but the inspiratory volume verification was not. From this work, it can be recommended that, if one wishes to apply a BVF on the Easy-On ultra-sonic spirometer to reduce cross infection risk, it should be placed distally only if expiratory volumes are of clinical importance and not inspiratory volumes.

Breath detection algorithms affect lung function outcomes (M. Oestreich, Switzerland)

M. Oestreich [11] presented a study about breath detection algorithms in children, as breath detection is essential in calculating outcomes in multiple breath washout (MBW). This is especially important in children where irregular breathing is frequent. The authors produced a novel algorithm for breath detection and compared it to algorithms used in MBW software from trials collected in children and adolescents. The results demonstrated that, in trials with normal

tidal breathing, there was no difference in outcomes between the software and the algorithm. However, greater challenges in breath detection were found in young children, as in 14% of trials in children aged 5 to 10 years the number of breaths detected differed by $\geq 10\%$ and the lung clearance index differed up to 40.5% (mean difference 3.6%). It was concluded that, as the MBW technique relies on cumulative analysis of all washout breaths, rejection of breaths should be limited. Findings demonstrated that the new breath detection algorithm, based on reversal and presence of CO₂, could be an alternative algorithm to those currently used.

Real-world application of spirometry quality control deep-learning algorithm (S. Stanojevic, Canada)

S. Stanojevic [12] described the results of a study that validated a recently developed spirometry quality control software based on a deep-learning-based algorithm [13]. Spirometry data were collected from a pre-school population and from a healthy adult population. Data were evaluated by the quality control software using 2005 ATS/ERS standards [14]. Manoeuvre acceptability and quality acceptability were compared with trained operators at time of testing. Overall, findings showed that the quality control software was in agreement with the technicians' review in the different populations. This study provides evidence for the clinical use of the software for reviewing quality of spirometry results, saving clinicians' time.

Use of artificial intelligence for improving quality control of oscillometry (C. Veneroni, Italy)

C. Veneroni [15] presented one study about the use of machine learning methods to reliably classify artefact free breaths in impulse oscillometry (IOS). This artefact detection is fundamental for situations when testing cannot be supervised. In total, 932 measurements were collected from 134 patients using a multifrequency stimulus at 5-11-19Hz following the ERS standards [16]. Results demonstrated that the feature selection performed by a wrapper method combined with an Adaboost tree model provided the best performance in classifying artefacts. Additionally, automatically classified tests differed from the manually classified ones by more than 0.5 cmH₂O*s/L in less than 5% of measurements. The authors concluded that supervised machine learning techniques are able to reliably detect IOS measurements and this would be of value in home monitoring settings where tests cannot be supervised.

Effects of respiratory muscle training on quality of life, physical function, and psychological state in post-COVID syndrome (T. C. Nuñez-Flores, Spain)

T. C. Nuñez-Flores [17] presented this late breaking work from a double-blind randomised controlled trial (RCT) involving post-COVID subjects, where the effects of high intensity home-

based respiratory muscle training over 8 weeks were studied on quality of life, physical and pulmonary function, psychological and cognitive state. The results showed that subjects in the experimental group improved quality of life in regard to breathlessness, muscle fatigue, respiratory muscle strength, cardiorespiratory endurance, anxiety and depression levels, posttraumatic stress disorder and cognition function when compared to the control group, but no between-group differences were observed in pulmonary function and peripheral muscle strength. This study demonstrates the benefits of home-based respiratory muscle training on quality of life and functional characteristics in people with post-COVID syndrome, where symptoms of breathlessness and fatigue are common.

Take-home messages

- To reduce cross-infection risk when using Easy-on ultra-sonic spirometer, the bacterial viral filter should be applied distally.
- Recently developed spirometry quality control software showed overall good agreement with technicians' review and, therefore, software could be used clinically to review spirometry data quality and save clinicians' time.
- A home-based respiratory muscle training programme can improve quality of life, symptoms, physical function, cognitive and psychological function of people with post-COVID syndrome in the short-term.

Group 9.02: Physiotherapists

Symposium: Sarcopenia in chronic respiratory disease: new definitions, assessment and treatment options

In this Symposium (session ID 210), four expert speakers provided an update on the latest knowledge on sarcopenia definition and diagnosis, its assessment and management, in the context of chronic respiratory diseases.

Update on recent definitions and diagnosis of sarcopenia (G. Bahat, Turkey)

G. Bahat started by explaining that sarcopenia was first clinically defined in 1998 solely by the presence of low muscle mass [18]. It was only in 2010 that a change in the paradigm happened, with the publication of several consensus reports integrating also decreased muscle function into the definition of sarcopenia [19-22]. Since the age-related decline in muscle strength begins before the decline in muscle mass, and the former is also 2-5 times faster [23], it is possible that by only including the presence of low muscle mass in the definition of sarcopenia, many cases

of older adults with reduced muscle strength would be overlooked. Therefore, low muscle strength has become the main characteristic of sarcopenia in the current consensus definitions (Table 1) [24-26]. G. Bahat concluded the talk by emphasising that future studies should include muscle strength in their sarcopenia definition.

(insert Table 1 about here)

Malnutrition and sarcopenia assessment – A. Schols (Netherlands)

A. Schols highlighted the importance of considering the different metabolic phenotypes since sarcopenia is observed in different metabolic syndromes, such as cachexia (characterised by an additional loss of body weight and fat mass) and sarcopenic obesity (characterised by adiposity) [27]. Different metabolic phenotypes result in different clinical risks (e.g., increased cardiovascular/mortality risk, impaired physical performance), which are stratified based on body composition and change in body weight [27]. For screening purposes, the SARC-F, the SARC-CalF or the sarcopenia index can be used [28]. Nevertheless, a thorough assessment of all sarcopenia domains, i.e., muscle mass (whole body magnetic resonance imaging, computed tomography, dual-energy X-ray absorptiometry, bioelectrical impedance, anthropometric measures), muscle strength (quadriceps strength, handgrip strength, chair stand test) and physical performance (gait speed, short physical performance battery, timed up and go test, 6-minute walk test) is recommended [28]. To capture malnutrition, additional measures such as the MUST and the SNAQ tools, information on dietary intake and nutrients deficiency are also important.

Underlying mechanisms and pharmacological approaches for treatment of sarcopenia (N. Greening, United Kingdom)

The third talk of the Symposium focused on skeletal muscle loss, particularly in chronic obstructive pulmonary disease (COPD). N. Greening explained that there are multiple triggers for sarcopenia in COPD and their underlying mechanisms are linked [29]. Disuse atrophy, hypoxia and inflammation are the most notable drivers of muscle loss [29]. To counteract this muscle loss, some anabolic therapies have been used, such as testosterone [30], bimagrumap [31] and selective androgen receptor modulator. Although these anabolic interventions result in increased muscle mass and some improvements in muscle strength, no improvements in the functional level (e.g., 6-minute walk test) of patients with COPD have been found [30, 31]. Nevertheless, it is possible that these anabolic interventions play a role in specific individuals and circumstances, such as during acute exacerbations of COPD, where it is crucial to prevent

the loss of muscle mass that occurs during hospitalisation [32]. N. Greening finished the talk highlighting that it would also be important to consider muscle endurance, neural adaptations and oxidative capacity when studying sarcopenia.

Modalities of pulmonary rehabilitation, including exercise training for sarcopenia in chronic lung disease (M. Maddocks, United Kingdom)

M. Maddocks emphasised that, overall, around 22% of the patients with COPD have sarcopenia [33]. The international clinical practice guidelines for sarcopenia strongly recommend the prescription of resistance training [34]; nevertheless, most of the evidence informing the guidelines comes from studies in non-sarcopenic patients. Moreover, there is wide variety in the exercise programmes delivered, thus, the exact way of designing an exercise programme for patients with sarcopenia remains to be determined [35]. Previous studies in patients with COPD have shown that: i) patients with sarcopenia respond well to pulmonary rehabilitation (PR) and 28% are even able to reverse it [36]; and ii) the combination of endurance or interval training with resistance training might be optimal in patients with sarcopenia [37]. Recently, electromyostimulation has also shown potential to improve muscle strength and function in sarcopenic patients [38]. M. Maddocks concluded the talk by drawing attention to the promising results of the “Sarcopenia and Physical frailty IN older people: multi-component Treatment strategies” (SPRINTT) trial [39], which will be published shortly.

Take-home messages

- Sarcopenia is defined by low muscle strength and low muscle quality/quantity, and it is essential that studies start assessing sarcopenia comprehensively, i.e., including muscle mass, muscle strength and physical performance measures.
- Anabolic interventions improve muscle mass in patients with COPD, but that does not translate into improvements in muscle function.
- Pulmonary rehabilitation is an evidence-based intervention for the management of sarcopenia in patients with chronic respiratory diseases.

Symposium: *Maintenance of the beneficial effects of pulmonary rehabilitation: the role of digital health*

This Symposium (session ID 348) featured presentations on the incorporation of new technologies to optimise maintenance of pulmonary rehabilitation (PR) outcomes, including a synthesis of the current state of knowledge and a patient's testimony.

Patient's perspective (J. M. Fourrier, France)

J.M. Fourrier described the experience as a patient with the use of new technologies for monitoring physical activity (PA). In his opinion, the most significant difficulties associated with activities of daily living are dyspnoea, pain and muscle fatigue, which can be managed by engaging in PA. J.M. Fourrier uses digital technology (Fitbit smartwatch) to help monitor daily exercise and health status. The user interface tracks the patient's achievement and enables him to connect with a coach who conducts group exercises in real time. J.M. Fourrier emphasised that he was not aware of any individuals dissatisfied with the rehabilitation programme; all patients experienced an improvement in quality of life. He concluded that exercise has become a natural part of his daily life, partly thanks to digital health technology, for which he is grateful.

Natural decline of (extra)pulmonary features in patients with COPD: a slippery slope (L.G.W. Vanfleteren, Sweden)

L.G.W. Vanfleteren outlined four areas of natural deterioration in patients with COPD: i) lung function decline; ii) exacerbations and the natural history of COPD; iii) alterations in body composition over time; and iv) comorbidities. With regards to disease exacerbations, there is a window of opportunity about 5 years after the first exacerbation, where PR programmes should be implemented. After this time, the risk of another exacerbation or even death increases dramatically. The factors that influence disease exacerbations were also recalled: prior exacerbations, worse dyspnoea, reduced lung function, comorbid cardiovascular disease, history of gastroesophageal reflux, comorbid depression, poorer quality of life, female sex, elevated white blood cell count and elevated eosinophil count. Concerning changes in body composition, L.G.W. Vanfleteren discussed the prospective ECRHS study which suggests that moderate and high weight gain over a 20-year period was associated with accelerated FVC and FEV₁ decline, while weight loss delayed the deterioration of the lungs. The presentation was concluded by pointing out the positive impact of PR on each of the listed areas of natural decline.

Concepts, goals, challenges and effects of maintenance programmes after pulmonary rehabilitation (C. Rochester, United States of America)

C. Rochester structured her presentation around 3 topics: i) rationale for maintenance PR; ii) the effects of maintenance programmes; and iii) challenges in the delivery of maintenance programmes (without digital health technologies), including areas for future research. The first section summarised the well-documented effects of PR for patients with different chronic respiratory diseases as well as factors that impact people's ability to continue exercise/active

after PR, i.e., motivation, home environment, weather, financial and family issues. A longitudinal study indicating different long-term (12 months) effects of PR was also presented [40]. It was mentioned that the form of maintenance training (supervised vs. unsupervised) also affects its effectiveness over time, and supervised programmes may decrease the risk of respiratory-related hospitalisation. The challenges of maintaining PR outcomes include a limited capacity of existing PR programmes, a limited number of healthcare professionals, funding and reimbursement limitations, travel/transport to maintenance programmes, as well as the low priority of insurance companies for preventive health. Future research areas may include: i) optimal maintenance strategies across different patient groups; ii) impact of patients' choice of options for maintenance programmes, iii) real-world cost-effectiveness of maintenance programmes, and iv) a combination of targeted maintenance/repetition of PR programmes, or novel models of maintenance using digital health technologies.

Take the active option: physical activity coaching programmes (H. Demeyer, Belgium)

H. Demeyer focused on the role of PA to maintain PR effects and discussed the relationship between changes in PA and in exercise capacity, as both are targeted through different interventions. Physical activity coaching programmes aim to increase PA focusing on low intensity activities in an unsupervised way, which can be done through digital technologies. Research on the effectiveness of coaching programmes showed that, in stable patients with COPD, such programmes yield a significant increase in PA, but effectiveness was less clear when implemented during PR. A small number of studies evaluating long-term PA coaching were also highlighted. The literature indicated that coaching programmes do not enhance exercise capacity (six-minute walking distance; 6MWD) beyond the minimal clinically important difference (MCID). This raises the question whether maintaining/improving PA provides sufficient benefit. A long-term comparison of PA and exercise capacity in patients participating in a 3-month PR programme only vs. patients participating in PR with an additional 3-month coaching programme during PR was also presented. In both cases, the trend was similar, with a decrease in both parameters in a similar manner 12 months after the intervention [41]. However, research conducted by the team of H. Demeyer (data not published) indicated that implementation of additional coaching sessions after completion of a 3-month PR programme resulted in the maintenance of PA levels, whereas the 'usual care' group not participating in coaching sessions after PR decreased their levels of PA. Nevertheless, a reduced exercise capacity was noted in both groups after completion of PR.

Role of digital health to improve maintenance (B. Caulfield, Ireland)

This presentation addressed the requirements for and components of successful PR maintenance strategies using digital tools. B. Caulfield started by recalling the results of the latest Cochrane meta-analysis [42] showing strong evidence for the effectiveness of supervised maintenance programmes on quality of life, but low effectiveness on disease exacerbations and hospitalisations in patients with COPD, and suggested that this area could be enhanced by digital technologies. P. Slevin, from B. Caulfield's team, performed a series of semi-structured interviews with patients with COPD and healthcare professionals about the use of digital technologies in the maintenance phase after PR [43-46]. Patients indicated that digital technologies provided them with an opportunity for self-management and enhanced communication with healthcare professionals. Clinicians indicated benefits in the availability of inter-appointment data and enhanced consultation efficiency, and recognised the role of digital technology to empower patients' self-management. Barriers to the adoption of digital technologies were also discussed. Patients identified primarily a lack of perceived usefulness and digital literacy, while clinicians mentioned the lack of quality data, little evidence base, and resource constraints. Based on these data, the requirements of digital health technologies in patients with COPD were determined. Such a hypothetical system must be effective in terms of health-related quality of life, exacerbations and functional status. It should also be safe, sustainable, appropriate to be implemented in patients' daily life. Therefore, supplementing the traditional management system, or partial replacement by digital health, can be carried out through devices addressing the following areas: measurement and monitoring, information and education, encouragement and engagement, peer support and augmented exercise.

Take-home messages

- Benefits of maintenance programmes after pulmonary rehabilitation are well-established; however, their effectiveness diminishes over time if there is no regular contact with the healthcare team.
- Available technology can be implemented in a comprehensive approach to monitoring patients' health condition, as well as for exercise training and physical activity coaching.
- Integrating digital health into the healthcare system requires further research that incorporates both patient and clinician perspectives.

Group 9.03: Nurses

Experts View: Hospital at home for the management of COVID-19

This Experts View session (session ID 16) highlighted the need to relieve hospitals during COVID-19 pandemic, giving patients access to hospital care and increasing the resilience of the healthcare systems. Safe alternatives to healthcare services were provided to patients with COVID-19 enabling treatment, monitoring and rehabilitation.

A Medicalized Hotel (C. Hernández, Spain)

C. Hernández presented a successful adaptation of a four-star hotel in Barcelona by a hospital-at-home team to provide hospital-level care to patients with COVID-19. In the period from March to May 2020, in the Hospital Clinic de Barcelona, 2,410 patients were hospitalised due to COVID-19, 21.4% of patients were transferred to Medicalized Hotel [47]. The main challenges were the multidisciplinary teamwork and there was particular importance given to separate clean and dirty areas. The equipment was also critical alongside professional experiences. The patients' experiences [48] were analysed very carefully, and the better experiences were related to patients who claimed to have been treated with respect. In summary, C. Hernández stated that the Medicalized Hotel is a safe alternative to conventional hospitals for patients with non-critical COVID-19 and can relieve hospitals from the overload.

Covid@Home (L. Brien, Ireland)

L. Brien described the rapid setup and performance of a virtual monitoring service for patients with COVID-19. This service allowed patients to be in their homes and access telehealth care performed by a multidisciplinary team and health technologies with the objective of their health monitoring. During 14 days, eligible patients had access to the programme, were monitored four times a day and had a phone service always available that they could use when necessary. After discharge, patients were referred to post-COVID rehabilitation as needed. This intervention alone was safe and responsible for reducing the length of hospital stay for patients with COVID-19, which lead to increasing bed availability. In addition, patient monitoring enabled the quick response to silent hypoxia and acute deterioration, which led to the reassurance of patients during the post-hospital period. L. Brien states that open communication between home teams and hospitals teams was decisive to the success of the interventions.

Take-home messages

- There are safe and adequate alternatives to the hospital regarding the treatment of patients with COVID-19.
- Teamwork and compromise were crucial elements regarding the reorganisation of healthcare solutions.

Oral Presentation: Nursing perspective during the COVID-19 pandemic

This oral presentation session (session ID 187) provided an overview of state-of-the-art advances in nursing interventions and experience during onset of COVID-19 pandemic, focusing on three main themes: impact on the nursing practice; innovation in healthcare delivery; and patients' perspective.

Living with the risk of being infected: COPD patients' experiences during the coronavirus pandemic (C.A. Mousing, Denmark)

C.A. Mousing and D. Sørensen [49] explored the perspectives of living with the risk of being infected with coronavirus in Danish patients with COPD (n=13), using individual semi-structured interviews. The authors found that living with the risk of being infected greatly affected everyday life of patients with moderate to severe COPD. There is a need to develop targeted interventions aimed at providing adequate information, supporting hope and maintaining warm distancing contact to ease symptoms and provide comfort to patients.

A negotiation of respiratory risk in the first phase of the Covid-19 pandemic (L. Welch, United Kingdom)

During the initial phase of the COVID-19 pandemic, various guidelines on infection control were published [50], which led to mixed messages in the early stages of the pandemic [51]. L. Welch [52] presented a nursing response to respiratory risk during the pandemic, where concerns were expressed regarding the working environment, the supply and availability of adequate personal protective equipment (PPE) and the unsafe working practice. Almost half of the respondents (48.6%) reported undertaking aerosol generating procedures. L. Welch reinforced that earlier and clearer guidance of infection control is required to ensure clinical safety and preparation for the future respiratory pandemics [51].

Awake proning in patients with COVID-19 (W. Stilma, Netherlands)

Prone positioning in selected patients with severe early-stage ARDS improves survival [53] and has been widely accepted only in recent years [54]. W. Stilma and colleagues [55] performed a clinical appraisal of the available literature and personal clinical experiences of healthcare workers in various settings around the world to formulate guidelines for awake prone positioning, with recommendations for its use. The authors concluded that awake proning is an attractive and safe adjunctive treatment for hypoxemia refractory to supplemental oxygen in patients with COVID-19 acute respiratory failure.

Non-Invasive Positive Pressure Ventilation training program for general wards Healthcare workers during Covid-19 outbreak (A. Godono, Italy)

The COVID-19 pandemic had a severe impact in the healthcare system and affected the availability of beds in hospitals and intensive care units [56]. A. Godono and colleagues [57] evaluated the efficiency of a non-invasive positive pressure ventilation (NPPV) training programme targeting healthcare professionals of general wards. The authors concluded that the training programme improved healthcare professionals' confidence levels in managing NPPV devices in non-intensive care units and could be a crucial element to improve patients' outcomes.

The impact of the COVID-19 pandemic on Portuguese rehabilitation nurses (M. Padilha, Portugal)

M. Padilha and R. Silva [58] assessed the impact of the pandemic on Portuguese rehabilitation nurses through an online questionnaire: the number of hours worked per week, the patient/nurse ratio, and the complexity of patients with COVID. Participants reported that the complexity of respiratory patients has increased while the satisfaction with the quality of nursing care has decreased during the pandemic. The COVID-19 pandemic posed several challenges to nurses, but rehabilitation nurses played a central role in stabilising, rehabilitating and reintegrating patients with COVID-19 [59].

Virtual consultations as follow up on home oxygen therapy for patients with COPD: A patient perspective (H. Schmidt, Denmark)

H. Schmidt [60] presented a qualitative study encompassing semi-structured interviews with 20 patients with COPD on long-term oxygen therapy to uncover patients' experiences with virtual consultation, thereby ensuring a patient-centred clinical practice. Findings revealed that virtual consultation was considered air- and energy-preserving due to avoiding transportation, waiting

time, handling oxygen tanks, and not having to involve others for help with transportation. Most participants emphasised their fear and vulnerability in relation to COVID-19, so being able to stay at home was a welcome initiative.

The SWISS Nurse; respiratory nursing care without any borders. An admission avoidance winter pressure initiative (E. Rickards, United Kingdom)

Over the last 7 years across the North of England, respiratory admissions have risen three times faster than general admissions [61]. The study presented by E. Rickards [62] introduced the role of a specialist respiratory nurse (Swiss nurse) within the emergency department, who provides ongoing support, equitable care for all patients and organises long-term chronic disease management. A total of 240 admissions were prevented after 5 months with the creation of this specialist respiratory nurse position. In conclusion, this initiative improves connections with the community respiratory teams.

Nurse-led community care (A. Madden, Ireland)

A. Madden et al. [63] implemented a Community Virtual Ward (VW) with the support of telemedicine to deliver specialist respiratory care in the community. Patients with exacerbations of asthma or COPD and/or at risk of deterioration were followed up by an Advanced Nurse Practitioner and supported by community nurses. In addition, an online electronic health record was accessible to the community and hospital clinicians. As the pandemic progressed, many countries reorganised their health care delivery systems and implemented digital communication tools to provide secure virtual care [64].

Was our step-down virtual covid ward safe and effective? (D. Jennings, United Kingdom)

D. Jennings and colleagues [65] demonstrated a VW model of care and its effectiveness on patients' outcomes. Eighty-nine patients were admitted to the VW, and findings showed that readmission rates within the 6-week period were similar to National figures (<10%). Mean mood and anxiety scores improved between day 1 and 14 and there was no increase in readmission rates for patients who required CPAP during admission and discharged patients with prescribed oxygen therapy. Patients' feedback on VW model of care was positive and the authors concluded that the system was very useful for patients, especially to those with chronic conditions.

Promoting physical activity by the Respiratory Nurses (S. Rocha, Portugal)

Pulmonary rehabilitation (PR) is the most effective non-pharmacologic treatment strategy to control symptoms and progression of COPD [66]. S. Rocha and colleagues [67] assessed the

impact and effectiveness of the respiratory nurses' interventions on the treatment of the patients involved in the programme (n=46). Interventions were adapted for each patient according to their individual characteristics. The average score of the PA domain of the London Chest Activity of Daily Living (LCADL) was 5.17 at baseline and 3.87 at the end of the programme (p<0.05). Findings showed the positive impact of nursing interventions in PR by improving the individual skills and self-care behaviour of patients with COPD.

Take-home messages

- Virtual consultations and follow-up are an effective measure in monitoring patients with chronic respiratory disease.
- There is a need for targeted interventions aimed at giving adequate information, supporting hope and maintaining warm distancing contact.
- Psychological support must be integrated as a part of the virtual model care for the respiratory patients.

Group 9.04: Psychologists and behavioural scientists

Oral Presentation: Psychological and behavioural issues in respiratory care

This oral presentation session (session ID 59) was the inaugural session with abstracts submitted to the novel Psychologists and Behavioural Scientists (9.04) group. The session focused on the role of psychological issues in health care providers, caregivers and patients; in COVID-19 and other respiratory conditions.

No gender differences in neural gating of respiratory sensations in COPD patients – preliminary findings (Y. Denutte, Belgium)

Based on the neural gating model of respiratory sensations (NGRS), Y. Denutte [68] presented the results of an electroencephalography (EEG) study of gender differences in dyspnoea perception. According to the NGRS, dyspnoea results from reduced neural gating of respiratory afferent signals into the cerebral cortex, and it was hypothesised that female patients would show reduced NGRS. EEG measurements were performed on 11 female and 15 male patients with COPD when presented with paired inspiratory occlusions. The results indicated no significant gender differences in NGRS, but larger studies are needed to confirm these findings. Identifying relevant biological, psychological, and behavioural predictors of differences in

dyspnoea perception is highly important in the holistic and patient-centred approach to COPD care.

Translation and initial validation of the Danish version of the COPD-Anxiety Questionnaire (S. Rubio-Rask, Denmark)

The COPD Anxiety Questionnaire was originally constructed in German to measure disease-specific anxiety in patients with COPD across five domains: i) fear of dyspnoea, ii) fear of disease progression, iii) fear of physical activity, iv) fear of social exclusion, v) sleep-related worries. On behalf of S. Rubio-Rask, I. Farver-Vestergaard [69] presented their Danish translation of the questionnaire (CAF-DK), as well as a study performing the initial validation of the CAF-DK in a sample of 260 patients with COPD (238 electronic responses; 22 paper responses). Results showed that Cronbach's alpha values for total- and domain-scales were comparable to those of the original CAF, as was the correlation coefficients between CAF-DK and concurrent measures (General Anxiety Disorder-7; COPD Assessment Test). In conclusion, the CAF-DK can be used to measure COPD-specific anxiety in Danish-speaking patients, yet further studies are needed to explore the psychometric properties of the CAF, compared with other disease-specific measures of anxiety in COPD.

Optimism, exacerbations, and health outcomes of pulmonary rehabilitation in patients with COPD (T. Janssens, Belgium)

Pulmonary rehabilitation (PR) is a cost-effective treatment for patients with COPD, but not all patients benefit from this treatment to the same extent. T. Janssens [70] presented the results of a prospective study focusing on the role of optimism as a predictor of treatment response across 3-week inpatient PR and 6-month follow-up in 370 patients with COPD. Results showed that higher levels of optimism at baseline were associated with lower depressive symptoms and better health status across all timepoints. Contrary to what was hypothesised, the positive associations between optimism and health status were limited to patients that did not experience an exacerbation during follow-up. These findings confirm the role of patient beliefs on outcome trajectories, but also highlight boundaries of these effects.

Technology-dependent children with respiratory problems on a home care program: Exploring the family experience (M. Galogavrou, Greece)

An increasing number of children have respiratory problems needing paediatric home care, which has a major impact on the parents and home environment. M. Galogavrou [71] presented the results of an observational study on parents' perceptions of paediatric health care services

and family well-being for children (n=51, 27 boys) depending on paediatric home care (mechanical ventilation n=25, oxygen dependent n=13, tracheostomy n=8, non-invasive ventilation n=5). Most parents had completed higher education, were employed, and married. Results showed high levels of satisfaction with care. Nevertheless, 27.4% of parents reported moderate or severe depressive symptoms and 35.2% reported moderate-severe anxiety symptoms. These findings suggest a need for attention to parental well-being in this population.

Psychological functioning and lifestyle behaviours of patients with idiopathic pulmonary fibrosis: a prospective study (A. Delameillieure, Belgium)

Idiopathic pulmonary fibrosis (IPF) is a rare and severe lung disease with a relatively poor prognosis, which beyond the physical symptoms, impacts psychological and behavioural factors. A. Delameillieure [72] presented the results of a prospective study on psychological functioning and lifestyle behaviours of patients with IPF, conducted at the University Hospitals Leuven, Belgium. The cohort of patients (n=66; 72% male; mean age 70 years) were included at IPF diagnosis and completed questionnaires assessing depression, anxiety, sun protection, PA and alcohol intake at each hospital visit during the following two years. Results showed that a considerable proportion of patients experienced symptoms of anxiety and depression, as well as suboptimal lifestyle behaviour patterns across the illness trajectory. However, response rates were low and conclusions should therefore be made with caution. Nonetheless, the identification of psychological and lifestyle factors across the illness trajectory is an important step towards providing IPF care that is more holistic and patient-centred.

Mental health during COVID-19 in Dutch adolescents and adults with cystic fibrosis (CF) or primary ciliary dyskinesia (PCD) and their caregivers (M. Verkleij, The Netherlands)

Patients with CF or PCD as well as their caregivers have a high risk to develop anxiety or depressive symptoms. Furthermore, COVID-19 related measures, such as national lockdowns and social distancing, have a potential for negative impact on mental health. M. Verkleij [73] presented the results of a prospective study investigating changes in anxiety, depression, and resilience in Dutch adolescents with CF or PCD (n=41) and caregivers of patients (n=69) during the second wave of the COVID-19 pandemic. Results showed overall high prevalence of anxiety symptoms (31-53% reporting at least mild depressive symptoms; PHQ-9>5, 33-36% reporting at least mild anxiety symptoms (GAD-7>5) during the second lockdown. However, anxiety and depressive symptoms were unchanged compared to pre-pandemic measures. These results highlight a continued mental health burden in patients with CF or PCD and their caregivers.

Psychological sequelae among health personnel treated for COVID-19 at the Hassan II hospital in Agadir (F. Bounoua, Morocco)

The COVID-19 pandemic has led to increased demands on healthcare professionals, especially healthcare workers in the front line of management of COVID-19. F. Bounoua [74] presented results of an observational study investigating anxiety and depressive symptoms as well as potential post-traumatic stress disorder (PTSD) in staff of a single Moroccan hospital treated for COVID-19 (n=67, 58% women, 32% with comorbidities, 3% needing intensive care treatment). From these, 64% of participants reported residual symptoms after recovery. Only 42% of participants did not report anxiety or depressive symptoms (measured using the Hospital Anxiety and Depression Scale, HADS), and 10% of participants showed symptoms of PTSD. These results show a strong psychological impact in this group of healthcare workers treated for COVID-19 and highlights the need for psychological support.

Impact of COVID-19 on mental health: a traumatic event (S. Houben-Wilke, the Netherlands)

Residual symptoms of COVID-19 have been reported, but longer-term symptoms and psychological burden in patients with residual symptoms remains unclear. On behalf of S. Houben-Wilke, M. Van Herck [75] presented the results of a prospective survey investigating symptoms of anxiety and depression (using HADS) as well as PTSD in a group of participants (n=239, 62 hospitalised) 3 and 6 months after confirmed COVID-19 diagnosis. Participants reported good health status before infection (79% in hospitalised vs. 90% in non-hospitalised patients, $p < 0.05$). Considering the whole sample, 35% of patients reported symptoms of anxiety 3 months after infection, which was not reduced at 6 months, and nearly 50% of patients reported symptoms of anxiety 3 months after infection with a small reduction after 6 months. Forty percent (40%) of patients were at risk for PTSD 3 months after infection, which was reduced to 30% 6 months after infection. Symptoms of anxiety, depression, and PTSD were comparable among hospitalised and non-hospitalised patients. Findings suggest that a substantial proportion of patients with residual symptoms after COVID-19 show psychological complications, and awareness of healthcare providers and access to care is needed to address these problems.

Sex differences in mental health of hospitalized patients with COVID-19 (Z. Barmparessou, Greece)

COVID-19 hospitalisation can pose a considerable mental burden on patients, also after hospital discharge. Z. Barmparessou [76] presented the results of a survey performed in a large COVID-19 reference hospital in Athens, Greece. Depression, anxiety, post-traumatic stress disorder

(PTSD), sleeping difficulties and quality of life were assessed in 131 patients 1-2 months after discharge from COVID-19 hospitalisation. Prevalence rates of clinically significant symptom levels were high (PTSD=50%; sleep disorder=41%; anxiety=35%; depression=20%), suggesting a considerable general mental burden on patients upon COVID-19 hospital discharge. Moreover, prevalence rates were significantly higher for female compared to male patients, in spite of females spending fewer days at the hospital. As this was a cross-sectional study with no non-COVID-19 control group, no causal inferences could be made. However, the results do highlight the potential need for mental health screening and follow-up in patients that have been hospitalised with COVID-19.

Take-home messages

- Respiratory conditions are associated with psychological distress in patients and caregivers.
- Psychological and behavioural aspects of the respiratory disease show associations with important health outcomes.
- Measuring psychological distress is a first step towards the identification of psychological healthcare needs in patients with respiratory disease.

Concluding remarks

This article provides valuable insights on the latest scientific data and emerging areas affecting the clinical practice of Allied Respiratory Professionals, presented at the Virtual 2021 ERS International Congress. We look forward to seeing you in Barcelona in 2022 for another highly successful ERS Congress!

Acknowledgments: Ana Machado's work is supported by Fundação para a Ciência e Tecnologia (reference: SFRH/BD/147200/2019) and by Bijzonder Onderzoeksfonds – Bilaterale Samenwerking (BOF BILA) from Hasselt University (BOF BILA reference: DOC/SCHL-BOSE/190/522). J. Cruz acknowledges the support of the Center for Innovative Care and Health Technology (ciTechCare), funded by Portuguese national funds provided by Fundação para a Ciência e Tecnologia (FCT) (UIDB/05704/2020).

References:

1. Savushkina, O., A. Cherniak, A. Zaitsev et al., Ventilation heterogeneity after COVID-19. Eur Respir J 2021; 58(suppl 65): OA2681.

2. Hurst, C., S. Yearsley, H. Stone et al., Evaluation of patient attendance to respiratory physiology in the midst of the COVID-19 Pandemic. *Eur Respir J* 2021; 58(suppl 65): OA2682.
3. National Health Service (NHS) Foundation Trust. *2019 Outpatients project – Results published*. Last updated: October 17 2019; Last accessed: October 19 2021]; Available from: <https://www.nhsbenchmarking.nhs.uk/news/2019-outpatients-project-results-published>.
4. Steenbruggen, I., P. De Laat, A.H.M. Van De Geijn et al., Implementation of the 2017 ERS Methacholine Challenge Standard. *Eur Respir J* 2021; 58(suppl 65): OA2683.
5. Coates, A.L., J. Wanger, D.W. Cockcroft et al., ERS technical standard on bronchial challenge testing: general considerations and performance of methacholine challenge tests. *Eur Respir J* 2017; 49(5): 1601526.
6. De Soomer, K., L. Casas, H. Vaerenberg et al., Evaluation of the Global Lung function Initiative reference values for spirometry and diffusion capacity. *Eur Respir J* 2021; 58(suppl 65): OA2684.
7. Quanjer, P.H., S. Stanojevic, T.J. Cole et al., Multi-ethnic reference values for spirometry for the 3–95-yr age range: the global lung function 2012 equations. *Eur Respir J* 2012; 40(6): 1324-1343.
8. Stanojevic, S., B.L. Graham, B.G. Cooper et al., Official ERS technical standards: Global Lung Function Initiative reference values for the carbon monoxide transfer factor for Caucasians. *Eur Respir J* 2017; 50(3): 1700010.
9. Mottram, C.D., T. Mccaskill, J. Road et al., Pulmonary airway resistance variability in normal biologic quality control subjects among pulmonary function testing systems across British Columbia. *Eur Respir J* 2021; 58(suppl 65): OA2685.
10. Robertson, L., R. Lowry, and K. Sylvester, Do bacterial/viral filters impact quality assurance verification of ultrasonic spirometers? *Eur Respir J* 2021; 58(suppl 65): OA2686.
11. Oestreich, M.-A., B. Frauchiger, P. Latzin et al., Breath detection algorithms effect lung function outcomes. *Eur Respir J* 2021; 58(suppl 65): OA2687.
12. Stanojevic, S., P. Powell, G. Hall et al., Real-world application of Spirometry Quality Control Deep-Learning Algorithm. *Eur Respir J* 2021; 58(suppl 65): OA2688.
13. Das, N., K. Verstraete, S. Stanojevic et al., Deep-learning algorithm helps to standardise ATS/ERS spirometric acceptability and usability criteria. *Eur Respir J* 2020; 56(6).
14. Miller, M.R., J. Hankinson, V. Brusasco et al., Standardisation of spirometry. *Eur Respir J* 2005; 26(2): 319-338.
15. Veneroni, C., A. Acciarito, P. Pompilio et al., Use of artificial intelligence for improving quality control of oscillometry. *Eur Respir J* 2021; 58(suppl 65): OA2689.

16. King, G.G., J. Bates, K.I. Berger et al., Technical standards for respiratory oscillometry. *Eur Respir J* 2020; 55(2): 1900753.
17. Del Corral Nuñez-Flores, T., R. Fabero-Garrido, G. Plaza-Manzano et al., Late Breaking Abstract - Effects of respiratory muscle training on quality of life, physical function, and psychological state in Post-COVID Syndrome. *Eur Respir J* 2021; 58(suppl 65): OA2690.
18. Baumgartner, R.N., K.M. Koehler, D. Gallagher et al., Epidemiology of sarcopenia among the elderly in New Mexico. *Am J Epidemiol* 1998; 147(8): 755-763.
19. Cruz-Jentoft, A.J., J.P. Baeyens, J.M. Bauer et al., Sarcopenia: European consensus on definition and diagnosis Report of the European Working Group on Sarcopenia in Older People A. J. Cruz-Gentoft et al. *Age Ageing* 2010; 39(4): 412-423.
20. Morley, J.E., A.M. Abbatecola, J.M. Argiles et al., Sarcopenia with limited mobility: an international consensus. *J Am Med Dir Assoc* 2011; 12(6): 403-409.
21. Muscaritoli, M., S. Anker, J. Argiles et al., Consensus definition of sarcopenia, cachexia and pre-cachexia: joint document elaborated by Special Interest Groups (SIG) "cachexia-anorexia in chronic wasting diseases" and "nutrition in geriatrics". *Clin Nutr* 2010; 29(2): 154-159.
22. International Working Group on Sarcopenia, Sarcopenia: an undiagnosed condition in older adults. Current consensus definition: prevalence, etiology, and consequences. *J Am Med Dir Assoc* 2011; 12(4): 249.
23. Mitchell, W.K., P.J. Atherton, J. Williams et al., Sarcopenia, dynapenia, and the impact of advancing age on human skeletal muscle size and strength; a quantitative review. *Front Physiol* 2012; 3: 260.
24. Bauer, J., J.E. Morley, A.M. Schols et al., Sarcopenia: a time for action. An SCWD position paper. *Journal of cachexia, sarcopenia and muscle* 2019; 10(5): 956-961.
25. Bhasin, S., T.G. Travison, T.M. Manini et al., Sarcopenia definition: the position statements of the sarcopenia definition and outcomes consortium. *J Am Geriatr Soc* 2020; 68(7): 1410-1418.
26. Cruz-Jentoft, A.J., G. Bahat, J. Bauer et al., Sarcopenia: revised European consensus on definition and diagnosis. *Age Ageing* 2019; 48(1): 16-31.
27. Schols, A.M., I.M. Ferreira, F.M. Franssen et al., *Nutritional assessment and therapy in COPD: a European Respiratory Society statement*. 2014, Eur Respiratory Soc.
28. van Bakel, S.I., H.R. Gosker, R.C. Langen et al., Towards Personalized Management of Sarcopenia in COPD. *Int J Chron Obstruct Pulmon Dis* 2021; 16: 25.

29. Langen, R., H. Gosker, A. Remels et al., Triggers and mechanisms of skeletal muscle wasting in chronic obstructive pulmonary disease. *The international journal of biochemistry & cell biology* 2013; 45(10): 2245-2256.
30. Casaburi, R., S. Bhasin, L. Cosentino et al., Effects of testosterone and resistance training in men with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2004; 170(8): 870-878.
31. Polkey, M.I., J. Praestgaard, A. Berwick et al., Activin type II receptor blockade for treatment of muscle depletion in chronic obstructive pulmonary disease. A randomized trial. *Am J Respir Crit Care Med* 2019; 199(3): 313-320.
32. McAuley, H.J., T.C. Harvey-Dunstan, M. Craner et al., Longitudinal changes to quadriceps thickness demonstrate acute sarcopenia following admission to hospital for an exacerbation of chronic respiratory disease. *Thorax* 2021; 76(7): 726-728.
33. Benz, E., K. Trajanoska, L. Lahousse et al., Sarcopenia in COPD: a systematic review and meta-analysis. *European respiratory review* 2019; 28(154).
34. Dent, E., J. Morley, A. Cruz-Jentoft et al., International clinical practice guidelines for sarcopenia (ICFSR): screening, diagnosis and management. *J Nutr Health Aging* 2018; 22(10): 1148-1161.
35. Cruz-Jentoft, A.J. and A.A. Sayer, Sarcopenia. *The Lancet* 2019; 393(10191): 2636-2646.
36. Jones, S.E., M. Maddocks, S.S. Kon et al., Sarcopenia in COPD: prevalence, clinical correlates and response to pulmonary rehabilitation. *Thorax* 2015; 70(3): 213-218.
37. Covey, M.K., E.G. Collins, S.I. Reynertson et al., Resistance training as a preconditioning strategy for enhancing aerobic exercise training outcomes in COPD. *Respir Med* 2014; 108(8): 1141-1152.
38. Teschler, M., M. Heimer, B. Schmitz et al., Four weeks of electromyostimulation improves muscle function and strength in sarcopenic patients: a three-arm parallel randomized trial. *Journal of Cachexia, Sarcopenia and Muscle* 2021.
39. SPRINTT trial consortium, *IMI1 Final Project Report Public Summary*.
40. Soicher, J.E., N.E. Mayo, L. Gauvin et al., Trajectories of endurance activity following pulmonary rehabilitation in COPD patients. *Eur Respir J* 2012; 39(2): 272-8.
41. Altenburg, W.A., N.H. ten Hacken, L. Bossenbroek et al., Short- and long-term effects of a physical activity counselling programme in COPD: a randomized controlled trial. *Respir Med* 2015; 109(1): 112-21.
42. Malaguti, C., S. Dal Corso, S. Janjua et al., Supervised maintenance programmes following pulmonary rehabilitation compared to usual care for chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* 2021; 8(8): Cd013569.

43. Slevin, P., T. Kessie, J. Cullen et al., A qualitative study of chronic obstructive pulmonary disease patient perceptions of the barriers and facilitators to adopting digital health technology. *Digital health* 2019; 5: 2055207619871729-2055207619871729.
44. Slevin, P., T. Kessie, J. Cullen et al., Exploring the potential benefits of digital health technology for the management of COPD: a qualitative study of patient perceptions. *ERJ open research* 2019; 5(2): 00239-2018.
45. Slevin, P., T. Kessie, J. Cullen et al., Exploring the barriers and facilitators for the use of digital health technologies for the management of COPD: a qualitative study of clinician perceptions. *QJM* 2020; 113(3): 163-172.
46. Slevin, P., T. Kessie, J. Cullen et al., A qualitative study of clinician perceptions regarding the potential role for digital health interventions for the management of COPD. *Health Informatics J* 2021; 27(1): 1460458221994888.
47. Torrallardona-Murphy, O., J.M. Pericàs, N. Rabaneda-Lombarte et al., Medicalized Hotel as an Alternative to Hospital Care for Management of Noncritical COVID-19. *Ann Intern Med* 2021; 174(9): 1338-1340.
48. Cervera, M., J. Escarrabill, E. Palou et al., Patient's Experiences in a Medicalized Hotel for Covid-19 Acute Care Support. An Observational Study with a Cross-Sectional Design. *Am J Respir Crit Care Med* 2021; 203: A3190.
49. Mousing, C.A. and D. Sørensen, Living with the risk of being infected: COPD patients' experiences during the coronavirus pandemic. *Eur Respir J* 2021; 58(suppl 65): OA1503.
50. Birgand, G., N.T. Mutters, J. Otter et al., Analysis of national and international guidelines on respiratory protection equipment for COVID-19 in healthcare settings. *medRxiv* 2020: 2020.04.23.20077230.
51. Bird, P., V. Badhwar, K. Fallon et al., High SARS-CoV-2 infection rates in respiratory staff nurses and correlation of COVID-19 symptom patterns with PCR positivity and relative viral loads. *J Infect* 2020; 81(3): 452-482.
52. Welch, L., N. Roberts, K. Lippiett et al., A negotiation of respiratory risk in the first phase of the Covid-19 pandemic. *Eur Respir J* 2021; 58(suppl 65): OA1504.
53. Scholten, E.L., J.R. Beitler, G.K. Prisk et al., Treatment of ARDS With Prone Positioning. *Chest* 2017; 151(1): 215-224.
54. Paul, V., S. Patel, M. Royse et al., Proning in Non-Intubated (PINI) in Times of COVID-19: Case Series and a Review. *J Intensive Care Med* 2020; 35(8): 818-824.
55. Stilma, W., L. Bos, L. Pisani et al., Awake prone in patients with COVID-19. *Eur Respir J* 2021; 58(suppl 65): OA1505.

56. Sánchez-Úbeda, E.F., P. Sánchez-Martín, M. Torrego-Ellacuría et al., Flexibility and Bed Margins of the Community of Madrid's Hospitals during the First Wave of the SARS-CoV-2 Pandemic. *Int J Environ Res Public Health* 2021; 18(7): 3510.
57. Godono, A., D. Gavetti, M. Clari et al., Non-Invasive Positive Pressure Ventilation training program for general wards Healthcare workers during Covid-19 outbreak. *Eur Respir J* 2021; 58(suppl 65): OA1506.
58. Padilha, M. and R. Silva, The impact of the COVID-19 pandemic on Portuguese rehabilitation nurses. *Eur Respir J* 2021; 58(suppl 65): OA1507.
59. Vitacca, M., M. Carone, E.M. Clini et al., Joint Statement on the Role of Respiratory Rehabilitation in the COVID-19 Crisis: The Italian Position Paper. *Respiration* 2020; 99(6): 493-499.
60. Schmidt, H.C., G. Tulinius, and H.M. Christensen, Virtual consultations as follow up on home oxygen therapy for patients with COPD: A patient perspective. *Eur Respir J* 2021; 58(suppl 65): OA1508.
61. British Lung Foundation. *Out in the cold – lung disease, the hidden driver of NHS winter pressure*. 2017; Last accessed: 25 November 2021. Available from: <https://www.blf.org.uk/policy/out-in-the-cold>.
62. Rickards, E., L. Ascough, S. Sibley et al., The SWISS Nurse; respiratory nursing care without any borders. An admission avoidance winter pressure initiative. *Eur Respir J* 2021; 58(suppl 65): OA1509.
63. Madden, A., B. Cushen, and C. Lewis, Nurse-led community care. *Eur Respir J* 2021; 58(suppl 65): OA1510.
64. Temesgen, Z.M., D.C. DeSimone, M. Mahmood et al., Health Care After the COVID-19 Pandemic and the Influence of Telemedicine. *Mayo Clin Proc* 2020; 95(9S): S66-S68.
65. Jennings, D., K. Hills, J. Briscoe et al., Late Breaking Abstract - Was our step-down virtual covid ward safe and effective? *Eur Respir J* 2021; 58(suppl 65): OA1511.
66. Wada, J.T., E. Borges-Santos, D.C. Porras et al., Effects of aerobic training combined with respiratory muscle stretching on the functional exercise capacity and thoracoabdominal kinematics in patients with COPD: a randomized and controlled trial. *Int J Chron Obstruct Pulmon Dis* 2016; 11: 2691-2700.
67. Rocha, S.M.P., M.F. Rodrigues, G. Marques et al., Late Breaking Abstract - Promoting physical activity by the Respiratory Nurses. *Eur Respir J* 2021; 58(suppl 65): OA1512.
68. Denutte, Y., T. Reijnders, R. Gosselink et al., No gender differences in neural gating of respiratory sensations in COPD patients – preliminary findings. *Eur Respir J* 2021; 58(suppl 65): OA183.

69. Rubio-Rask, S., I. Farver-Vestergaard, O. Hilberg et al., Translation and initial validation of the Danish version of the COPD-Anxiety Questionnaire. *Eur Respir J* 2021; 58(suppl 65): OA184.
70. Janssens, T., W. Geidl, J. Carl et al., Optimism, exacerbations and health outcomes of pulmonary rehabilitation in patients with COPD. *Eur Respir J* 2021; 58(suppl 65): OA185.
71. Galogavrou, M., E. Hatzigorou, P. Vantsi et al., Technology-dependent children with respiratory problems on a home care program: Exploring the family experience. *Eur Respir J* 2021; 58(suppl 65): OA186.
72. Delameillieure, a., F. Dobbels, K. Leceuvre et al., Psychological functioning and lifestyle behaviours of patients with idiopathic pulmonary fibrosis: a prospective study. *Eur Respir J* 2021; 58(suppl 65): OA187.
73. Verkleij, M., L. Noij, S. Hashimoto et al., Mental health during COVID-19 in Dutch adolescents and adults with cystic fibrosis (CF) or primary ciliary dyskinesia (PCD) and their caregivers. *Eur Respir J* 2021; 58(suppl 65): OA188.
74. Bounoua, f., H. Moubachir, and H. Serhane, Psychological sequelae among health personnel treated for COVID 19 at the Hassan II hospital in Agadir. *Eur Respir J* 2021; 58(suppl 65): OA189.
75. Houben-Wilke, S., Y. Goërtz, J. Delbressine et al., Impact of COVID-19 on mental health: a traumatic event. *Eur Respir J* 2021; 58(suppl 65): OA190.
76. Barmparaessou, Z., S. Pappa, A. Pappas et al., Sex differences in mental health of hospitalized patients with COVID-19. *Eur Respir J* 2021; 58(suppl 65): OA191.

Figures




Figure 1. Classification of airway hyperresponsiveness (AHR) according to the European Respiratory Society (ERS) 2017 technical standard [5], presented at the oral communication “Implementation of the 2017 ERS Methacholine Challenge Standard” (I. Steenbruggen, The Netherlands).

Tables

Table 1. Current definitions of sarcopenia.

	EWGSOP2 [26]	SCWD [24]	SDOC [25]
Probable sarcopenia	Low muscle strength		
Sarcopenia	Low muscle strength + Low muscle quantity/quality	Low muscle strength + Low muscle quantity/quality	Low muscle strength + Low physical performance
Severe sarcopenia	Low muscle strength + Low muscle quantity/quality + Low physical performance		
Primary sarcopenia	Age-related, with no other evident specific cause	Age-related, with no other evident specific cause	
Secondary sarcopenia	Secondary to a systemic disease (e.g., chronic respiratory disease)	Secondary to a systemic disease (e.g., chronic respiratory disease)	

EWGSOP, European working group in sarcopenia in older people; SCWD, Society of sarcopenia, cachexia and wasting disorders; SDOC, Sarcopenia definition and outcomes consortium.