Early View

Original research article

The interrelationship of chronic cough and depression: a prospective population-based study

Johnmary T. Arinze, Amy Hofman, Emmely W de Roos, Maria A. J. de Ridder, Katia M. C. Verhamme, Bruno Stricker, Guy G. Brusselle, Annemarie I. Luik

Please cite this article as: Arinze JT, Hofman A, de Roos EW, *et al.* The interrelationship of chronic cough and depression: a prospective population-based study. *ERJ Open Res* 2022; in press (<https://doi.org/10.1183/23120541.00069-2022>).

This manuscript has recently been accepted for publication in the *ERJ Open Research*. It is published here in its accepted form prior to copyediting and typesetting by our production team. After these production processes are complete and the authors have approved the resulting proofs, the article will move to the latest issue of the ERJOR online.

Copyright ©The authors 2022. This version is distributed under the terms of the Creative Commons Attribution Non-Commercial Licence 4.0. For commercial reproduction rights and permissions contact permissions@ersnet.org

The interrelationship of chronic cough and depression: a prospective population-based study

Johnmary T. Arinze, MD^{1,2}, Amy Hofman, MSc³, Emmely W de Roos, MD^{1,3}, Maria A. J. de Ridder, PhD², Katia M. C. Verhamme, MD, PhD^{2,5}, Bruno Stricker, MD, PhD³, Guy G. Brusselle, MD, PhD^{1,3,4}, Annemarie I. Luik, PhD^{3,5*}

Affiliations:

¹ Department of Respiratory Medicine, Ghent University Hospital, Ghent, Belgium

² Department of Medical Informatics, Erasmus University Medical Center (MC), Rotterdam, The Netherlands

³ Department of Epidemiology, Erasmus University Medical Center (MC), Rotterdam, The Netherlands

⁴ Department of Respiratory Medicine, Erasmus University Medical Center (MC), Rotterdam, The Netherlands

⁵ Department of Bioanalysis, Faculty of Pharmaceutical Sciences, Ghent University, Ghent, Belgium.

*Department of Epidemiology, Erasmus University Medical Center (MC), Rotterdam, The Netherlands.

Electronic address: a.luik@erasmusmc.nl

Word count: 3,013

ABSTRACT

Background Chronic cough is a debilitating medical condition that is often complicated by psychomorbidities such as depressive symptoms. Nevertheless, little is known about the impact of chronic cough on the risk of developing depression. Therefore, we investigated the association between chronic cough and prevalent, incident, as well as recurrent depression in a population-based sample of middle-aged and older persons.

Methods Within the Rotterdam Study, a population-based cohort, we defined chronic cough as reporting daily coughing for at least 3 months. Depression was assessed using the Center for Epidemiologic Studies – Depression scale, clinical interviews, and medical records. Associations between chronic cough and depression were determined with linear, logistic and Cox regression analyses.

Results The study included 5,877 participants (mean age $\pm$ standard deviation: 72 ± 8 years, 59% female) who contributed 37,287 person-years of follow-up. At baseline, participants with chronic cough reported more depressive symptoms (adjusted standardized mean difference 0.15, 95% CI 0.07 – 0.22) compared to those without chronic cough. Over time, chronic cough was associated with an increased risk of depression in participants with a history of depression (HR 1.45, 95% CI 1.13 – 1.84) but not in those without a history of depression (HR 0.91, 95% CI 0.68 – 1.22).

Conclusions Adults with chronic cough have a disproportionate burden of depressive symptoms and an increased risk of recurrent depression. This highlights the importance of screening for depression in patients with chronic cough.

Introduction

Chronic cough, defined as a cough lasting at least eight weeks, is a common medical condition affecting about 1 – 12% of the general adult population[1, 2]. It is debilitating, often complicating associated clinical disorders such as asthma and chronic obstructive pulmonary disease (COPD), and necessitating visits to primary care and referral to specialist clinics[3-7].

Individuals with chronic cough frequently experience physical and psychosocial conditions including social isolation, physical exhaustion, sleep complaints, stress incontinence, cough syncope, and a lower quality of life[8-13]. Also, mental health complaints including depressive symptoms and major depressive disorder are common in people with chronic cough[5, 9, 13-15]. Depression additionally contributes to the overall clinical burden of the diseases commonly underlying chronic cough[13, 16]. Potentially, chronic cough may even have a negative impact on the risk of developing depression[17, 18]. The complex nature of risk factors for depression is such that the vulnerability for onset and recurrent depressive events differs with regards to triggers[18]. Some of these factors, such as low socio-economic status and reduced work ability are also more prevalent in patients with chronic cough[19, 20].

Previous studies have highlighted the link between chronic cough and depression[17]. Clinical studies have shown that treatment of chronic cough also decreases depressive symptoms in patients with chronic cough[9, 13], which may suggest that chronic cough could be a causal factor for developing depressive symptoms. For example, French *et al* studied the impact of cough treatment on depressive symptoms in 55 women with a 7-year mean cough duration and reported a significant reduction in depressive symptoms after three and six months of treating chronic cough[9]. Similarly, Dicipinigaitis *et al* followed 81 chronic cough

patients for three months and found that cough treatment significantly reduced both cough severity and depressive symptom scores[13].

So far, most studies on chronic cough and depression involved patients referred to specialist cough clinics, whose clinical profiles may differ from that of the general population[9, 13, 16]. Furthermore, the evidence on the burden of depression in chronic cough in population-based samples has come mainly from cross-sectional studies and provided little insight into the temporal association between chronic cough and depression[5, 15].

We therefore conducted a population-based observational cohort study in middle-aged and older adults to investigate the association between chronic cough and prospectively assessed depressive symptoms and disorders over time.

Methods

Study design and study population

The Rotterdam Study is a prospective population-based cohort study on the epidemiology of chronic diseases among 14,926 middle-aged and older adults (aged ≥ 45 years) living in the well-defined Ommoord district, a suburb of the city of Rotterdam, the Netherlands[21]. The Rotterdam Study data are acquired through home interviews and clinical examinations at the research center every 3 to 6 years, and from medical records of the general practitioners (GPs), hospitals, pharmacies, and nursing homes.

Within this study, we included participants who had chronic cough data collected during the home interview between January 2002 and December 2005, and had cross-sectional data on depressive symptoms from the home interview and/or data on incident depressive events via the follow-up of medical records up until February 2012, see figure 1.

The Medical Ethics Committee of the Erasmus Medical Centre (registration number MEC 02.1015) and by the Dutch Ministry of Health, Welfare and Sport (Population Screening Act WBO, license number 1071272-159521-PG) approved the Rotterdam Study. The Rotterdam Study has been registered with the Netherlands National Trial Register (NTR; www.trialregister.nl) and into the WHO International Clinical Trials Registry Platform (ICTRP; www.who.int/ictrp/network/primary/en/) under the joint catalogue number NTR6831. All participants provided written informed consent to participate in the study and to have their information obtained from treating physicians[21].

Chronic cough

Chronic cough was assessed using the question “In the last 2 years, did you cough almost every day for three consecutive months or more?”[22]. Chronic cough was defined as daily coughing lasting for at least 3 months, in line with most epidemiological studies[1].

Depression

Depression was assessed in three ways. First, during the home interviews, depressive symptoms were assessed with the Dutch version of the Center for Epidemiologic Studies - Depression Scale (CES-D). The CES-D scale ranges from 0 through 60 and assesses the burden of self-reported depressive symptoms with higher scores denoting more severe symptoms[23, 24]. A score of 16 and above was defined as clinically relevant depressive symptoms. This measure of depression was used in our cross-sectional analyses. Second, if participants had a score of 16 or above, they were invited for a semi-structured interview by a trained professional using the Dutch version of the Schedules for Clinical Assessment in Neuropsychiatry (SCAN)[25] to identify depressive disorders. Based on these interviews, participants were diagnosed as having no depression, depressive symptoms, a depressive syndrome (i.e. mild depressive disorder or dysthymia), or major depressive disorder according to the Statistical Manual of Mental Disorders, 4th revised edition (DSM-IV-TR) criteria[26]. Third, we used medical records, including data from GP’s notes, specialist reports, and hospital discharge letters, to assess depression in between examination rounds, coding both depressive symptoms and diagnoses of depressive disorders. Participants were categorized as having a history of depression if they experienced a depressive event during five years prior to or at baseline.

For our longitudinal analyses, we combined data of the CES-D and SCAN across multiple research rounds and medical records to indicate any incident event of depression, including depressive symptoms, depressive syndrome, and major depressive disorder. Within this

project we focused on the first occurring event. Participants were continuously followed from baseline (2002 – 2005) until the occurrence of the first event of depressive condition, death or the end of study (February 2012), whichever came first.

Other variables

Other variables relevant to the association of chronic cough with depression were assessed at baseline including participants' age and sex. Body mass index (BMI) was calculated as the ratio of weight in kilogram to height in squared meter as measured with calibrated scales. Smoking was assessed in the interview and divided into three categories: never, past, and current smoking. Asthma was physician-diagnosed, and COPD cases were validated using spirometry data and medical records. The diagnoses of asthma and/or COPD at baseline were grouped as obstructive airway disease. Cases of lung cancer were ascertained with the Dutch cancer registry and diabetes cases validated using laboratory and medical records. Heart failure was defined using the European Society of Cardiology criteria[27]. Pharmacy data were used to determine the use of Angiotensin Converting Enzyme (ACE) inhibitors (Anatomical Therapeutic Chemical code (ATC) C09A, C09B). Gastro-esophageal reflux disease (GERD) and chronic rhinosinusitis were defined with pharmacy data as a proxy. In particular, participants who received more than two prescriptions of medications for acid-related disorders such as peptic ulcer or reflux disease (ATC A02B) were considered to have GERD. Chronic rhinosinusitis was also defined as having received at least three prescriptions of nasal steroids (ATC R01AD) within one year before baseline.

Statistical analysis

Baseline characteristics were presented with descriptive statistics, and compared between participants according to prevalent chronic cough status. Normally distributed variables were presented as means with standard deviations and compared using independent t-tests. A

Mann-Whitney tests were performed for skewed continuous variables and their median and interquartile range were reported. Categorical data were compared using Chi-square test.

We performed active imputation of 874 (14.9%) missing BMI data using fully conditional specification (10 imputed datasets). Prior to performing linear regression analyses, the CES-D scores were square-root normalized and standardized to unit variance using the z-transformation. To assess the cross-sectional association of chronic cough with depressive symptoms (CES-D) at baseline, we performed a linear regression which was adjusted for age and sex (Model 1), and additionally for BMI, smoking, chronic rhinosinusitis, GERD, obstructive airway disease, lung cancer, diabetes, and heart failure (Model 2). Analyses were repeated after stratification for prevalent respiratory conditions (chronic rhinosinusitis, obstructive airway disease, and lung cancer). Next, we studied the association between chronic cough and clinically relevant depressive symptoms at baseline using logistic regression in the sex-age adjusted model (Model 1) and the confounder adjusted model (Model 2).

The longitudinal association between chronic cough and incident depression (including depressive symptoms, depressive syndrome and major depressive disorder to maximize power) was estimated using Cox regression and adjusted for age and sex (Model 1), and additionally for Model 2 plus history of depressive events (Model 2a) in all eligible participants. The analyses were repeated and stratified for history of depression in the past 5 years based on the incident data (yes/no) since history of depression might confound the risk of recurrent depression. We additionally adjusted for baseline CES-D scores (Model 3) in the stratified analysis.

All statistical analyses were performed using SPSS statistical software version 24 (IBM SPSS Statistics for Windows; IBM Corp, Armonk, New York, USA). Statistical significance was set at a p-value of < 0.05 .

Results

Baseline characteristics

A total of 5,877 participants were included in this study (Table 1). Participants were mainly female (n=3,460, 58.9%), predominantly elderly (mean age 72.3 ± 7.9 years), 13.9% (n=818) were current smokers, and 23.5% (n=1,381) used ACE inhibitors. At baseline, 32.1% (n=1884) of the participants had at least one comorbidity. Within the sample, the prevalences of comorbidities were as follows: 14.4% (n=848) for asthma and/or COPD, 14.5% (n=850) for GERD, 0.3% (n=20) for lung cancer, 2.6% (n=152) for chronic rhinosinusitis, 2.6% (n=153) for diabetes, and 5.7% (n=336) for heart failure. Nearly 13% (n=755) of the participants reported clinically relevant depressive symptoms, whereas 21.9% (n=1,285) had previously suffered from depression.

Approximately 12% of the participants had chronic cough at baseline (n=729). Compared to participants without chronic cough (n=5,148), those with chronic cough were more often current smokers (25.2% vs. 12.3%, $p<0.001$), and had more comorbidities (51.9% vs. 29.3%, $p<0.001$) such as chronic rhinosinusitis (4.4% vs. 2.3%, $p=0.001$), GERD (18.4% vs. 13.9%, $p=0.001$), obstructive airway disease (34.7% vs. 11.6%, $p<0.001$), lung cancer (1.2% vs. 0.2%, $p<0.001$), and heart failure (8.6% vs. 5.3%, $p<0.001$).

Chronic cough and baseline depressive symptoms

Participants with chronic cough reported a 0.25 SD higher depressive symptoms score than those without chronic cough after adjusting for age and sex (adjusted standardized mean difference 0.25, 95% Confidence Interval (CI) 0.18 – 0.33). The association remained significant after further adjustment for BMI, smoking, chronic rhinosinusitis, GERD, obstructive airway disease, lung cancer, diabetes, and heart failure (adjusted standardized mean difference 0.15, 95% CI 0.08 – 0.22); (see table 2). This suggests that persons with

CES-D scores of 4, 17, and 40 would have 1, 2, and 3 points higher CES-D scores, respectively, if they had chronic cough, independent of any related medical conditions. Stratified analyses demonstrated that chronic cough was associated with more depressive symptoms in participants without respiratory disease (adjusted standardized mean difference 0.19, 95% CI 0.10 – 0.28), but not in those with respiratory disease (adjusted standardized mean difference 0.06, 95% CI -0.08 – 0.19) (Supplemental table 1). Persons with CES-D scores of 3, 11, and 24 would have 1, 2, and 3 points higher CES-D scores, respectively, if they had chronic cough not explained by a respiratory disease.

Chronic cough was also significantly associated with more clinically relevant depressive symptoms after adjusting for age and sex (Odds Ratio (OR) 1.39, 95% CI 1.12 – 1.73), but the effect size decreased after further adjustment for BMI, smoking, chronic rhinosinusitis, GERD, obstructive airway disease, lung cancer, diabetes, and heart failure (OR 1.12, 95% CI 0.89 – 1.41); (see table 2). Stratified analyses showed that chronic cough was associated with more clinically relevant depressive symptoms in participants who did not have respiratory disease (OR 1.35, 95% CI 1.02 – 1.77), but not in those with respiratory disease (OR 0.78, 95% CI 0.52 – 1.17). (Supplemental table 2).

Chronic cough and risk of developing depression

Over a total observation period of 37,287 person-years, 900 out of 5,799 participants (with complete follow-up data) developed depressive symptoms (n=670), a depressive syndrome (n=112) or a major depressive disorder (n=118). Chronic cough was significantly associated with an increased risk of any event of depression after adjustment for confounders (Hazard Ratio (HR) 1.22, 95% CI 1.01 – 1.46, table 3).

However, stratification for a history of depression demonstrated that chronic cough was only associated with an increased risk of recurrent depression in participants with a history of

depression (HR 1.45, 95% CI 1.13 – 1.84), see table 3. In contrast, chronic cough was not associated with an increased risk of depression in participants without history of depression at baseline (adjusted HR 0.91, 95% CI 0.68 – 1.22, table 3, figure 2).

Discussion

In this prospective population-based study of middle-aged and older persons, we observed that chronic cough was cross-sectionally associated with more depressive symptoms. Moreover, chronic cough increased the risk of depressive events in persons with a history of depression, but not in persons without a history of depression.

Our findings corroborate previous studies indicating that persons with chronic cough have a substantially higher burden of depressive symptoms[9, 13], independent of commonly associated risk factors and comorbidities such as smoking and asthma[5, 15, 16]. In our study, chronic cough was also associated with higher burden of depressive symptoms after adjusting for preexisting respiratory diseases including chronic rhinosinusitis, asthma, COPD, and lung cancer, albeit with smaller effect sizes. This is line with previous research. For example, Heo *et al* found that, compared to persons without chronic cough, patients with chronic cough had more severe depressed mood regardless of their smoking and lung function status[15].

Our stratified analyses showed that chronic cough is associated with depressive symptoms, particularly in persons whose chronic cough is not explained by a known respiratory disease. Although this might be in part explained by more power in the analyses including those with chronic cough not explained by respiratory disease, we also see substantially larger effect sizes for the association of chronic cough not explained by a known respiratory disease and depression. This is consistent with the findings of Hulme *et al*, who examined the psychological profiles of patients presenting with chronic cough and observed that individuals without a pulmonary disease had significantly more depressive symptoms than those with explainable causes of chronic cough as well as those without cough[16]. Chronic cough is refractory in approximately 50% of patients with treatable traits for cough[29] and

remains unexplained in about 40% of affected persons[30]. We might speculate that some patients may suffer psychological distress due to suboptimal response to cough treatment and lack of definitive diagnosis despite several health visits for cough.

Chronic cough also increased the risk of depression in our study participants, but only in those with a history of depression. Studies have shown that recurrent episodes of depressive disorders have a predilection for clinico-demographic factors relevant in chronic cough[18], and the overall disease burden of chronic cough negatively impacts the risk of depression[8, 10, 12, 14, 31]. Although we were unable to assess any bidirectional effects in this study, we hypothesize that, while depression is more common in chronic cough, it is usually mild rather than severe, and that chronic cough might only lead to depression in those who are already more vulnerable to depression or exacerbate existing depressive symptoms.

Plausible explanation for chronic cough as a risk factor for depression could be neurobiological; for example patients with cough hypersensitivity show increased neural activity in periaqueductal gray[32, 33], a region involved in emotional processing of stress and pain, and also implicated in chronic depression[33, 34]. Furthermore, the psychological determinants of chronic cough[17] and its associated negative social effects[35] highlight the potential concerted role of psychosocial factors on depressive events in chronic cough patients. In particular, cough-related psychosocial problems, such as altered social identity, social embarrassment and isolation, preoccupation and attentional focus, obsessive and unsuccessful attempts to suppress cough,[35, 36] have a negative impact on depression[16]. Indeed, these psychosocial factors may play a role in mediating the pathological link between chronic cough and depression. Another potential connection between chronic cough and depression is impaired modulation of excitatory and inhibitory neurotransmission in the brain[32, 37]. Indeed, centrally acting neuromodulators such as gabapentin and opioids are effective therapeutic options in both conditions[2, 38, 39]. Notably, during cough therapy,

there is a concurrent reduction in cough intensity and depressive symptoms[9, 13]. Therefore, recognizing and treating both conditions may improve the overall clinical management outcomes and patient satisfaction.

Our study has several strengths. The results were based on a large sample from the general population, and the prospective longitudinal assessment of depressive events was based on a combination of questionnaires, clinical interviews and medical records. However, some limitations need to be considered. First, we lacked the power to further explore the impact of chronic cough on specific incident depressive events. Second, although we have information on history of depression up till 5 years before the baseline assessment, these participants might have suffered from depression before this period. Third, recurrent bouts of depression are not unusual in people with a history of depression, and they may be more easily recognized by a GP who is often seen for a complaint of chronic cough. Fourth, our population-based study included older adults and elderly participants whose clinical profiles may differ from those of patients presenting to health care facilities. Consequently, our findings may not be generalizable to hospital-based cohorts. Fifth, chronic pain might be a major confounding factor in the chronic cough-depression link[40]. However, because we did not have chronic pain data during this study period in the Rotterdam Study, we could not adjust for it in our analyses. Sixth, the overlapping symptoms of chronic cough and depression may artificially have inflated the CES-D score. Furthermore, we acknowledge the potential confounding effect of female sex on the association of chronic cough and depressive symptoms and have adjusted for it in our analyses. Lastly, we were unable to assess potential bi-directionality of the association, that is including the potential effects of depression on chronic cough.

In conclusion, middle-aged and older persons with chronic cough have a disproportionate burden of depressive symptoms, particularly those with chronic cough not explained by a

respiratory disease (chronic rhinosinusitis, obstructive airway disease, and lung cancer). Moreover, in individuals who have a history of depression, chronic cough increases the risk of recurrent depressive events. This emphasizes the importance of screening for depression in patients with chronic cough.

References

1. Song WJ, Chang YS, Faruqi S, Kim JY, Kang MG, Kim S, Jo EJ, Kim MH, Plevkova J, Park HW, Cho SH, Morice AH. The global epidemiology of chronic cough in adults: a systematic review and meta-analysis. *Eur Respir J* 2015; 45(5): 1479-1481.
2. Morice AH, Millqvist E, Bieksiene K, Birring SS, Dicpinigaitis P, Domingo Ribas C, Hilton Boon M, Kantar A, Lai K, McGarvey L, Rigau D, Satia I, Smith J, Song W-J, Tonia T, van den Berg JWK, van Manen MJG, Zacharasiewicz A. ERS guidelines on the diagnosis and treatment of chronic cough in adults and children. *European Respiratory Journal* 2020; 55(1): 1901136.
3. Schappert SM. National Ambulatory Medical Care Survey: 1991 summary. *Adv Data* 1993(230): 1-16.
4. Chung KF, Pavord ID. Prevalence, pathogenesis, and causes of chronic cough. *Lancet* 2008; 371(9621): 1364-1374.
5. Sohn KH, Song WJ, Kim SH, Jang HC, Kim KW, Chang YS. Chronic cough, not asthma, is associated with depression in the elderly: a community-based population analysis in South Korea. *Korean J Intern Med* 2019; 34(6): 1363-1371.
6. Koo H-K, Park S-W, Park J-W, Choi HS, Kim T-H, Yoon HK, Yoo KH, Jung K-S, Kim DK. Chronic cough as a novel phenotype of chronic obstructive pulmonary disease. *International journal of chronic obstructive pulmonary disease* 2018; 13: 1793-1801.
7. Çolak Y, Afzal S, Lange P, Laursen LC, Nordestgaard BG, Dahl M. Role and Impact of Chronic Cough in Individuals with Asthma From the General Population. *J Allergy Clin Immunol Pract* 2019; 7(6): 1783-1792 e1788.
8. Irwin RS, Dudiki N, French CL, Panel CEC. Life-Threatening and Non-Life-Threatening Complications Associated With Coughing: A Scoping Review. *Chest* 2020.
9. French CL, Crawford SL, Bova C, Irwin RS. Change in Psychological, Physiological, and Situational Factors in Adults After Treatment of Chronic Cough. *Chest* 2017; 152(3): 547-562.
10. French CL, Irwin RS, Curley FJ, Krikorian CJ. Impact of chronic cough on quality of life. *Arch Intern Med* 1998; 158(15): 1657-1661.
11. Chamberlain SA, Garrod R, Douiri A, Masefield S, Powell P, Bücher C, Pandyan A, Morice AH, Birring SS. The impact of chronic cough: a cross-sectional European survey. *Lung* 2015; 193(3): 401-408.
12. Dicpinigaitis P. Prevalence of urinary incontinence in women seeking evaluation for chronic cough. *European Respiratory Journal* 2020; 56(suppl 64): 1286.
13. Dicpinigaitis PV, Tso R, Banauch G. Prevalence of depressive symptoms among patients with chronic cough. *Chest* 2006; 130(6): 1839-1843.
14. Adams RJ, Appleton SL, Wilson DH, Taylor AW, Ruffin RE. Associations of physical and mental health problems with chronic cough in a representative population cohort. *Cough (London, England)* 2009; 5: 10-10.
15. Heo IR, Kim J-Y, Go S-I, Kim TH, Ju SM, Yoo J-W, Lee SJ, Cho YJ, Jeong YY, Lee JD, Kim HC. Chronic Cough is Associated with Depressive Mood in Women Regardless of Smoking Status and Lung Function. *The Clinical Respiratory Journal* 2021; n/a(n/a).
16. Hulme K, Deary V, Dogan S, Parker SM. Psychological profile of individuals presenting with chronic cough. *ERJ Open Research* 2017; 3(1): 00099-02016.
17. Vertigan AE. Somatic cough syndrome or psychogenic cough-what is the difference? *Journal of thoracic disease* 2017; 9(3): 831-838.
18. Burcusa SL, Iacono WG. Risk for recurrence in depression. *Clin Psychol Rev* 2007; 27(8): 959-985.

19. Won HK, Lee JH, An J, Sohn KH, Kang MG, Kang SY, Morice AH, Cho SH, Song WJ. Impact of Chronic Cough on Health-Related Quality of Life in the Korean Adult General Population: The Korean National Health and Nutrition Examination Survey 2010-2016. *Allergy Asthma Immunol Res* 2020; 12(6): 964-979.
 20. Johansson H, Johannessen A, Holm M, Forsberg B, Schlünssen V, Jøgi R, Clausen M, Lindberg E, Malinovski A, Emilsson Ö I. Prevalence, progression and impact of chronic cough on employment in Northern Europe. *Eur Respir J* 2021; 57(6).
 21. Ikram MA, Brusselle G, Ghanbari M, Goedegebure A, Ikram MK, Kavousi M, Kieboom BCT, Klaver CCW, de Knecht RJ, Luik AI, Nijsten TEC, Peeters RP, van Rooij FJA, Stricker BH, Uitterlinden AG, Vernooij MW, Voortman T. Objectives, design and main findings until 2020 from the Rotterdam Study. *Eur J Epidemiol* 2020; 35(5): 483-517.
 22. Arinze JT, de Roos EW, Karimi L, Verhamme KMC, Stricker BH, Brusselle GG. Prevalence and incidence of, and risk factors for chronic cough in the adult population: the Rotterdam Study. *ERJ Open Res* 2020; 6(2).
 23. Beekman AT, van Limbeek J, Deeg DJ, Wouters L, van Tilburg W. [A screening tool for depression in the elderly in the general population: the usefulness of Center for Epidemiological Studies Depression Scale (CES-D)]
- Een screeningsinstrument voor depressie bij ouderen in de algemene bevolking: de bruikbaarheid van de Center for Epidemiologic Studies Depression Scale (CES-D). *Tijdschr Gerontol Geriatr* 1994; 25(3): 95-103.
24. Radloff LS. The CES-D Scale: A Self-Report Depression Scale for Research in the General Population. *Applied Psychological Measurement* 1977; 1(3): 385-401.
 25. Wing JK, Babor T, Brugha T, Burke J, Cooper JE, Giel R, Jablenski A, Regier D, Sartorius N. SCAN. Schedules for Clinical Assessment in Neuropsychiatry. *Arch Gen Psychiatry* 1990; 47(6): 589-593.
 26. Segal DL. Diagnostic and Statistical Manual of Mental Disorders (DSM-IV-TR). The Corsini Encyclopedia of Psychology; pp. 1-3.
 27. Bleumink GS, Knetsch AM, Sturkenboom MC, Straus SM, Hofman A, Deckers JW, Wittman JC, Stricker BH. Quantifying the heart failure epidemic: prevalence, incidence rate, lifetime risk and prognosis of heart failure The Rotterdam Study. *Eur Heart J* 2004; 25(18): 1614-1619.
 28. Ohno S, Takahashi K, Inoue A, Takada K, Ishihara Y, Tanigawa M, Hirao K. Smallest detectable change and test-retest reliability of a self-reported outcome measure: Results of the Center for Epidemiologic Studies Depression Scale, General Self-Efficacy Scale, and 12-item General Health Questionnaire. *J Eval Clin Pract* 2017; 23(6): 1348-1354.
 29. Ryan NM, Vertigan AE, Birring SS. An update and systematic review on drug therapies for the treatment of refractory chronic cough. *Expert Opin Pharmacother* 2018; 19(7): 687-711.
 30. Haque RA, Usmani OS, Barnes PJ. Chronic idiopathic cough: a discrete clinical entity? *Chest* 2005; 127(5): 1710-1713.
 31. Johansson H, Johannessen A, Holm M, Forsberg B, Schlünssen V, Jøgi R, Clausen M, Lindberg E, Malinovski A, Emilsson ÖI. Prevalence, progression and impact of chronic cough on employment in Northern Europe. *European Respiratory Journal* 2021; 57(6): 2003344.
 32. Ando A, Smallwood D, McMahon M, Irving L, Mazzone SB, Farrell MJ. Neural correlates of cough hypersensitivity in humans: evidence for central sensitisation and dysfunctional inhibitory control. *Thorax* 2016; 71(4): 323.

33. Buhle JT, Kober H, Ochsner KN, Mende-Siedlecki P, Weber J, Hughes BL, Kross E, Atlas LY, McRae K, Wager TD. Common representation of pain and negative emotion in the midbrain periaqueductal gray. *Soc Cogn Affect Neurosci* 2013; 8(6): 609-616.
34. Ho YC, Lin TB, Hsieh MC, Lai CY, Chou D, Chau YP, Chen GD, Peng HY. Periaqueductal Gray Glutamatergic Transmission Governs Chronic Stress-Induced Depression. *Neuropsychopharmacology* 2018; 43(2): 302-312.
35. Hulme K, Dogan S, Parker SM, Deary V. 'Chronic cough, cause unknown': A qualitative study of patient perspectives of chronic refractory cough. *J Health Psychol* 2019; 24(6): 707-716.
36. Van den Bergh O, Van Diest I, Dupont L, Davenport PW. On the psychology of cough. *Lung* 2012; 190(1): 55-61.
37. Sanacora G, Gueorguieva R, Epperson CN, Wu YT, Appel M, Rothman DL, Krystal JH, Mason GF. Subtype-specific alterations of gamma-aminobutyric acid and glutamate in patients with major depression. *Arch Gen Psychiatry* 2004; 61(7): 705-713.
38. Yovell Y, Bar G, Mashiah M, Baruch Y, Briskman I, Asherov J, Lotan A, Rigbi A, Panksepp J. Ultra-Low-Dose Buprenorphine as a Time-Limited Treatment for Severe Suicidal Ideation: A Randomized Controlled Trial. *Am J Psychiatry* 2016; 173(5): 491-498.
39. Yasmin S, Carpenter LL, Leon Z, Siniscalchi JM, Price LH. Adjunctive gabapentin in treatment-resistant depression: a retrospective chart review. *J Affect Disord* 2001; 63(1-3): 243-247.
40. Arinze JT, Verhamme KMC, Luik AI, Stricker B, van Meurs JBJ, Brusselle GG. The interrelatedness of chronic cough and chronic pain. *Eur Respir J* 2021; 57(5).

Funding/ acknowledgement

This study was supported by Merck Sharp & Dohme (MSD) research grant for investigator-initiated study.

Table 1: Baseline Characteristics of the study population

Baseline Characteristics	Total (n = 5,877)	No chronic cough (n = 5,148)	Chronic cough (n = 729)	p-value
Age, mean years $\pm$ SD	72.3 $\pm$ 7.9	72.2 $\pm$ 7.9	72.7 $\pm$ 7.7	0.081
Female, <i>n</i> (%)	3,460 (58.9)	3,031 (58.9)	429 (58.8)	0.988
BMI, mean kg/m ² $\pm$ SD	27.6 $\pm$ 4.1	27.6 $\pm$ 4.0	27.6 $\pm$ 4.4	0.645
Smoking, <i>n</i> (%)				
▪ Never	1,876 (31.9)	1,693 (32.9)	183 (25.1)	<0.001
▪ Past	3,180 (54.1)	2,818 (54.8)	362 (49.7)	
▪ Current	818 (13.9)	634 (12.3)	184 (25.2)	
ACE inhibitor use, <i>n</i> (%)	1,381 (23.5)	1,153 (22.4)	228 (31.3)	<0.001
CESD scores, median (<i>IQR</i>)	4 (1 – 10)	4 (1 – 9)	6 (2 – 12)	<0.001
CESD score, score ≥ 16 (%)	755 (12.9)	636 (12.4)	119 (16.4)	0.002
History of depressive events, <i>n</i> (%)	1,285 (21.9)	1,095 (21.3)	190 (26.1)	0.003
Baseline comorbidities, <i>n</i> (%)	1,884 (32.1)	1,506 (29.3)	378 (51.9)	<0.001
▪ Chronic rhinosinusitis	152 (2.6)	120 (2.3)	32 (4.4)	0.001
▪ GERD	850 (14.5)	716 (13.9)	134 (18.4)	0.001
▪ Obstructive airway disease	848 (14.4)	595 (11.6)	253 (34.7)	<0.001
▪ Lung cancer	20 (0.3)	11 (0.2)	9 (1.2)	<0.001
▪ Diabetes	153 (2.6)	133 (2.6)	20 (2.7)	0.800
▪ Heart failure	336 (5.7)	273 (5.3)	63 (8.6)	<0.001
SD – Standard deviation; IQR – Interquartile range; BMI – Body mass index; GERD – Gastroesophageal reflux disease; ACE – Angiotensin converting enzyme; CESD - Center for Epidemiological Studies Depression Scale; COPD – Chronic obstructive pulmonary disease; Obstructive airway disease – COPD and/or asthma				

Table 2: The cross-sectional association between chronic cough and depressive symptoms at baseline (n=5,852)

	CES-D scores median (IQR)	Model 1 standardized mean adjusted difference (95% CI) in depressive symptoms at baseline	Model 2 standardized mean adjusted difference (95% CI) in depressive symptoms at baseline
No chronic cough (n=5,128)	4 (1 – 9)	Ref.	Ref.
Chronic cough (n=724)	6 (2 – 12)	0.25 (0.18 – 0.33)	0.15 (0.08 – 0.22)
	CES-D score ≥ 16 (n=755, 12.9%)	Model 1 OR (95% CI)	Model 2 OR (95% CI)
No chronic cough (n=5,128)	636 (12.4%)	Ref.	Ref.
Chronic cough (n=724)	119 (16.4%)	1.39 (1.12 – 1.73)	1.12 (0.89 – 1.41)
CES-D scores were square root-transformed and z-standardized; Model 1 – adjusted for age and sex; Model 2 – adjusted for age, sex, BMI, smoking, chronic rhinosinusitis, GERD, obstructive airway disease, lung cancer, diabetes, and heart failure. 25 participants did not have CES-D score data.			

Table 3: The longitudinal association between chronic cough and incident/ recurrent depression

All eligible participants regardless of history of depression (n=5,799)	Incident depression (n=900)	Model 1 HR (95% CI)	Model 2 HR (95% CI)	Model 2a HR (95% CI)
No chronic cough (n=5,080)	758	Ref.	Ref.	Ref.
Chronic cough (n=719)	142	1.40 (1.17 – 1.68)	1.22 (1.01 – 1.46)	1.22 (1.01 – 1.46)
Participants without a history of depression (n=4,516)	Incident depression (n=466)	Model 1 HR (95% CI)	Model 2 HR (95% CI)	Model 3 HR (95% CI)
No chronic cough (n=3,987)	408	Ref.	Ref.	Ref.
Chronic cough (n=529)	58	1.11 (0.84 – 1.46)	0.96 (0.73 – 1.28)	0.91 (0.68 – 1.22)
Participants with a history of depression (n=1,283)	Recurrent depression (n=434)	Model 1 HR (95% CI)	Model 2 HR (95% CI)	Model 3 HR (95% CI)
No chronic cough (n=1,093)	350	Ref.	Ref.	Ref.
Chronic cough (n=109)	84	1.53 (1.21 – 1.95)	1.46 (1.14 – 1.86)	1.45 (1.13 – 1.84)
Model 1 – adjusted for age and sex; Model 2 – adjusted for age, sex, BMI, smoking, chronic rhinosinusitis, GERD, obstructive airway disease, diabetes, and heart failure; Model 2a – adjusted for age, sex, BMI, smoking, chronic rhinosinusitis, GERD, obstructive airway disease, diabetes, heart failure, and history of depressive event; Model 3 – adjusted for age, sex, BMI, smoking, chronic rhinosinusitis, GERD, obstructive airway disease, diabetes, heart failure, and baseline depressive symptoms (CESD scores). Depression includes any event of depressive symptoms, depressive syndromes and major depressive disorder. 78 participants had no follow-up data on incident/recurrent depression.

STROBE Study Flow chart (RS-I RS-II)

Summary of risk of incident/ recurrent depression among study participants

Supplemental Table 1: The cross-sectional association between chronic cough and depressive symptoms at baseline (stratified)

Participants with respiratory disease (n=950)	CES-D scores median (IQR)	Model 1 standardized mean adjusted difference (95% CI) in depressive symptoms at baseline	Model 2 standardized mean adjusted difference (95% CI) in depressive symptoms at baseline
No chronic cough (n=680)	5 (2 – 12)	Ref.	Ref.
Chronic cough (n=270)	7 (3 – 13)	0.12 (-0.02 – 0.25)	0.06 (-0.08 – 0.19)
Participants without respiratory disease (n = 4,902)	CES-D scores median (IQR)	Model 1 standardized mean adjusted difference (95% CI) in depressive symptoms at baseline	Model 2b standardized mean adjusted difference (95% CI) in depressive symptoms at baseline
No chronic cough (n=4,448)	4 (1 – 9)	Ref.	Ref.
Chronic cough (n=454)	5 (2 – 11)	0.24 (0.14 – 0.33)	0.19 (0.10 – 0.28)
CES-D scores were square root-transformed and z-standardized; Respiratory disease= chronic rhinosinusitis, obstructive airway disease, or lung cancer; Model 1 – adjusted for age and sex; Model 2 – adjusted for age, sex, BMI, smoking, chronic rhinosinusitis, GERD, obstructive airway disease, lung cancer, diabetes, and heart failure; Model 2b – adjusted for age, sex, BMI, smoking, GERD, diabetes, and heart failure. 25 participants did not have CES-D score data.			

Supplemental Table 2: The cross-sectional association between chronic cough and clinically relevant depressive symptoms at baseline (stratified)

Participants with respiratory disease (n=950)	CES-D score $\geq$ 16 (n=165, 17.4%)	Model 1 OR (95% CI)	Model 2 OR (95% CI)
No chronic cough (n=680)	121 (17.8%)	Ref.	Ref.
Chronic cough (n=270)	44 (16.3%)	0.89 (0.60 – 1.31)	0.78 (0.52 – 1.17)
Participants without respiratory disease (n = 4,902)	CES-D score $\geq$ 16 (n=590, 12.0%)	Model 1 OR (95% CI)	Model 2b OR (95% CI)
No chronic cough (n=4,448)	515 (11.6%)	Ref.	Ref.
Chronic cough (n=454)	75 (16.5%)	1.49 (1.14 – 1.96)	1.35 (1.02 – 1.77)
Model 1 – adjusted for age and sex; Model 2 – adjusted for age, sex, BMI, smoking, chronic rhinosinusitis, GERD, obstructive airway disease, lung cancer, diabetes, and heart failure; Model 2b – adjusted for age, sex, BMI, smoking, GERD, diabetes, and heart failure. 25 participants did not have CES-D score data.			