



Early View

Review

COVID-19 Therapies for inpatients: a review and quality assessment of clinical guidelines

David A. Wohl, Aprille A. Espinueva, Lauren Dau, Chen-Yu Wang, Alexandra Lachmann, Rujuta A. Bam, Aaditya Rawal, Kerris Chappell-Smith, Juergen K. Rockstroh

Please cite this article as: Wohl DA, Espinueva AA, Dau L, *et al.* COVID-19 Therapies for inpatients: a review and quality assessment of clinical guidelines. *ERJ Open Res* 2022; in press (<https://doi.org/10.1183/23120541.00236-2022>).

This manuscript has recently been accepted for publication in the *ERJ Open Research*. It is published here in its accepted form prior to copyediting and typesetting by our production team. After these production processes are complete and the authors have approved the resulting proofs, the article will move to the latest issue of the ERJOR online.

Copyright ©The authors 2022. This version is distributed under the terms of the Creative Commons Attribution Non-Commercial Licence 4.0. For commercial reproduction rights and permissions contact permissions@ersnet.org

COVID-19 Therapies for Inpatients: A Review and Quality Assessment of Clinical Guidelines

David A. Wohl (MD),^{1,2} Aprille A. Espinueva (PharmD),³ Lauren Dau (PharmD),³ Chen-Yu Wang (PhD),³ Alexandra Lachmann (MSPAS),³ Rujuta A. Bam (MS),³ Aaditya Rawal (MSc),⁴ Kerris Chappell-Smith (BA),⁵ Juergen K. Rockstroh (MD)^{6,7}

¹Gillings School of Global Public Health, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA; ²School of Medicine, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA; ³Gilead Sciences Inc., Foster City, CA, USA; ⁴Costello Medical, Boston, MA, USA; ⁵Costello Medical, Cambridge, UK; ⁶Department of Internal Medicine, University Hospital Bonn, Bonn, Germany; ⁷German Centre for Infection Research, Partner Site Cologne-Bonn, Bonn, Germany.

Correspondence to: David A. Wohl, wohl@med.unc.edu, (919) 843-2723

Short title: COVID-19 Guidelines Review

Trial registration: N/A

Funding: Gilead Sciences Inc.

Key words: COVID-19; Guideline; AGREE-II; baricitinib; dexamethasone; remdesivir; casirivimab/imdevimab; tocilizumab

SUMMARY: Take Home Message (256 character limit [including spaces])

This targeted review of COVID-19 treatment guidelines aimed to understand the heterogeneity in quality, recommendations, and evidence bases for therapies commonly used among COVID-19 inpatients

ABSTRACT

Due to condensed development processes, expanding evidence and differences in healthcare system characteristics, many COVID-19 guidelines differ in their quality and treatment recommendations, which has consequences for clinical practice. This review aimed to identify COVID-19 treatment guidelines, assess their quality, and summarise their recommendations. Guidelines were identified for five therapies most commonly used among inpatients with COVID-19 (remdesivir, dexamethasone, tocilizumab, baricitinib, and casirivimab/imdevimab) from 11 countries. Guideline quality was assessed using the Appraisal of Guidelines for Research and Evaluation II (AGREE-II) tool. Full details of recommendations and supporting evidence were analysed for high-quality guidelines, defined as those scoring $\geq 50\%$ in Domain 3 (Rigour of Development) of AGREE-II. Overall, guidelines differed substantially in their quality and, even among high-quality guidelines using the same evidence, recommendations regarding specific therapeutics varied. Potential reasons for this heterogeneity, including the availability and consistency of clinical data, visibility of trial endpoints, and context-specific factors, are discussed.

INTRODUCTION

The clinical research response to the novel severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) pandemic has been unprecedented, rapidly yielding highly effective and safe vaccines, as well as data that guide the use of therapeutics across the spectrum of coronavirus disease-2019 (COVID-19).[1, 2] Monoclonal antibodies (mAbs), direct-acting antiviral therapies, corticosteroids, interleukin-6 (IL-6) antagonists, and Janus Kinase (JAK) inhibitors are commonly used to treat patients with COVID-19, following clinical trial results indicating their benefit in certain populations.[3-7] Yet, with the rapid generation of large amounts of data and sometimes conflicting clinical results, deciding on the best practice has been challenging.

To provide clinicians with evidence-based recommendations for managing patients with COVID-19 in their regions, many governmental or scientific organizations (both national and international) developed clinical treatment guidelines.[8-10] However, to deliver recommendations in a timely manner, guideline development groups often had to use rapid, condensed and less meticulous methods, reducing a process that often takes years to just a few weeks. Premature or draft datasets were sometimes utilised and, with an ever-expanding body of clinical data, evidence to support recommendations may have become outdated. Further, variable treatment availability and healthcare system characteristics could have led to differences in preferences or priorities. Such challenges have given rise to guidelines that are heterogeneous in their recommendations, quality and rigour; this can lead to confusion and uncertainty among clinicians, and disparate management practices around the globe.[11, 12] There is a need to identify high-quality COVID-19 treatment guidelines and provide clinicians with a synthesis of recommendations, especially those applicable to hospital settings.

Using a targeted literature review approach, this *de novo* review aims to: a) identify treatment guidelines for therapies most commonly used in patients hospitalised with COVID-19, b) systematically assess the quality of included guidelines, and c) compare the recommendations, evidence sources, and considered outcomes of guidelines deemed to be of high-quality.

METHODS

Scope of Review

COVID-19 treatment guidelines applicable to hospital settings were reviewed, with a focus on five key therapies used to treat COVID-19 in hospitalised patients: remdesivir, dexamethasone, tocilizumab, baricitinib, and casirivimab/imdevimab. While some of these therapies may be used to treat non-hospitalised patients, this review focused only on their use in hospital settings.

This review focused on guidelines from governmental or scientific organizations within countries that had the resource capacity and the continued need (due to recent epidemiologic disease burden) to develop and regularly update COVID-19 treatment guidelines. To this end, countries with the largest economies in their regions (within the top four highest 2020 total gross domestic product in the Americas, Asia, Europe, and the Middle East and Africa)[13] as well as the most severe, recent impact of COVID-19 (within the top 50 highest reported bi-weekly COVID-19 deaths globally, as of January 1, 2021) were identified.[14] Using these criteria, 11 countries were prioritised: Brazil, Canada, France, Germany, India, Italy, Japan, Mexico, Turkey, United Kingdom (UK), and the United States (US). Guidelines from specific regional or international organizations with the resources and relevant influence to develop and update guidelines were also included. These comprised the World Health Organization (WHO), the European Respiratory Society (ERS), and the Surviving Sepsis Campaign (SSC). Initial searches to identify guidelines were conducted in August 2021; included guidelines were screened for updates on January 4, 2022.

Search strategy and selection criteria

Due to rapid development and regular updating, the most recent COVID-19 treatment guidelines are often not published in journals or indexed in electronic databases. Therefore, targeted searches of continually updated websites, such as guideline repositories and libraries, were conducted (**Supplementary Table 1**).

Following an initial visual search, each source was queried using search terms tailored to the functionality, specificity and language of each source (**Supplementary Table 2; Supplementary Figure 1**). Initial searches were conducted in August 2021.

Identified guidelines were screened against eligibility criteria (**Supplementary Table 3; Supplementary Figure 2**) and prioritised for inclusion. Priority was given to guidelines which: were applicable to the broadest possible patient population and to an entire country or region, assessed multiple therapies of interest, and directly assessed clinical data (rather than summarizing information from other guidelines).

Quality Assessments and Data Extraction

The quality of each included guideline was assessed by two reviewers using the Appraisal of Guidelines for Research and Evaluation II (AGREE-II) instrument.[15] Designed by an international team of guideline developers and researchers, AGREE-II has been used to rigorously assess the quality of a broad range of treatment guidelines, including those for infectious diseases.[16-20] The instrument includes 23 items across six domains: Domain 1: Scope and Purpose; Domain 2: Stakeholder Involvement; Domain 3: Rigour of Development; Domain 4: Clarity of Presentation; Domain 5: Applicability; and Domain 6: Editorial Independence (**Supplementary Table 4**).[15] For each item, a score from 1 (strongly disagree) to 7 (strongly agree) was allocated. Any differences between reviewers of ≥ 2 points were resolved by discussion. A standardised domain score between 0% and 100% (low to high quality) was then calculated for each of the six domains.

For guidelines considered high-quality, the full details of recommendations were extracted by a single reviewer and verified by a second reviewer. High-quality guidelines were defined as those scoring $\geq 50\%$ in Domain 3 (Rigour of Development) which determines whether recommendations are based on robust methodologies and reliable, up-to-date evidence. Similar thresholds of high quality have been used in other published AGREE-II assessments of guidelines.[21-23] Extracted details included the therapy, characteristics of the applicable population (for example, supplemental oxygen requirement, disease severity, oxygen saturation), type of recommendation (for, against, clinical trials only, or insufficient evidence), and the evidence and outcomes used to support recommendations. Recommendations were categorised by the applicable population: hospitalised but do not require supplemental oxygen ("mild" disease); hospitalised and require low-flow supplemental oxygen ("moderate" disease); hospitalised and require non-invasive ventilation (NIV) or high-flow oxygen ("severe" disease); and hospitalised and

require invasive mechanical ventilation (IMV) or extracorporeal membrane oxygenation (ECMO; “critical” disease).

Guideline Updates

Included guidelines were screened on January 4, 2022, for updates since the initial searches in August 2021. Quality assessments were not repeated; however, data were re-extracted for updated guidelines. Where several versions of the same guideline had been published, only the most recent versions were considered.

Role of the Funding Source

Gilead Sciences Inc. provided funding to Costello Medical Inc. to conduct this study, under the direction of all authors. Five authors (AAE, LD, CYW, AL and RAB) are employees of Gilead Sciences Inc., and were involved in the study conception and design, the interpretation of data, in revising and approving the article and in the decision to submit the paper for publication. JKR and DAW were not compensated for their time in conducting this study.

RESULTS

In total, 96 eligible treatment guidelines were identified; of these, 73 did not fulfil the prioritisation criteria, leaving 23 which were prioritised for quality assessment. Three were from international organizations and 20 were country-specific guidelines (**Supplementary Table 5**).

Guideline Quality

There was considerable variability between the overall quality of guidelines, as well as the Domain scores within guidelines (**Supplementary Table 5**). The highest scores were generally achieved in Domain 4 (Clarity of Presentation) while the lowest scores were shown in Domains 5 and 6 (Applicability and Editorial Independence, respectively). A total of 13 guidelines achieved a score of $\geq 50\%$ in Domain 3 (Rigour of Development), meeting the criteria for high-quality and full data extraction; of these, three were international guidelines and ten were country-specific (from nine different countries). A broad geographical range was represented, including international guidelines, and guidelines from Brazil, Canada, Germany, India, Italy, Japan, Mexico, the UK, and the US.[8-10, 24-41] **Supplementary Table 6** provides a list of these 13 high-quality guidelines and a brief overview of their recommendations by therapy and applicable population.

High-Quality Guideline Recommendations by Therapeutic

Remdesivir

All 13 high-quality guidelines discussed remdesivir as monotherapy for use in patients hospitalised with COVID-19. Many also discussed the use of remdesivir within combination regimens. A summary of recommendations, supporting evidence and outcomes considered is presented in **Table 1**.

Of the guidelines that discussed the use of remdesivir in mild (N=9)[8-10, 24, 26, 36, 38, 39] and critical (N=12)[9, 10, 34, 36, 37, 39, 41] disease, most recommended against the use of remdesivir monotherapy (mild: n=6[8, 9, 24, 26, 36, 38]; critical: n=10[9, 10, 24, 26, 34, 36-39, 41]). Others suggested that there was insufficient evidence or that remdesivir (alone or in combination) should only be used in clinical trials[8, 10, 35, 39]. In moderate and severe disease, there was a mix of recommendations regarding remdesivir monotherapy, but remdesivir in combination with other therapies (particularly dexamethasone or baricitinib) was more frequently recommended. The WHO[9], Brazilian Ministry of Health (MoH)[38], and Indian Clinical Infectious Disease Society (CIDS)[26] guidelines did not recommend remdesivir in any disease severity. The ERS international guidelines[10] and the Association of the Scientific Medical Societies in Germany (AWMF)[24] guidelines recommended against remdesivir in some disease severities and reported insufficient evidence for others.

All guidelines, besides those from the Government of Mexico[35], reported considering evidence from the ACTT-1,[5] WHO Solidarity,[42] and Wang 2020[43] trials. No guidelines recommending against remdesivir considered the SIMPLE Severe trial,[44] but three which recommended for remdesivir considered results from this study: COVID-19 Advisory Ontario,[34] Italian Society of Anti-Infective Therapy (SITA)/Italian Society of Pulmonology (SIP),[40] and the National Institute for Health and Care Excellence (NICE)[8]. Only two organizations considered results from Mahajan 2021,[45] both of which recommended against the use of remdesivir in hospitalised patients[24, 26].

Mortality and safety outcomes were considered by most guidelines (mortality: n=12[8-10, 24, 26, 34, 36-41]; safety: n=11[8-10, 24, 26, 35-39, 41]), regardless of recommendation. However, the guidelines that recommended against remdesivir

more frequently considered cost (n=4)[9, 10, 24, 38] and those which recommended for remdesivir more frequently considered clinical recovery (n=7)[8, 34, 36, 37, 39-41]. Regardless of recommendation, few guidelines considered time to discharge (n=5)[8, 9, 26, 34, 38, 39] and time to intensive care unit (ICU) admission (n=2)[34, 38].

Dexamethasone

All 13 high-quality guidelines discussed the use of dexamethasone in patients hospitalised with COVID-19 (**Table 2**); of these, all recommended dexamethasone as monotherapy in severe and/or critical disease while most (n=10)[8-10, 24, 35-39] recommended against its use in mild disease. Many also recommended dexamethasone in combination with other therapies. For example, in the severe and critical populations, several guidelines recommended tocilizumab in combination with dexamethasone if patients had rapidly progressive disease (severe: n=8[8, 10, 24, 27, 33, 35, 36, 40]; critical: n=7[8, 10, 27, 33, 35, 36, 39]). A smaller number of guidelines also recommended remdesivir with dexamethasone (n=2)[8, 39] or baricitinib with dexamethasone (n=3)[27, 36, 39]. The NIH guidelines recommended the use of baricitinib or tocilizumab in combination with remdesivir and dexamethasone.[39]

Except those from the Brazilian MoH,[38] all guidelines considered evidence from the RECOVERY trial (n=12).[3] Most also considered information from DEXA-COVID19 (n=8)[46] and CoDEX (n=10).[47] In addition, several guidelines considered evidence from trials which investigated corticosteroids other than dexamethasone; for example, REMAP-CAP (n=9),[48] CAPE COVID (n=9),[49] COVID STEROID (n=7),[50] and GLUCOCOVID (n=6).[51]

The NIH and ERS guidelines considered the broadest range of outcomes, including several 'other' outcomes such as viral clearance and duration of fever.[10, 39] All guidelines recommending dexamethasone in severe and/or critical disease but against its use in mild disease (n=10)[8-10, 24, 27, 35-39] considered mortality outcomes and most considered safety outcomes. Few guidelines considered clinical recovery (n=1)[37] and time to ICU admission (n=2)[10, 39].

Tocilizumab

Overall, 12 of the 13 high-quality guidelines discussed the use of tocilizumab in patients hospitalised with COVID-19 (**Table 3**); the international SSC guidelines were the only not to discuss its use.[41] Three guidelines recommended against tocilizumab in mild disease.[10, 24, 25] Meanwhile, in severe disease, ten guidelines recommended tocilizumab[8-10, 24, 25, 30, 35, 36, 39, 40]; eight of these explicitly recommended its use in combination (predominantly with dexamethasone).[8, 10, 24, 25, 30, 35, 36, 40] The Brazilian MoH guidelines reported insufficient evidence to formulate a recommendation for the use of tocilizumab (as a monotherapy) in severe COVID-19.[38] In moderate and critical COVID-19, guidelines generally recommended for the use of tocilizumab (n=5[8, 10, 30, 35, 37] and n=8[8-10, 25, 30, 35, 36, 39], respectively), predominantly in combination with dexamethasone. However, two guidelines in each disease severity category recommended against its use as a monotherapy (AWMF and Indian CIDS guidelines recommended against use in moderate COVID-19[24, 25]; AWMF and Brazilian MoH guidelines recommended against its use in severe COVID-19[24, 38]). Seven guidelines specifically recommend tocilizumab in patients who have progressive disease and systemic inflammation (as measured by increased levels of c-reactive protein).[8, 24, 25, 30, 35, 36, 39, 40]

Regardless of recommendation, six different trials (RECOVERY,[3] TOCIBRAS,[52] CORIMUNO- 19,[53] COVACTA,[54] EMPACTA,[55] and Stone 2020[56]) were considered by the large majority of guidelines (n=10).[8, 10, 24, 25, 30, 36-40] The Government of Mexico did not consider any evidence and the WHO guidelines only considered meta-analyses of trials.[9, 35] In terms of evidence considered, there were no distinct differences between guidelines which consistently recommended tocilizumab and those which recommended against its use in some populations.

All guidelines considered mortality and safety outcomes. The majority also considered progression to invasive mechanical ventilation (IMV; n=11)[8-10, 24, 25, 30, 36, 38-40], either as an individual outcome or as part of a composite outcome. Discharge, clinical recovery, and time to ICU admission were each considered by approximately half of the guidelines. There were no clear differences in the outcomes that were considered between guidelines which consistently recommended tocilizumab and those which recommended against its use in some populations.

Baricitinib

Nine high-quality guidelines discussed the use of baricitinib among patients hospitalised with COVID-19 (**Table 4**).[9, 24, 28, 32, 35-37, 39, 40] Five guidelines discussed the use of baricitinib in mild disease, though recommendations were relatively inconsistent.[24, 28, 32, 35, 39] In moderate, severe and critical COVID-19, guidelines generally recommended baricitinib as a monotherapy (n=3[24, 32, 37]; n=3[9, 37, 39]; n=2[9, 32], respectively). Some also recommended its use in combination with dexamethasone and/or remdesivir (n=2[28, 39]; n=4[28, 36, 39, 40]; n=2[28, 39], respectively). The NIH guidelines discussed the use of baricitinib in combination with tocilizumab in severe and critical COVID-19, but recommended against this combination therapy except in clinical trial settings.[39] Across all disease severities, the Government of Mexico guidelines stated that baricitinib should be examined in the context of clinical trials only, preferably in combination with remdesivir.[35]

Evidence from the ACTT-2 trial[6] was considered by all organizations, while COV-BARRIER[57] was considered by all guidelines except those from SITA/SIP.[40]

All nine guidelines considered mortality and safety outcomes and seven (all except the IDSA[36] and Government of Mexico[35] guidelines) considered clinical recovery. No guidelines considered time to ICU admission and only three considered cost. The Government of Mexico guidelines[35], which stated the baricitinib should be examined in clinical trials only, considered the fewest outcomes while CIDS[28] and AWMF[24] considered the broadest range.

Casirivimab/Imdevimab

In total, 10 high-quality guidelines discussed the use of casirivimab/imdevimab in patients hospitalised with COVID-19, though the J-SSCG did not make a clear recommendation (**Table 5**).[8, 9, 24, 29, 31, 35, 37-40] Overall, there was no clear consensus between guidelines regarding casirivimab/imdevimab recommendations. However, within guidelines, the same recommendation was often given across all disease severities for which casirivimab/imdevimab were discussed. AWMF[24] discussed casirivimab/imdevimab in mild and moderate disease, recommending its use in both; the Brazilian MoH guidelines[38] recommended against the use of casirivimab/imdevimab in all four severities; the NICE, COVID-19 Advisory Ontario

and Indian CIDS guidelines[8, 29, 31] recommended its use in seronegative patients only; and the Government of Mexico and J-SSCG guidelines[35, 37] suggested that there was insufficient evidence to formulate a recommendation.

All guidelines which recommended casirivimab/imdevimab considered evidence from the RECOVERY trial[3] and two[9, 24] also considered REGN-COV 2067.[7]

Guidelines recommending against casirivimab/imdevimab did not consider either of these trials.[38-40] The NIH strictly followed guidance in the US Food and Drug Administration (FDA) Emergency Use Authorization (EUA) document that casirivimab/imdevimab should only be used in non-hospitalised patients.

Guidelines which recommended casirivimab/imdevimab more frequently considered mortality, discharge, safety outcomes and progression to IMV than those which recommended against its use. Guidelines which recommended for the use of casirivimab/imdevimab typically considered a broader range of clinical outcomes than those which recommended against its use.

DISCUSSION

During the COVID-19 pandemic, guideline bodies around the globe have attempted to provide clinicians with recommendations for treating patients hospitalised with COVID-19. Assessments using the AGREE-II tool revealed considerable heterogeneity in the quality of the 23 guidelines included in this review. Even within guidelines, there was often substantial variation in quality across AGREE-II Domains, with few guidelines achieving consistently high scores. Overall, 13 guidelines passed the threshold for high quality, scoring $\geq 50\%$ in Domain 3 (Rigour of Development).

For dexamethasone and tocilizumab, recommendations across the 13 high-quality guidelines were relatively consistent. Meanwhile, recommendations for remdesivir, baricitinib and casirivimab/imdevimab varied between guidelines. Recommendations relating to remdesivir were particularly varied in moderate and severe disease categories, while baricitinib recommendations were most varied in mild COVID-19. Across all disease severity categories, there was no agreement on recommendations for casirivimab/imdevimab.

There was a general trend between the number of guidelines discussing a particular therapeutic and the availability and extent of clinical trial data. Remdesivir[5, 42-45, 58], dexamethasone[3, 46, 47, 59, 60], and tocilizumab[52-56, 61-67] have been investigated in several clinical trials and were discussed by almost all guidelines, while baricitinib[6, 57] and casirivimab/imdevimab[7, 62] had been investigated in fewer trials and were discussed within fewer guidelines. For all therapies except remdesivir, there also appeared to be a trend between the availability of clinical trial data and the consensus between recommendations; a greater number of clinical trials was typically associated with increased consensus. It may be argued that when clinical data are not readily available, guideline development groups either make no recommendation or fill data vacuums with input from expert opinion, which can be highly subjective and inconsistent. When data become available, such recommendations may require changing or updating.

Even when clinical trial data are available, guideline development groups may not choose to use all data to inform decisions. For example, multiple trials[3, 46, 47, 59, 60] examined dexamethasone in COVID-19 patients, but most guidelines considered just a few studies (or often considered trials of other corticosteroids) and have not

updated their dexamethasone recommendations since mid-2020. This is likely due to the publication of strong supportive evidence for the use of dexamethasone early in the pandemic,[3] and later evidence corroborating earlier findings.[47] However, if trial results are less strong or consistent, considering different data (due to the timing of evidence publication, for example) could lead to differences in recommendations. For instance, many of the remdesivir trials came to contradictory conclusions: ACTT-1[5] and SIMPLE Moderate[58] had broadly positive results; the WHO Solidarity trial[42] suggested negative outcomes; and Wang 2020[43] was insufficiently powered, stopping early with negative outcomes.

There were no clear trends between recommendations and the consideration of specific outcomes. However, where recommendations were more consistent, high-visibility endpoints were considered. Guidelines relating to dexamethasone more frequently considered mortality, for example. Notably, for dexamethasone recommendations, few outcomes other than mortality were considered, likely because death is the most visible and important clinical endpoint; when mortality data are available, other outcomes are less influential. As the number of approved therapies increased over the course of the pandemic, and more outcomes were included and achieved in trials, it is likely that these were increasingly favoured over less visible outcomes.

The timing of a study and its data cuts may also influence the specific outcomes to consider. Some ordinal scales, for instance, were developed over the course of trials, and mortality outcomes were sometimes only available as post-hoc analyses. Consequently, the timing of guideline publication and/or guideline updates may have influenced which data cuts and outcomes were considered in guideline development. Further, the timing of the study may influence how outcomes were defined. For example, patients enrolled in the ACTT-1 trial[5] (which was conducted early in the pandemic) were sometimes required to remain hospitalised, despite not requiring supplemental oxygen or ongoing medical care, due to infection-control measures. As such, discharge-related outcomes required re-defining.[5]

The inconsistency between recommendations, even when considering the same evidence, suggests that subjective factors, such as cost, accessibility, alignment with other guidelines and COVID-19 variants, also had an impact. Therefore, while the current review provides novel information to clinicians, they should be aware of and

consider the totality of evidence prior to making treatment decisions. Additionally, clinicians should recognize when there is a possibility that non-clinical factors may play a role in influencing clinical practice decisions. For instance, dexamethasone is a relatively low-cost therapy, which may reinforce the preference to recommend its use. Meanwhile, remdesivir is more expensive, which, when combined with inconsistent data, could support decisions to recommend against its widespread use. Access may have contributed to the AWMF's[24] recommendation for casirivimab/imdevimab, given the purchase of a large stock of this therapy by Germany's Health Ministry.[68] In resource-limited settings, alignment with the WHO guidelines[9] may have an influential role; in their recommendations against remdesivir, the Brazilian MoH and Indian CIDS referenced the WHO guidelines, which also did not recommend this therapy. Finally, guideline groups may have shaped casirivimab/imdevimab recommendations on the most prevalent COVID-19 variant in circulation. This is an important consideration which was not investigated here but will likely play an increasing role in COVID-19 treatment guidelines, particularly for monoclonal antibodies. Finally, it would be important to look back at the actions and processes taken during the COVID-19 pandemic to see what can be learned for future situations.

Limitations

There are limitations to this analysis that should be considered when interpreting its findings. Foremost, the review focused only on five major therapies, although others (such as sarilumab and tofacitinib) have been recommended for use among hospitalised COVID-19 patients.[8, 9, 36] The review does not consider recommendations relevant to non-hospitalised patients, who arguably make up the greatest proportion of individuals infected with COVID-19. [69] Further, while a broad geographical range was represented by the included guidelines, large geographical regions (for instance, the Middle East and Africa) were left unrepresented.

Though the review looked to identify the most up-to-date guidelines, the rapid development and regular updating of recommendations inevitably meant that more recent information may not have been captured in this review. For instance, treatment guidelines such as those developed by the European Society of Clinical Microbiology and Infectious Diseases[70] were not captured, as they were published

after the initial guideline identification cut-off. Similarly, impactful data from recent clinical trials may have been missed. One example is the PINETREE trial (NCT04501952), which is referenced by multiple guidelines to support remdesivir recommendations in patients at high risk for progression to severe disease in hospital and ambulatory settings.[71] Other recent results from trials evaluating mAbs, oral antivirals and JAK inhibitors have also been considered, but these data were not incorporated into guidelines at the time of review.

The AGREE-II tool is just one method for assessing the quality of guidelines, and has limitations in its approach. While the use of the AGREE-II Domain 3 to determine high-quality is supported by other published guideline assessments, this definition disregards other characteristics which contribute to guideline quality, such as stakeholder engagement or clarity of presentation. Given the variability in performance across Domains, an alternative definition of high-quality would have led to a different set of guidelines undergoing full data extraction. Additionally, since AGREE-II does not address in detail the methodology used for guideline development (e.g. analyses of bias and variance), analysis of the methodology could also result in a different set of included guidelines. AGREE-II also does not consider the quality of the evidence included in the guidelines, and thus the current review did not aim to make a judgement of the quality of the evidence included in any given guideline, but simply to assess the quality of the guideline in a systematic manner. Another potential limitation of AGREE-II is the lack of analysis of population/intervention/comparison/outcome (PICO) questions and formulations across guidelines; differences recommendations could arise due to differences in PICO questions. Further, though AGREE-II has been used to assess many types of guidelines, its applicability to those developed in health emergencies such as pandemics has not yet been verified. Finally, the findings of this review are themselves limited by the transparency of guidelines regarding their justifications for recommendations, and the rapid pace at which guidelines, clinical trial results, and recommendations are being updated in line with evolving data and COVID-19 variants.

ACKNOWLEDGEMENTS

The authors acknowledge Craig Brooks-Rooney, Isabel Katz, Aashna Shah, and Yasemin Hasimoglu, Costello Medical Inc., USA, and Hannah Luedke, Sara Steeves,

Vittoria Vardanega, Dario Gregori, and Jacques Morgan, Costello Medical Consulting Ltd., UK, for their assistance in conducting this review. Support for this assistance was funded by Gilead Sciences Inc. This work was sponsored by Gilead Sciences Inc.

CONCLUSION

This review identified substantial heterogeneity in the quality of guidelines for the therapeutic treatment of patients hospitalised with COVID-19. Further, even among high-quality guidelines, it was found that recommendations regarding specific therapeutics varied, despite using the same clinical trials and specific outcomes. These findings suggest that unreported, subjective factors may also play a role, particularly where evidence is limited or conflicting, and call for guideline groups to justify their recommendations more transparently. In response to health emergencies, greater global collaboration to produce, synthesise and update evidence, along with country- or region-specific efforts to develop locally-relevant recommendations, may provide clearer and more consistent guidance to clinicians.

AUTHORS' CONTRIBUTIONS

Substantial contributions to study conception and design: **DAW; AAE; LD; CYW; AL; RAB; AR; JKR**; substantial contributions to analysis and interpretation of the data: **DAW; AAE; LD; CYW; AL; RAB; AR; KCS; JKR**; drafting the article or revising it critically for important intellectual content: **DAW; AAE; LD; CYW; AL; RAB; AR; KCS; JKR**; final approval of the version of the article to be published: **DAW; AAE; LD; CYW; AL; RAB; AR; KCS; JKR**.

DISCLOSURES

DAW: Research funding from Gilead Sciences, Merck & Co, and Eli Lilly to university; honoraria for consulting, advisory board, and educational events from Gilead Sciences.

AAE; LD; CYW; AL; RAB: Employees of Gilead Sciences Inc., Foster City, CA, USA.

AR: Employee of Costello Medical Inc., Boston, MA, USA.

KCS: Employee of Costello Medical Consulting Ltd., Cambridge, UK.

JKR: Consulting fees from Gilead Sciences, Janssen, Merck, and ViiV Healthcare Limited; payment for lectures, presentations, speakers bureaus, manuscript writing or educational events from Gilead Sciences, Janssen, Merck, Theratechnologies and

ViiV Healthcare Limited; participation on a Data Safety Monitoring Board or Advisory Board for Abivax, Galapagos, Gilead Sciences, Merck, and ViiV Healthcare Limited.

REFERENCES

1. *BMJ Best Practice. Coronavirus disease 2019 (COVID-19). 2022.*
<https://bestpractice.bmj.com/topics/en-gb/3000201> (accessed March 29 2022).
2. Thorlund, K., et al., *A real-time dashboard of clinical trials for COVID-19.* Lancet Digit. Health, 2020. **2**(6): p. e286-e287.
3. *RECOVERY Collaborative Group. Dexamethasone in hospitalized patients with Covid-19.* N. Engl. J. Med, 2021. **384**(8): p. 693-704.
4. *RECOVERY Collaborative Group. Tocilizumab in patients admitted to hospital with COVID-19 (RECOVERY): a randomised, controlled, open-label, platform trial.* Lancet, 2021. **397**(10285): p. 1637-1645.
5. Beigel, J.H., et al., *Remdesivir for the treatment of Covid-19.* N. Engl. J. Med, 2020. **383**(19): p. 1813-1826.
6. Kalil, A.C., et al., *Baricitinib plus remdesivir for hospitalized adults with Covid-19.* N. Engl. J. Med, 2021. **384**(9): p. 795-807.
7. Weinreich, D.M., et al., *REGN-COV2, a neutralizing antibody cocktail, in outpatients with Covid-19.* N. Engl. J. Med, 2021. **384**(3): p. 238-251.
8. *National Institute for Health and Care Excellence. COVID-19 rapid guideline: Managing COVID-19 (version 19.1). Published 10.01.22.*
9. Agarwal, A., et al., *A living WHO guideline on drugs for covid-19.* BMJ, 2020. **370**.
10. Chalmers, J.D., et al., *Management of hospitalised adults with coronavirus disease 2019 (COVID-19): a European Respiratory Society living guideline.* Eur Respir J, 2021. **57**(4).
11. Singh, J.A. and R. Ravinetto, *COVID-19 therapeutics: how to sow confusion and break public trust during international public health emergencies.* J Pharm Policy Pract, 2020. **13**(1): p. 47.
12. Pulla, P., *Covid-19: India's slow moving treatment guidelines are misleading and harming patients.* BMJ, 2021. **372**: p. n278.
13. *The World Bank. GDP (current US\$).*
<https://data.worldbank.org/indicator/NY.GDP.MKTP.CD> (accessed March 29, 2022).
14. *Our World in Data. Coronavirus Pandemic (COVID-19) – the data.*
<https://ourworldindata.org/coronavirus-data> (accessed March 29, 2022).
15. Brouwers, M.C., et al., *AGREE II: advancing guideline development,*

- reporting and evaluation in health care. Cmaj, 2010. 182(18): p. E839-42.*
16. Doniselli, F.M., et al., *A critical appraisal of the quality of low back pain practice guidelines using the AGREE II tool and comparison with previous evaluations: a EuroAIM initiative. Eur. Spine J, 2018. 27(11): p. 2781-2790.*
 17. Polus, S., et al., *Appraisal of WHO Guidelines in Maternal Health Using the AGREE II Assessment Tool. PloS one, 2012. 7(8): p. e38891.*
 18. Sabharwal, S., et al., *High Methodologic Quality But Poor Applicability: Assessment of the AAOS Guidelines Using The AGREE II Instrument. Clin. Orthop. Relat. Res., 2014. 472(6): p. 1982-1988.*
 19. Zhao, S., et al., *A quality evaluation of guidelines on five different viruses causing public health emergencies of international concern. Ann Transl Med, 2020. 8(7): p. 500.*
 20. Zhu, Q., et al., *How about the quality and recommendation on prevention, diagnosis, and treatment of HIV/AIDS guidelines developed by WHO: A protocol for systematic review. Medicine (Baltimore), 2020. 99(52): p. e23638.*
 21. Kim, S.G., et al., *Guidelines for the diagnosis and treatment of H elicobacter pylori infection in K orea, 2013 revised edition. J. Gastroenterol. Hepatol, 2014. 29(7): p. 1371-1386.*
 22. Arevalo-Rodriguez, I., et al., *Alzheimer's disease dementia guidelines for diagnostic testing: a systematic review. AJADD, 2013. 28(2): p. 111-119.*
 23. Smith, C.A., et al., *A systematic critical appraisal of clinical practice guidelines in juvenile idiopathic arthritis using the appraisal of guidelines for research and evaluation II (AGREE II) instrument. PloS one, 2015. 10(9): p. e0137180.*
 24. Kluge, S., et al., *S2k-Leitlinie–Empfehlungen zur stationären Therapie von Patienten mit COVID-19. Pneumologie, 2021. 75(02): p. 88-112.*
 25. *Covid Management Guidelines India Group - Anti-inflammatory Working Group. Tocilizumab. May 24, 2021. <https://indiacovidguidelines.org/tocilizumab/> (accessed March 29, 2022).*
 26. *Covid Management Guidelines India Group - Anti-viral Working Group. Remdesivir. June 1, 2021. <https://indiacovidguidelines.org/remdesivir/> (accessed March 29, 2022).*
 27. *Covid Management Guidelines India Group - Anti-inflammatory Working Group. Systemic Corticosteroids. July 15, 2021. <https://indiacovidguidelines.org/systemic-corticosteroids/> (accessed March 29, 2022).*
 28. *Covid Management Guidelines India Group – Anti-inflammatory and*

- Antibody working Group – Baricitinib. September 17th, 2021.*
<https://indiacovidguidelines.org/baricitinib/> (accessed March 29, 2022).
29. *Covid Management Guidelines India Group – Anti-body Working Group. Casirivimab – Imdevimab. August 16, 2021.*
<https://indiacovidguidelines.org/casirivimab-imdevimab-moderate-to-severe/?preview=true> (accessed March 29, 2022).
 30. Morris, A., N. Stall, and P. Bobos, *Tocilizumab for hospitalized patients with COVID-19. Science Briefs of the Ontario COVID-19 Science Advisory Table. 2021; 2 (11).* 2021, Updated on March 2, 2021. Version 1.0 is available under Additional
 31. Bailey, J., A. Morris, and S. Bean, *Evidence-based recommendations on the use of anti-SARS-CoV-2 monoclonal antibodies (casirivimab+ imdevimab, and sotrovimab) for adults in Ontario.* Science Briefs of the Ontario COVID-19 Science Advisory Table, 2021. **2**: p. 51.
 32. Hempel, A., S. Jevtic, and S. Vandersluis, *Baricitinib for hospitalized patients with COVID-19.* Science Briefs of the Ontario COVID-19 Science Advisory Table, 2022. **3**: p. 53.
 33. Jüni, P., A. Odutayo, and U. Allen, *Dexamethasone in patients hospitalized for COVID-19.* Science Briefs of the Ontario COVID-19 Science Advisory Table, 2020. **1**(1).
 34. Morris, A., P. Jüni, and A. Odutayo, *Remdesivir for hospitalized patients with COVID-19.* Science Briefs of the Ontario COVID-19 Science Advisory Table, 2021. **2**: p. 27.
 35. *Gobierno de México. Guía clínica para el tratamiento de la COVID-19 en México. August 2, 2021.* https://coronavirus.gob.mx/wp-content/uploads/2021/08/GuiaTx_COVID19_ConsensoInterinstitucional_2021.08.03.pdf (accessed March 29, 2022).
 36. *Infectious Diseases Society of America. IDSA guidelines on the treatment and management of patients with COVID-19 (v.5.6.0).* 2021.
 37. Yamakawa, K., et al., *Japanese rapid/living recommendations on drug management for COVID-19: updated guidelines (September 2021).* Acute Med. Surg., 2021. **8**(1): p. e706.
 38. *Ministério da Saúde. Diretrizes Brasileiras para Tratamento Hospitalar do Paciente com COVID-19 – Capítulo 2: Tratamento Farmacológico. Maio 2021.* 2021.
 39. *National Institute of Health. Coronavirus disease 2019 (COVID-19) treatment guidelines. January 5, 2022.*
<https://www.covid19treatmentguidelines.nih.gov/> (accessed January 11, 2022). 2021.

40. Bassetti, M., et al., *Clinical management of adult patients with COVID-19 outside intensive care units: guidelines from the Italian Society of Anti-Infective Therapy (SITA) and the Italian Society of Pulmonology (SIP)*. *Infect Dis Ther*, 2021. **10**(4): p. 1837-1885.
41. Alhazzani, W., et al., *Surviving sepsis campaign guidelines on the management of adults with coronavirus disease 2019 (COVID-19) in the ICU: first update*. *Crit. Care Med.*, 2021. **49**(3): p. e219-e234.
42. Pan, H., et al., *Consortium WST (2021) Repurposed Antiviral Drugs for Covid-19-Interim WHO Solidarity Trial Results*. *N. Engl. J. Med.* **384**(6): p. 497-511.
43. Wang, Y., et al., *Remdesivir in adults with severe COVID-19: a randomised, double-blind, placebo-controlled, multicentre trial*. *The Lancet*, 2020. **395**(10236): p. 1569-1578.
44. Goldman, J.D., et al., *Remdesivir for 5 or 10 days in patients with severe Covid-19*. *N. Engl. J. Med.*, 2020. **383**(19): p. 1827-1837.
45. Mahajan, L. and G. AP Singh, *Clinical outcomes of using remdesivir in patients with moderate to severe COVID-19: A prospective randomised study*. *Indian J. Anaesth*, 2021. **65**(Suppl 1): p. S41.
46. *ClinicalTrials.gov. Efficacy of Dexamethasone Treatment for Patients With ARDS Caused by COVID-19 (DEXA-COVID19)(NCT04325061)*. 2021. <https://clinicaltrials.gov/ct2/show/NCT04325061> (accessed March 29, 2022).
47. Tomazini, B.M., et al., *Effect of dexamethasone on days alive and ventilator-free in patients with moderate or severe acute respiratory distress syndrome and COVID-19: the CoDEX randomized clinical trial*. *JAMA*, 2020. **324**(13): p. 1307-1316.
48. Angus, D.C., et al., *Effect of hydrocortisone on mortality and organ support in patients with severe COVID-19: the REMAP-CAP COVID-19 corticosteroid domain randomized clinical trial*. *Jama*, 2020. **324**(13): p. 1317-1329.
49. Dequin, P.-F., et al., *Effect of hydrocortisone on 21-day mortality or respiratory support among critically ill patients with COVID-19: a randomized clinical trial*. *JAMA*, 2020. **324**(13): p. 1298-1306.
50. Munch, M.W., et al., *Low-dose hydrocortisone in patients with COVID-19 and severe hypoxia: The COVID STEROID randomised, placebo-controlled trial*. *Acta Anaesthesiol. Scand.*, 2021. **65**(10): p. 1421-1430.
51. Corral-Gudino, L., et al., *Methylprednisolone in adults hospitalized with COVID-19 pneumonia*. *Wiener klinische Wochenschrift*, 2021. **133**(7): p. 303-311.
52. Veiga, V.C., et al., *Effect of tocilizumab on clinical outcomes at 15 days in*

- patients with severe or critical coronavirus disease 2019: randomised controlled trial.* BMJ, 2021. **372**.
53. Hermine, O., et al., *Effect of tocilizumab vs usual care in adults hospitalized with COVID-19 and moderate or severe pneumonia: a randomized clinical trial.* JAMA internal medicine, 2021. **181**(1): p. 32-40.
 54. Rosas, I.O., et al., *Tocilizumab in hospitalized patients with severe Covid-19 pneumonia.* N. Engl. J. Med, 2021. **384**(16): p. 1503-1516.
 55. Salama, C., et al., *Tocilizumab in patients hospitalized with Covid-19 pneumonia.* N. Engl. J. Med, 2021. **384**(1): p. 20-30.
 56. Stone, J.H., et al., *Efficacy of tocilizumab in patients hospitalized with Covid-19.* N. Engl. J. Med, 2020. **383**(24): p. 2333-2344.
 57. Marconi, V.C., et al., *Efficacy and safety of baricitinib for the treatment of hospitalised adults with COVID-19 (COV-BARRIER): a randomised, double-blind, parallel-group, placebo-controlled phase 3 trial.* The Lancet Respiratory Medicine, 2021. **9**(12): p. 1407-1418.
 58. Spinner, C.D., et al., *Effect of remdesivir vs standard care on clinical status at 11 days in patients with moderate COVID-19: a randomized clinical trial.* JAMA, 2020. **324**(11): p. 1048-1057.
 59. Jamaati, H., et al., *No clinical benefit of high dose corticosteroid administration in patients with COVID-19: a preliminary report of a randomized clinical trial.* Eur. J. Pharmacol, 2021. **897**: p. 173947.
 60. Ranjbar, K., et al., *Methylprednisolone or dexamethasone, which one is superior corticosteroid in the treatment of hospitalized COVID-19 patients: a triple-blinded randomized controlled trial.* BMC infectious diseases, 2021. **21**(1): p. 1-8.
 61. *EU Clinical Trials Register. PreToVid (2020-001375-32).*
<https://www.clinicaltrialsregister.eu/ctr-search/trial/2020-001375-32/NL>
(accessed March 30, 2022).
 62. *RECOVERY Collaborative Group. Casirivimab and imdevimab in patients admitted to hospital with COVID-19 (RECOVERY): a randomised, controlled, open-label, platform trial.* Lancet, 2022. **399**(10325): p. 665-676.
 63. Gordon, A.C., et al., *Interleukin-6 receptor antagonists in critically ill patients with Covid-19.* N. Engl. J. Med, 2021.
 64. Salvarani, C., et al., *Effect of tocilizumab vs standard care on clinical worsening in patients hospitalized with COVID-19 pneumonia: a randomized clinical trial.* JAMA internal medicine, 2021. **181**(1): p. 24-31.
 65. Soin, A.S., et al., *Tocilizumab plus standard care versus standard care in patients in India with moderate to severe COVID-19-associated cytokine*

- release syndrome (COVINTOC): an open-label, multicentre, randomised, controlled, phase 3 trial.* Lancet Respir. Med., 2021. **9**(5): p. 511-521.
66. Talaschian, M., et al., *Tocilizumab Failed to Reduce Mortality in Severe COVID-19 Patients: Results From a Randomized Controlled Clinical Trial.* Research Square, 2021.
 67. Wang, D., et al., *Tocilizumab in patients with moderate or severe COVID-19: a randomized, controlled, open-label, multicenter trial.* Front. Med., 2021. **15**(3): p. 486-494.
 68. *European Biotechnology. Germany orders antibody therapies from USA. 2021. <https://european-biotechnology.com/up-to-date/latest-news/news/germany-orders-antibody-therapies-from-usa.html> (accessed March 29, 2022).*
 69. Menachemi, N., et al., *How Many SARS-CoV-2-Infected People Require Hospitalization? Using Random Sample Testing to Better Inform Preparedness Efforts.* J Public Health Manag Pract, 2021. **27**(3): p. 246-250.
 70. Bartoletti, M., et al., *ESCMID COVID-19 Living guidelines: drug treatment and clinical management.* Clin Microbiol Infect, 2021.
 71. Gottlieb, R.L., et al., *Early remdesivir to prevent progression to severe Covid-19 in outpatients.* N. Engl. J. Med, 2022. **386**(4): p. 305-315.
 72. Siemieniuk, R., et al., *A living WHO guideline on drugs for covid-19.* BMJ, 2020. **370**: p. m3379.
 73. Afra, K., L.Y. Chen, and D. Sweet, *Tocilizumab for hospitalized patients with COVID-19.* CMAJ, 2021. **193**(15): p. E521-E521.

Table 1: Summary of Evidence and Outcomes Reported by Guidelines Discussing Remdesivir

Guidelines	RCTs						Outcomes Considered							
	ACTT-1 (NCT04280705)[5]	SIMPLE Severe (NCT04292899)[44]	SIMPLE Moderate (NCT04292730)[58]	WHO Solidarity (NCT04315948/ISRCTN83971151)[42]	Wang 2020 (NCT04257656)[43]	Mahajan 2021 (NCT NR)[45]	Cost	Mortality	Clinical recovery	Discharge	Safety	Time to ICU admission	Progression to IMV	Other
Guidelines Recommending for the Use of Remdesivir in At Least One Population Described														
SSC	✓		✓	✓	✓		✓	✓	✓		✓			
COVID-19 Advisory Ontario	✓	✓	✓	✓	✓			✓	✓	✓		✓	✓	✓ ^a
SITA/SIP	✓	✓	✓	✓	✓			✓	✓					
J-SSCG	✓		✓	✓	✓		✓	✓	✓		✓		✓	
Government of Mexico											✓			
NICE	✓	✓	✓	✓	✓			✓	✓	✓	✓		✓	✓ ^b
IDSA	✓			✓	✓			✓	✓		✓			
NIH	✓		✓	✓	✓			✓	✓		✓		✓	✓ ^c
Guidelines Recommending Against the Use of Remdesivir in Any Population Described*														

Guidelines	RCTs						Outcomes Considered							
	ACTT-1 (NCT04280705)[5]	SIMPLE Severe (NCT04292899)[44]	SIMPLE Moderate (NCT04292730)[58]	WHO Solidarity (NCT04315948/ISRCTN83971151)[42]	Wang 2020 (NCT04257656)[43]	Mahajan 2021 (NCT NR)[45]	Cost	Mortality	Clinical recovery	Discharge	Safety	Time to ICU admission	Progression to IMV	Other
ERS	✓		✓	✓	✓		✓	✓			✓			✓ ^d
WHO	✓		✓	✓	✓			✓		✓	✓			✓ ^e
MoH Brazil	✓		✓	✓	✓		✓	✓	✓	✓	✓	✓	✓	✓ ^f
AWMF	✓		✓	✓	✓	✓	✓	✓			✓		✓	✓ ^g
CIDS	✓		✓	✓	✓	✓	✓	✓	✓	✓	✓		✓	✓ ^h

✓: guideline explicitly mentions evidence or outcome. Empty cell: guideline does not explicitly mention evidence or outcome. *Includes guidelines which only provided recommendations against remdesivir or recommended against remdesivir for at least one disease severity category and reported insufficient evidence for any other categories.

Organizations: AWMF: Association of the Scientific Medical Societies in Germany; CIDS: Clinical Infectious Disease Society; ERS: European Respiratory Society; IDSA: Infectious Disease Society of America; J-SSCG: Japanese Clinical Practice Guidelines for Management of Sepsis and Septic Shock; MoH: Ministry of Health; NICE: National Institute for Health and Care Excellence; NIH: National Institutes of Health; SIP: Italian Society of Pulmonology; SITA: Italian Society of Anti-Infective Therapy; SSC: Surviving Sepsis Campaign; WHO: World Health Organization.

Abbreviations: ICU: intensive care unit; IMV: invasive mechanical ventilation; NCT: National Clinical Trial; NR: not reported; RCT: randomised controlled trial.

Other outcomes considered: ^aICU length of stay was considered; however, published data precluded the pooling of trials for ICU length. Need for oxygen support was considered. Clinical improvement outcomes were considered. ^bSeptic shock within 28 or 30 days; acute respiratory failure or acute respiratory distress syndrome (ARDS) within 28 and 30 days. ^cClinical improvement at Day 15, clinical status distribution on Day 11 + Day 14. ^dAdditional endpoints which were searched for by the guideline development committee but were either not studied or data was not found in an extractable format were: deterioration in those not requiring ventilation at start of treatment; requirement for oxygen; hospital admission; ICU length of stay; need for non-invasive ventilation; hospital length of stay; severity of symptoms; improvement in oxygen saturations or arterial blood gases; relapse and Duration of fever. ^eViral clearance (7 days); acute kidney injury; delirium; time to clinical improvement; duration of ventilation. ^fLack of

benefit in patients using IMV. ^gClinical worsening: new need for non-invasive mechanical ventilation or high-flow oxygen at day 28; clinical worsening: new need for oxygen by mask or nasal prongs at day 28; quality of life. ^hProgression to oxygen therapy; progression to NIV or high-flow oxygen.

Sources: AWMF, October 2021;[24] CIDS, June 2021;[26] COVID-19 Advisory Ontario, May 2021;[34] ERS, April 2021;[10] Government of Mexico, August 2021;[35] IDSA, November 2021;[36] J-SSCG, September 2021;[37] MoH Brazil, May 2021;[38] NICE, January 2022;[8] NIH, December 2021;[39] SITA/SIP, May 2021;[40] SSC, March 2021;[41] WHO, January 2022.[72]

Table 2: Summary of Evidence and Outcomes Reported by Guidelines Discussing Dexamethasone

Guidelines	RCTs					Outcomes Considered							
	DEXA-COVID19 (NCT04325061)[46]	RECOVERY (NCT04381936)[3]	CoDEX (NCT04327401)[47]	Jamaati 2021 (IRCT20151227025726N17)[59]	Ranjbar 2021 (IRCT20200204046369N1)[60]	Cost	Mortality	Clinical recovery	Discharge	Safety	Time to ICU admission	Progression to IMV	Other
Guidelines Recommending for the Use of Dexamethasone in At Least One Population Described													
ERS	✓	✓	✓			✓	✓		✓	✓	✓		✓ ^a
SSC	✓	✓	✓			✓	✓			✓			✓ ^b
WHO	✓	✓	✓				✓		✓	✓		✓	
MoH Brazil						✓	✓		✓	✓			
COVID-19 Advisory Ontario		✓	✓				✓					✓	
AWMF		✓			✓	✓	✓			✓		✓	✓ ^c
CIDS		✓				✓	✓			✓		✓	

Guidelines	RCTs					Outcomes Considered							
	DEXA-COVID19 (NCT04325061)[46]	RECOVERY (NCT04381936)[3]	CoDEX (NCT04327401)[47]	Jamaati 2021 (IRCT20151227025726N17)[59]	Ranjbar 2021 (IRCT20200204046369N1)[60]	Cost	Mortality	Clinical recovery	Discharge	Safety	Time to ICU admission	Progression to IMV	Other
SITA/SIP	✓	✓	✓				✓						
J-SSCG	✓	✓	✓			✓	✓	✓					
Government of Mexico	✓	✓	✓	✓	✓		✓			✓			
NICE	✓	✓	✓	✓	✓		✓		✓	✓		✓	✓ ^d
IDSA	✓	✓	✓				✓		✓	✓			
NIH		✓	✓				✓		✓	✓	✓	✓	✓ ^e

✓: guideline explicitly mentions evidence or outcome. Empty cell: guideline does not explicitly mention evidence or outcome.

Organizations: AWMF: Association of the Scientific Medical Societies in Germany; CIDS: Clinical Infectious Disease Society; ERS: European Respiratory Society; IDSA: Infectious Disease Society of America; J-SSCG: Japanese Clinical Practice Guidelines for Management of Sepsis and Septic Shock; MoH: Ministry of Health; NICE: National Institute for Health and Care Excellence; NIH: National Institutes of Health; SIP: Italian Society of Pulmonology; SITA: Italian Society of Anti-Infective Therapy; SSC: Surviving Sepsis Campaign; WHO: World Health Organization.

Abbreviations: ARDS: acute respiratory distress syndrome; ICU: intensive care unit; IMV: invasive mechanical ventilation; NCT: National Clinical Trial; NR: not reported; RCT: randomised controlled trial.

Other outcomes considered: ^aAdditional endpoints not included in the evidence table which were searched for but were either not studied or data was not found in an extractable format were: clinical resolution or cure (also includes the reverse i.e patients not cured); time to clinical improvement or resolution on an ordinal scale; requirement for oxygen; hospital admission; ordinal scale or clinical status at day 28; ICU length of stay; need for non-invasive ventilation; deterioration in those not requiring ventilation at start of treatment; severity of symptoms; improvement in oxygen saturations or arterial blood gases; relapse; viral clearance (negative SARS-CoV-2 test) and duration of fever. ^bExamined evidence from non-COVID-19 ARDS patients as well. ^cVentilator-free days 28 days; quality of life, including fatigue and neurological status long term; hospital-acquired infections during treatment per 28 days. ^dOutcomes assessed by end of treatment: gastrointestinal bleeding, bacterial co-infections, hyperglycemia, neuromuscular weakness, and neuropsychiatric effects. ^eNeed for insulin; positive blood cultures at Day 7; sepsis by Day 28; score on 6-point WHO ordinal scale at Day 15; mean Sequential Organ Failure Assessment (SOFA) score; need for rescue strategies; need or oxygenation; organ-support free-days at Day 21; duration of fever; virus clearance time; proportion prescribed antibiotics and antifungal therapy.

Sources: AWMF, October 2021;[24] CIDS, July 2021;[27] COVID-19 Advisory Ontario, August 2021;[33] ERS, April 2021;[10] Government of Mexico, August 2021;[35] IDSA, October 2021;[36] J-SSCG, September 2021;[37] MoH Brazil, May 2021;[38] NICE, January 2022;[8] NIH, December 2021;[39] SITA/SIP, May 2021;[40] SSC, March 2021;[41] WHO, January 2022.[72]

Table 3: Summary of Evidence and Outcomes Reported by Guidelines Discussing Tocilizumab

Guidelines	RCTs												Outcomes Considered							
	REMAP-CAP (NCT02735707)[63]	RECOVERY (NCT04381936)[4]	TOCIBRAS (NCT04403685)[52]	CORIMUNO-19 (NCT04331808)[53]	COVACTA(NCT04320615)[54]	EMPACTA (NCT04372186)[55]	Salvarani 2021 (NCT04346355)[64]	Stone 2020 (NCT04356937)[56]	Wang 2020 (ChiCTR2000029765)[67]	COVINTOC (CTRI/2020/05/025369)[65]	PreToVid (EU-CTR-2020-001375-32)[61]	Talaschian 2021[66]	Cost	Mortality	Clinical recovery	Discharge	Safety	Time to ICU admission	Progression to IMV	Other
Guidelines Recommending for the Use of Tocilizumab in At Least One Population Described																				
ERS	✓	✓	✓	✓	✓	✓	✓	✓						✓			✓		✓	✓ ^a
WHO														✓	✓	✓	✓		✓	✓ ^b
COVID-19 Advisory Ontario	✓	✓	✓	✓	✓	✓	✓	✓	✓					✓		✓	✓		✓	✓ ^c
AWMF	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓				✓	✓	✓	✓		✓	✓ ^d
CIDS	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓			✓	✓	✓	✓	✓	✓	✓	✓ ^e
SITA/SIP		✓	✓	✓	✓	✓	✓	✓		✓				✓	✓		✓	✓ [*]	✓ ^{**}	✓ ^f
J-SSCG	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓		✓			✓			
Government of Mexico														✓			✓		✓	

Guidelines	RCTs											Outcomes Considered								
	REMAP-CAP (NCT02735707)[63]	RECOVERY (NCT04381936)[4]	TOCIBRAS (NCT04403685)[52]	CORIMUNO-19 (NCT04331808)[53]	COVACTA(NCT04320615)[54]	EMPACKTA (NCT04372186)[55]	Salvarani 2021 (NCT04346355)[64]	Stone 2020 (NCT04356937)[56]	Wang 2020 (ChiCTR2000029765)[67]	COVINTOC (CTRI/2020/05/025369)[65]	PreToVid (EU-CTR-2020-001375-32)[61]	Talaszian 2021[66]	Cost	Mortality	Clinical recovery	Discharge	Safety	Time to ICU admission	Progression to IMV	Other
NICE	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓				✓	✓		✓	✓	✓	✓ ^g
IDSA	✓	✓	✓	✓	✓	✓	✓	✓						✓			✓	✓	✓	✓ ^h
NIH	✓	✓	✓	✓	✓	✓	✓	✓		✓				✓			✓		✓	✓ ⁱ
Guidelines Recommending Against the Use of Tocilizumab in Any Population Described[†]																				
MoH Brazil	✓	✓	✓	✓	✓	✓	✓	✓						✓			✓	✓	✓	

✓: Guideline explicitly mentions evidence or outcome. Empty cell: Guideline does not explicitly mention evidence or outcome. *Composite endpoint of ICU admission, death, or clinical worsening led to enrolment discontinuation for futility. **Composite of death, mechanical ventilation, and clinical worsening. †Includes guidelines which only provided recommendations against tocilizumab or recommended against tocilizumab for at least one disease severity category and reported insufficient evidence for any other categories.

Organizations: AWMF: Association of the Scientific Medical Societies in Germany; CIDS: Clinical Infectious Disease Society; ERS: European Respiratory Society; IDSA: Infectious Disease Society of America; J-SSCG: Japanese Clinical Practice Guidelines for Management of Sepsis and Septic Shock; MoH: Ministry of Health; NICE: National Institute for Health and Care Excellence; NIH: National Institutes of Health; SIP: Italian Society of Pulmonology; SITA: Italian Society of Anti-Infective Therapy; SSC: Surviving Sepsis Campaign; WHO: World Health Organization.

Abbreviations: ICU: intensive care unit; IMV: invasive mechanical ventilation; NCT: National Clinical Trial; NR: not reported; RCT: randomised controlled trial.

Other outcomes considered: ^aClinical worsening. ^bDuration of IMV. ^cComposite or mortality or invasive mechanical ventilation was considered; respiratory or cardiovascular organ support-free days was considered. ^dNeed for new hemodialysis/hemofiltration. ^eVentilator-free days. ^fDisease progression; Scoring higher than 5 on day 4 on a 10 points ordinal clinical scale. ^gOrdinal scale combining in-hospital mortality and days free of organ support by Day 21, days free of organ support in survivors by Day 21. ^hClinical deterioration (follow up: range 14 days to 30 days). ⁱProgression to high-flow oxygen or NIV.

Sources: AWMF, October 2021;[24] CIDS, May 2021;[25] COVID-19 Advisory Ontario, March 2021;[73] ERS, April 2021;[10] Government of Mexico, August 2021;[35] IDSA, October 2021;[36] J-SSCG, September 2021;[37] MoH Brazil, May 2021;[38] NICE, January 2022;[8] NIH, December 2021;[39] SITA/SIP, May 2021;[40] WHO, January 2022.[72]

Table 4: Summary of Evidence and Outcomes Reported by Guidelines Discussing Baricitinib

Guidelines	RCTs		Outcomes Considered							
	ACTT-2 (NCT04401579)[6]	COV-BARRIER (NCT04421027)[57]	Cost	Mortality	Clinical recovery	Discharge	Safety	Time to ICU admission	Progression to IMV	Other
Guidelines Recommending for the Use of Baricitinib in At Least One Population Described										
WHO	✓	✓		✓	✓	✓	✓		✓	✓ ^a
COVID-19 Advisory Ontario	✓	✓		✓	✓		✓		✓	✓ ^b
AWMF	✓	✓	✓	✓	✓	✓	✓		✓	
CIDS	✓	✓	✓	✓	✓	✓	✓		✓	✓ ^c
SITA/SIP	✓			✓	✓		✓			✓ ^d
J-SSCG	✓	✓	✓	✓	✓		✓			
IDSA	✓	✓		✓		✓	✓		✓	✓ ^e

Guidelines	RCTs		Outcomes Considered							
	ACTT-2 (NCT04401579)[6]	COV-BARRIER (NCT04421027)[57]	Cost	Mortality	Clinical recovery	Discharge	Safety	Time to ICU admission	Progression to IMV	Other
NIH	✓	✓		✓	✓		✓		✓	
Guidelines Recommending the Use of Baricitinib in Clinical Trials Only										
Government of Mexico	✓	✓		✓			✓			

✓: Guideline explicitly mentions evidence or outcome. Empty cell: Guideline does not explicitly mention evidence or outcome.

Organizations: AWMF: Association of the Scientific Medical Societies in Germany; CIDS: Clinical Infectious Disease Society; IDSA: Infectious Disease Society of America; J-SSCG: Japanese Clinical Practice Guidelines for Management of Sepsis and Septic Shock; NIH: National Institutes of Health; SIP: Italian Society of Pulmonology; SITA: Italian Society of Anti-Infective Therapy; WHO: World Health Organization.

Abbreviations: ICU: intensive care unit; IMV: invasive mechanical ventilation; NCT: National Clinical Trial; NR: not reported; RCT: randomised controlled trial.

Other outcomes considered: ^aDuration of IMV. ^bProgression to new NIV or high-flow oxygen. ^cClinical status at day 15. ^dNeed for NIV. ^eDisease progression (follow up: 28 days), IMV-free days (follow up; 60 days).

Sources: AWMF, October 2021;[24] CIDS, September 2021;[28] COVID-19 Advisory Ontario, January 2022;[32] Government of Mexico, August 2021;[35] IDSA, October 2021;[36] J-SSCG, September 2021;[37] MoH Brazil, May 2021;[38] NICE, January 2022;[8] NIH, December 2021;[39] SITA/SIP, May 2021;[40] WHO, January 2022.[72]

Table 5: Summary of Evidence and Outcomes Reported by Guidelines Discussing Casirivimab/Imdevimab

Guidelines	RCTs		Outcomes Considered							
	REGN-COV 2067 (NCT04425629)[7]	RECOVERY (NCT04381936)[62]	Cost	Mortality	Clinical recovery	Discharge	Safety	Time to ICU admission	Progression to IMV	Other
Guidelines Recommending for the Use of Casirivimab/Imdevimab in At Least One Population Described										
WHO	✓	✓	✓*	Y		✓	✓		✓	✓ ^a
COVID-19 Advisory Ontario		✓		✓		✓	✓		✓**	✓ ^b
AWMF	✓	✓		✓		✓	✓		✓**	✓ ^c
CIDS (seronegative)		✓	✓	✓		✓			✓	✓ ^d
NICE		✓		✓		✓	✓		✓	✓ ^e
Guidelines Recommending the Use of Casirivimab/Imdevimab in Clinical Trials Only										
Government of Mexico				✓			✓			✓ ^f
Guidelines Recommending Against the Use of Casirivimab/Imdevimab in Any Population Described[†]										
MoH Brazil										
SITA/SIP										
NIH										

✓: Guideline explicitly mentions evidence or outcome. Empty cell: Guideline does not explicitly mention evidence or outcome. *Qualitatively considered. **Composite endpoint of need for IMV or death. [†]Includes guidelines which only provided recommendations against casirivimab/imdevimab or recommended against casirivimab/imdevimab for at least one disease severity category and reported insufficient evidence for any other categories.

Organizations: AWMF: Association of the Scientific Medical Societies in Germany; CIDS: Clinical Infectious Disease Society; MoH: Ministry of Health; NICE: National Institute for Health and Care Excellence; NIH: National Institutes of Health; SIP: Italian Society of Pulmonology; SITA: Italian Society of Anti-Infective Therapy; WHO: World Health Organization.


Abbreviations: ICU: intensive care unit; IMV: invasive mechanical ventilation; NCT: National Clinical Trial; NR: not reported; RCT: randomised controlled trial.

Other outcomes considered: ^aAdmission to hospital; time to symptom improvement. ^bTime-weighted average daily change from baseline viral load in nasopharyngeal samples from day 1-7. ^cNeed for dialysis; neurological function; viral clearance. ^dProgression to NIV; progression to organ replacement therapy. ^eWithin 28 days of randomization: Non-invasive mechanical ventilation. ^fReduction in hospitalizations.

Sources: AWMF, October 2021;[24] CIDS, August 2021;[29] COVID-19 Advisory Ontario, November 2021;[31] Government of Mexico, August 2021;[35] IDSA, October 2021;[36] J-SSCG, September 2021;[37] MoH Brazil, May 2021;[38] NICE, January 2022;[8] NIH, December 2021;[39] SITA/SIP, May 2021;[40] WHO, January 2022.[72]


SUPPLEMENTARY MATERIAL

Supplementary Figure 1: Summary of Search Strategy for the COVID-19 Guideline Review



Footnotes: Specific search strings were altered based on the syntax allowed for a specific website (for example, Boolean operators may not be used in certain websites).

Supplementary Figure 2: Flow Chart for COVID-19 Guideline Inclusion/Exclusion



¹Hits were relevant if they were guidelines.

²Priority was given to guidelines which: were applicable to the broadest possible patient population and to an entire country or region, assessed multiple therapies of interest, and directly assessed clinical data.

³Agree-II Domain 3: Rigour of Development.

Supplementary Table 1: International and Country-Specific Sources

Country/Region	Source
All Countries/Regions	Google searches ¹
International	The Guidelines International Network (GIN) International Guideline Library (https://g-i-n.net/)
	Evidence Search (https://www.evidence.nhs.uk/)
	GuidelineCentral (https://www.guidelinecentral.com/)
	The Turning Research Into Practice (TRIP) database (https://www.tripdatabase.com/)
	World Health Organization (WHO) (https://www.who.int/)
	The International Society for Infectious Disease (ISID) (https://isid.org/)
	European Society of Clinical Microbiology and Infectious Diseases (ESCMID) (https://www.escmid.org/)
	European Respiratory Society (ERS) (https://www.ersnet.org/guidelines/)
	British Medical Journal (BMJ) (https://bestpractice.bmj.com/info/us/coronavirus_covid-19)
	Surviving Sepsis Campaign (SSC) (https://www.sccm.org/SurvivingSepsisCampaign/Resources/Resource-Library)
Brazil	Brazilian Ministry of Health (https://www.gov.br/saude/pt-br)
Canada	Clinical Practice Guidelines (CPG) Infobase (https://joulecma.ca/cpg/homepage)
	Government of Canada COVID-19 Website (https://www.canada.ca/en/public-health/services/diseases/coronavirus-disease-covid-19.html)
	Ontario COVID-19 Science Advisory Table (https://covid19-sciencetable.ca/)
France	Haute Autorité de Santé (HAS) (https://www.has-sante.fr/)
	La Société de Réanimation de Langue Française (SRLF) (https://www.srlf.org/recommandations-referentiels-epp)
	Ministère des Solidarités et de la Santé (https://solidarites-sante.gouv.fr/)
	Haut Conseil de la Santé Publique (HCSP) (https://www.hcsp.fr/Explore.cgi/Accueil)
Germany	Das Portal der wissenschaftlichen Medizin (https://www.awmf.org/leitlinien/aktuelle-leitlinien.html#FA)
	Robert Koch Institut (RKI) (https://www.rki.de/DE/Content/InfAZ/N/Neuartiges_Coronavirus/nCoV.html)

Country/Region	Source
	Federal Ministry of Health (https://www.bundesgesundheitsministerium.de/)
India	Ministry of Health and Family Welfare (MoHFW) (https://www.mohfw.gov.in/)
	Clinical Infectious Diseases Society (CIDS) (http://www.cidsindia.org/)
	Indian Council of Medical Research (ICMR) (https://www.icmr.gov.in/)
Italy	AGENAS National Agency for Regional Health Services (https://www.agenas.gov.it/)
	Agenzia Italiana del Farmaco (AIFA) (https://www.aifa.gov.it/)
	Ministry of Health (https://www.salute.gov.it/portale/nuovocoronavirus/homeNuovoCoronavirus.jsp)
	Società Italiana di Malattie Infettive e Tropicali (SIMIT) (https://www.simit.org/)
Japan	Ministry of Health, Labour and Welfare (MHLW) (https://www.mhlw.go.jp/index.html)
	The Japanese Association for Infectious Diseases (JAID) (https://www.kansensho.or.jp/)
Mexico	Mexican Ministry of Health (https://coronavirus.gob.mx/)
Turkey	Türkiye Enfeksiyon Hastalıkları ve Klinik Mikrobiyoloji Uzmanlık Derneği (EKMUD) (https://ekmud.org.tr/)
	Ministry of Health (https://www.saglik.gov.tr/?_Dil=1)
	COVID-19 Data Portal (https://covid19.tubitak.gov.tr/)
UK	National Institute for Health and Care Excellence (NICE) (https://www.nice.org.uk/)
	UK Department of Health & Social Care (https://www.gov.uk/government/organisations/department-of-health-and-social-care)
US	Centers for Disease Control and Prevention (CDC) (https://www.cdc.gov/)
	Infectious Diseases Society of America (IDSA) (https://www.idsociety.org/)

Abbreviations: UK, United Kingdom; US, United States.

¹ For all countries and regions, Google searches were conducted to supplement country-specific websites and guideline repositories.

Supplementary Table 2: Search Terms by Language

Language	Search Terms
English	<ul style="list-style-type: none"> • Guideline • COVID-19 • Baricitinib • Dexamethasone • Remdesivir • REGEN-COV • Casirivimab and imdevimab • Tocilizumab • [Country of interest]
French	<ul style="list-style-type: none"> • Ligne directrice • COVID-19 • Baricitinib • Dexaméthasone • Remdesivir • REGEN-COV • Casirivimab et imdevimab • Tocilizumab • [France]
German	<ul style="list-style-type: none"> • Leitlinie • COVID-19 • Baricitinib • Dexamethason • Remdesivir • REGEN-COV • Casirivimab und Imdevimab • Tocilizumab • [Deutschland]
Italian	<ul style="list-style-type: none"> • Linee guida • COVID-19 • Baricitinib • Desametasone • Remdesivir • REGEN-COV • Casirivimab e imdevimab • Tocilizumab • [Italia]
Japanese	<ul style="list-style-type: none"> • ガイドライン • 新型コロナウイルス • COVID-19 • バリシチニブ • デキサメタゾン • レムデシビル • REGEN-COV • カシリビマブとイムデビマブ • トシリズマブ • [日本]
Portuguese	<ul style="list-style-type: none"> • Diretriz • COVID-19 • Baricitinibe • Dexametasona • Remdesivir • REGEN-COV • Casirivimab e imdevimab • Tocilizumab • Brasil
Spanish	<ul style="list-style-type: none"> • Directriz • COVID-19 • Baricitinib • Dexametasona • Remdesivir • REGEN-COV • Casirivimab e imdevimab • Tocilizumab

Language	Search Terms
	<ul style="list-style-type: none">• [Mexico]
Turkish	<ul style="list-style-type: none">• Rehber• COVID-19• Barisitinib• Deksmetazon• Remdesivir• REGEN-COV• Casirivimab ve imdevimab• Tocilizumab• [Türkiye]

Supplemental Table 3: Eligibility Criteria

Category	Inclusion Criteria	Exclusion Criteria
Population	<ul style="list-style-type: none"> Patients with a primary diagnosis of COVID-19 	<ul style="list-style-type: none"> Patients without a primary diagnosis of COVID-19
Interventions	<p>Includes recommendations regarding at least one of the following therapies:</p> <ul style="list-style-type: none"> Baricitinib Dexamethasone Remdesivir Casirivimab/imdevimab Tocilizumab 	<p>Does not include recommendations regarding any of the following therapies:</p> <ul style="list-style-type: none"> Baricitinib Dexamethasone Remdesivir Casirivimab/imdevimab Tocilizumab
Comparators	<ul style="list-style-type: none"> Any or none 	<ul style="list-style-type: none"> N/A
Outcomes	<ul style="list-style-type: none"> Guidelines must discuss the clinical management of COVID-19 in hospitalized patients (including ICU) 	<ul style="list-style-type: none"> Guidelines that do not discuss the clinical management of hospitalized patients with COVID-19 Guidelines that focus on the management of conditions other than COVID-19
Publication type	<ul style="list-style-type: none"> Official clinical practice guidelines or treatment guidelines 	<ul style="list-style-type: none"> Publications other than official clinical practice guidelines or treatment guidelines
Other considerations	<ul style="list-style-type: none"> Specifically produced for use in any of the countries of interest (Brazil, Canada, France, Germany, India, Italy, Japan, Mexico, Turkey, UK or the US) International guidelines 	<ul style="list-style-type: none"> Guidelines not specifically produced for use in the countries of interest or on an international level

Abbreviations: COVID-19, coronavirus disease 2019; ICU, intensive care unit; N/A, not applicable; UK, United Kingdom; US, United States.

Supplementary Table 4: AGREE-II Domains and Items

Domain	Item No.	Item Title
Domain 1: Scope and Purpose	1	The overall objective(s) of the guideline is (are) specifically described
	2	The health question(s) covered by the guideline is (are) specifically described
	3	The population (patients, public, etc.) to whom the guideline is meant to apply is specifically described
Domain 2: Stakeholder Involvement	4	The guideline development group includes individuals from all relevant professional groups
	5	The views and preferences of the target population (patients, public, etc.) have been sought
	6	The target users of the guideline are clearly defined
Domain 3: Rigour of Development	7	Systematic methods were used to search for evidence
	8	The criteria for selecting the evidence are clearly described
	9	The strengths and limitations of the body of evidence are clearly described
	10	The methods for formulating the recommendations are clearly described
	11	The health benefits, side effects, and risks have been considered in formulating the recommendations
	12	There is an explicit link between the recommendations and the supporting evidence
	13	The guideline has been externally reviewed by experts prior to its publication
	14	A procedure for updating the guideline is provided
Domain 4: Clarity of Presentation^a	15	The recommendations are specific and unambiguous
	16	The different options for management of the condition or health issue are clearly presented
	17	Key recommendations are easily identifiable
Domain 5: Applicability	18	The guideline describes facilitators and barriers to its application
	19	The guideline provides advice and/or tools on how the recommendations can be put into practice
	20	The potential resource implications of applying the recommendations have been considered
	21	The guideline presents monitoring and/or auditing criteria
Domain 6: Editorial Independence	22	The views of the funding body have not influenced the content of the guideline
	23	Competing interests of guideline development group members have been recorded and addressed

Footnote: ^aFor this domain, this did not include details of the specific content of the recommendations themselves, as this was captured as part of the full extractions for the high-quality guidelines.

Supplementary Table 5: Summary of AGREE-II Quality Assessments (Standardized Domain Scores)*

Organization	Domain 1 Scope and Purpose	Domain 2 Stakeholder Involvement	Domain 3 Rigour of Development	Domain 4 Clarity of Presentation	Domain 5 Applicability	Domain 6 Editorial Independence	Overall	Key Weaknesses
AWMF (Germany, 2021)	83%	94%	91%	97%	40%	100%	100%	-
SITA/SIP (Italy, 2021)	94%	67%	91%	78%	15%	79%	83%	-
CIDS (India, 2021)	47%	64%	88%	100%	58%	67%	83%	-
IDSA (US, 2021)	92%	67%	86%	100%	33%	100%	75%	Lacks monitoring criteria
WHO (International, 2021)	83%	100%	83%	100%	79%	71%	100%	-
J-SSCG (Japan, 2021)	100%	67%	76%	94%	35%	88%	83%	-
ERS (International, 2021)	100%	53%	74%	100%	73%	58%	83%	-
SSC (International, 2021)	100%	64%	66%	100%	63%	100%	83%	-
MoH (Brazil, 2021)	100%	52%	62%	86%	58%	21%	75%	Lacks conflict of interest statement and funding source
NIH (US, 2021)	61%	67%	60%	100%	54%	46%	92%	-
NICE (UK, 2021)	94%	53%	57%	100%	58%	17%	75%	Lacks conflict of interest statement; limited detail for methods; lacked systematic approach
Government of Mexico (Mexico, 2021)	81%	50%	57%	100%	19%	8%	67%	Lacks external review, information regarding facilitators/barriers, implementation, source implication and funding statement
COVID-19 Advisory Ontario (Canada, 2021)	97%	42%	50%	100%	73%	33%	83%	-
EKMUD Non-Antiviral (Turkey, 2021)	8%	19%	24%	36%	0%	0%	17%	Not systematic or methodical; recommendations are ambiguous
HCSP (France, 2021)	17%	39%	23%	78%	4%	42%	50%	Lacks methodology for gathering evidence and setting recommendations; lacks clear target users and target population
COVRIIN (Germany, 2021)	28%	25%	22%	83%	4%	0%	42%	Lacks detail on methodology

Organization	Domain 1 Scope and Purpose	Domain 2 Stakeholder Involvement	Domain 3 Rigour of Development	Domain 4 Clarity of Presentation	Domain 5 Applicability	Domain 6 Editorial Independence	Overall	Key Weaknesses
FADOI (Italy, 2021)	53%	31%	20%	56%	0%	0%	17%	Lacks systematic methods and clarity on methods, conflict of interest and funding
JAID (Japan, 2021)	39%	11%	15%	39%	42%	0%	33%	Lacks methodology
MHLW (Japan, 2021)	31%	36%	15%	39%	42%	0%	33%	Lacks methodology
STAKOB (Germany, 2021)	36%	14%	14%	69%	27%	0%	50%	Lacks methodology and discussion of strength of evidence used
MoH (Turkey, 2021)	25%	19%	13%	89%	13%	0%	42%	Lacks details on methods, links to evidence, external review, conflict of interest statement, funding statement
EKMUD Antiviral (Turkey, 2021)	14%	3%	13%	61%	4%	0%	17%	Not systematic or methodical; recommendations are ambiguous
Government of India MHFW (India, 2021)	14%	0%	3%	83%	25%	0%	17%	Lacks explicit links to evidence, methodology, conflicts of interest, funding information

*Table is based on the guidelines identified in the initial searches (August 2021) and does not include updates; guidelines are ordered by Domain 3 standardized score; greyed out rows denote guidelines which scored <50% in Domain 3 and therefore were not considered to be of high quality.

Abbreviations: AWMF: Association of the Scientific Medical Societies in Germany; CIDS: Clinical Infectious Disease Society; COVID-19: coronavirus disease 2019; COVRIIN: Specialist Group of Intensive Care Medicine, Infectious Diseases and Emergency Medicine; EKMUD: Infectious Diseases & Clinical Microbiology Specialty Society of Turkey; ERS: European Respiratory Society; FADOI: Federation of Associations of Internist Hospital Managers; HCSP: High Council of Public Health; IDSA: Infectious Diseases Society of America; JAID: Japanese Association for Infectious Diseases; J-SSCG: Japanese Clinical Practice Guidelines for Management of Sepsis and Septic Shock; MHFW: Ministry of Health and Family Welfare; MHLW: Ministry of Health, Labour and Welfare; MoH: Ministry of Health; NICE: National Institute for Health and Care Excellence; NIH: National Institutes of Health; SIP: Italian Society of Pulmonology; SITA: Italian Society of Anti-Infective Therapy; SSC: Surviving Sepsis Campaign; STAKOB: Standing Working Group of Competence and Treatment Centers for Highly Contagious and Life-Threatening Diseases; WHO: World Health Organization.

Supplementary Table 6: Overview of High-Quality Guidelines and Recommendations*

Organization (Country, Date)	Mild (Hospitalized; no supplemental oxygen)	Moderate (Hospitalized; low-flow supplemental oxygen)	Severe (Hospitalized; NIV or high-flow oxygen)	Critical (Hospitalized; IMV or ECMO)
ERS (International, April 2021)[10]	RV DX TZ	RV DX TD	RV DX TD	RV DX TD
SSC (International, March 2021)[41]			RV DX	RV DX
WHO (International, January 2022)[9]	RV DX CI	RV DX	CI RV DX TZ BA	CI RV DX TZ BA CI
MoH Brazil (Brazil, May 2021)[38]	RV DX CI	RV DX	CI RV DX TZ	CI RV DX TZ CI
COVID-19 Advisory Ontario (Canada, August 2021)[30–34]	BA CI	RV DX TD BA	CI RV DX TD	CI RV DX TD BA CI
AWMF (Germany, October 2021)[24]	RV DX TZ BA CI	RV DX TZ BA	CI RV DX TD	RV DX TZ
CIDS (India, September 2021)[25–29]	RV DX TZ BA	RV DX TZ BD	CI RV DX TD BD	CI RV DX TD BD CI
SITA/SIP (Italy, May 2021)[40]	CI	RV DX	CI RV DX TD BR	CI
J-SSCG (Japan, September 2021)[37]	DX	RV DX TZ BA	RV DX TZ BA	RV DX TZ BA CI
Government of Mexico (Mexico, August 2021)[35]	DX BR CI	RV DX TD BR	CI RV DX TD BR	CI DX TD BR CI
NICE (UK, January 2022)[8]	RV DX CI	RV DX TD	RV DX TD DR	CI RV DX TD DR CI
IDSA (US, November 2021)[36]	RV DX	RV DX	RV DX TD BD BR	RV DX TD
NIH (US, December 2021)[8]	RV DX BD CI	RV DX BD DR CI	RV DX TZ BA DR BD BT BDR CI	RV DX TD BT BD BDR CI

○ Monotherapy
 □ Combination therapy
 ● Recommended
 ● Not recommended
 ● Recommended only in some sub-populations
 ● Insufficient evidence or clinical trials only

*This table provides a simplified overview of recommendations only; for more comprehensive details, see subsequent tables within the results section of this article or refer to the original guideline documents.

Therapies: BA: baricitinib; BD: baricitinib + dexamethasone; BDR: baricitinib + dexamethasone + remdesivir; BR: baricitinib + remdesivir; BT: baricitinib + tocilizumab; CI: casirivimab + imdevimab; DR: dexamethasone + remdesivir; DX: dexamethasone; RV: remdesivir; TD: tocilizumab + dexamethasone; TZ: tocilizumab.

Organizations: AWMF: Association of the Scientific Medical Societies in Germany; CIDS: Clinical Infectious Disease Society; ERS: European Respiratory Society; IDSA: Infectious Diseases Society of America; J-SSCG: Japanese Clinical Practice Guidelines for Management of Sepsis and Septic Shock; MoH: Ministry of Health; NICE: National Institute for Health and Care Excellence; NIH: National Institutes of Health; SIP: Italian Society of Pulmonology; SITA: Italian Society of Anti-Infective Therapy; SSC: Surviving Sepsis Campaign; WHO: World Health Organization.