

Early View

Original article

Is Laryngeal Hypersensitivity the Basis for Unexplained or Refractory Chronic Cough?

Krishna M. Sundar, Amanda Carole Stark, Nan Hu, Julie Barkmeier-Kraemer

Please cite this article as: Sundar KM, Stark AC, Hu N, *et al.* Is Laryngeal Hypersensitivity the Basis for Unexplained or Refractory Chronic Cough?. *ERJ Open Res* 2021; in press (<https://doi.org/10.1183/23120541.00793-2020>).

This manuscript has recently been accepted for publication in the *ERJ Open Research*. It is published here in its accepted form prior to copyediting and typesetting by our production team. After these production processes are complete and the authors have approved the resulting proofs, the article will move to the latest issue of the ERJOR online.

Is Laryngeal Hypersensitivity the Basis for Unexplained or Refractory Chronic Cough?

An observational study of cough hypersensitivity and laryngeal abnormalities in chronic cough using a novel interdisciplinary approach

Krishna M. Sundar^{1*} MD

Amanda Carole Stark^{2,3*} M.S., CCC-SLP

Nan Hu, PhD⁴

Julie Barkmeier-Kraemer^{3,5} PhD., CCC-SLP

Departments of Internal Medicine¹, National Center for Voice and Speech², Department of Rehabilitation³, Department of Otolaryngology⁵ University of Utah, Department of Biostatistics, Robert Stempel College of Public Health and Social Work and Division of Public Health, Florida International University, Miami, Florida, USA⁴, ,

* Authors contributed equally to the manuscript

Krishna M. Sundar, MD – Professor (Clinical), Department of Medicine, University of Utah, 375, Chipeta Way, Salt Lake City, UT 84108; krishna.sundar@hsc.utah.edu

Amanda Carole Stark, SLP – Research Associate, National Center for Voice and Speech, University of Utah, 2901 S Campus Dr Suite 2120, Salt Lake City, UT 84112.

Amanda.stark@hsc.utah.edu

Nan Hu, PhD, Associate Professor, Department of Biostatistics, 11200 SW 8th Street AHC5 Miami, FL 33199; nhu@fiu.edu

Julie Barkmeier-Kraemer, PhD, Professor, Department of Otolaryngology, 50 North Medical Dr, SOM3C120, Salt Lake City, UT 84132; JulieB.Kraemer@hsc.utah.edu

Corresponding author – Krishna M. Sundar, MD; Professor (Clinical), Department of Medicine, University of Utah. Phone – 801-581-2016; Fax – 801587-3349; krishna.sundar@hsc.utah.edu

TAKE HOME SUMMARY

Chronic refractory cough is frequently characterized by cough hypersensitivity and laryngeal dysfunction. An interdisciplinary team approach that includes laryngeal evaluation can identify laryngeal hypersensitivity along with other cough aetiologies.

ABSTRACT

Background – Refractory chronic cough (RCC) and unexplained chronic cough (UCC) are common problems seen in primary care and subspecialty clinics. The role of cough hypersensitivity and laryngeal dysfunction in contributing to the persistence of cough in RCC/UCC is not well recognized.

Methods – Data of patients with RCC and UCC evaluated in 2019 by an interdisciplinary cough clinic led by a pulmonologist and speech-language pathology team was reviewed. Patients completed validated questionnaires including the Leicester cough questionnaire (LCQ), voice handicap index (VHI) and dyspnea index questionnaire (DI) at initial encounter. Presence of cough hypersensitivity was based upon a history of allotussia and hypertussia. Laryngeal dysfunction was diagnosed in those with a history of laryngeal paresthesias, throat clearing, voice abnormalities, upper airway dyspnoea and documentation of functional or anatomic laryngeal abnormalities on nasoendoscopy.

Findings – Of the 60 UCC/RCC patients analyzed, 75% of patients were female and 85% were over 40 years of age. Cough hypersensitivity was documented in all patients and multiple cough triggers occurred in 75% of patients. 95%, 50% and 25% of patients reported laryngeal paresthesias, voice abnormalities and upper airway dyspnoea, respectively. Significant associations between LCQ and VHI and DI scores occurred when adjusting for age, gender, ethnicity and body mass index. Laryngeal functional abnormalities were documented on 44/60 patients on nasoendoscopy.

Interpretation – Hypertussia, allotussia and laryngeal dysfunction are common in patients with RCC and UCC. Evaluation of UCC and RCC can delineate laryngeal hypersensitivity and allows appropriate treatment to be directed at this phenotype.

Funding – None

INTRODUCTION

Chronic cough is one of the most common clinical problems encountered in primary care and subspecialty clinics [1]. In the majority of patients, the underlying aetiology is thought to be gastroesophageal reflux disease (GERD/GORD), rhinosinusitis or cough variant asthma (CVA) when an empiric diagnostic approach to evaluate chronic cough is used [2]. While this approach to chronic cough attempts to link specific causal aetiologies to the occurrence of cough, a substantial proportion of patients persist with their cough despite conventional treatment(s) for these conditions [3]. Terminology such as unexplained chronic cough (UCC), refractory chronic cough (RCC) and idiopathic cough highlight the diagnostic and therapeutic problems in 5-42% of chronic cough patients [3].

To understand the persistence of cough in such individuals, the phenomenon of cough hypersensitivity has been proposed [4]. Cough hypersensitivity is characterized by decreased cough thresholds to noxious and non-noxious stimuli [5]. In addition to abnormal triggering of cough, abnormalities in laryngeal sensation and function are frequently reported in patients with UCC and RCC [6].

This study tested the hypothesis that symptoms suggestive of laryngeal dysfunction and cough hypersensitivity underlie the clinical symptomatology of UCC/RCC. To test this hypothesis, patient data from an interdisciplinary team-based evaluation of UCC/RCC patients was analyzed. The specific aims of this study included determination of the following:

1. Frequency of cough hypersensitivity symptoms documented during an initial clinical encounter in those diagnosed with RCC or UCC.
2. Frequency of coexisting voice, breathing or swallowing abnormalities in those with RCC or UCC using validated questionnaires.

3. Frequency of laryngeal dysfunction documented during nasoendoscopy examination in those with RCC or UCC.

METHODS

Study design, setting and patients

This was a one-year cohort study conducted according to the STROBE statement on recommendations for observational studies [7]. Patients were identified for inclusion in this study based on an electronic medical record review of all individuals seen in the interdisciplinary Cough Clinic at the University of Utah Voice Disorders Center from January 1, 2019 to December 31, 2019. Individuals with chronic cough are referred to this clinic from primary care providers, gastroenterologists, pulmonologists, allergists and otolaryngologists within the University of Utah Health system, or from local or regional locations. Candidates for this study were excluded if they were smokers, or diagnosed with the following: a) chronic lung disease, b) congestive heart failure, c) neurological conditions including Parkinson's disease or stroke, d) head and neck cancer, or e) underwent prior laryngeal surgery. All patients included had normal chest radiology and pulmonary function tests (PFTs) or had abnormalities that did not contribute to the occurrence of chronic cough as determined by the team pulmonologist.

Data Sources

All patients were evaluated by an interdisciplinary team that included a pulmonologist (KMS) and speech-language pathology (SLP) team members clinically specialized in voice, cough/airway, and swallowing disorders. Abnormal nasoendoscopy findings were reviewed by a laryngologist during a weekly multidisciplinary voice panel. Prior to the initial visit at the cough clinic, patients completed validated quality of life self-report questionnaires. These included:

- 1) Leicester Cough Questionnaire (LCQ) [8],
- 2) Voice handicap index (VHI) [9],
- 3) Dyspnea Index (DI) [10]
- 4) Cough visual analog scale score (VAS).

Instrumental voice and airway assessment was done per the American Speech Language and Hearing Association (ASHA) recommended protocol [11]. Data extracted from electronic medical records included patient demographics, cough characteristics, treatments rendered, speech assessment results and nasoendoscopic imaging findings.

Variables measured

Cough Hypersensitivity: In this study, the presence of cough hypersensitivity was defined by the report of coughing in response to stimuli that typically do not lead to cough (allotussia), and exaggerated coughing in response to typical stimuli that elicit cough at lower doses (hypertussia) [12]. Thus, all reported triggers of coughing such as strong smells/perfumes, changes in body position, eating, temperature change, exercise, talking etc. were included in the analysis.

Laryngeal dysfunction: Categorization of symptoms and signs of laryngeal dysfunction was done as below

1. *Laryngeal paresthesias* – report of globus sensation, a tickle or irritation in the throat.
2. *Presence or absence of throat clearing*
3. *Laryngeal sensorimotor abnormality* as indicated by self-reported symptoms of dysphonia, upper airway dyspnoea, and dysphagia indicating impaired laryngeal function (e.g. aspiration, or frequent occurrence of coughing relative to swallowing).

4. *Anatomic or physiologic laryngeal abnormalities* – based on nasoendoscopic examination reported by the treating SLP and confirmed by a laryngologist when a structural, or neurological abnormality of the larynx was present.

Outcomes from therapies rendered

This study did not aim to assess outcomes from therapies rendered since objective outcome measures (e.g., VAS or LCQ total score) were not consistently available during follow-up visits across patients in this study. Patients were commonly treated with behavioural speech therapy (BST) based upon defined treatment protocols [13]. Patients also underwent screening and evaluation for obstructive sleep apnoea (OSA) that was based upon existing guidelines [14]. Patient outcomes following treatment were determined based upon subjective reports of improvement during follow-up visits or phone calls.

Statistical analyses

Data were summarized as mean and standard deviation (SD) for continuous variables, and frequency (n) and percentage (%) for categorical variables. Univariate linear regression was used to examine the association between continuous outcomes (LCQ, VAS, VHI, DI and number of triggers) and single predictor variables (laryngeal paresthesia, and throat clearing). Univariate logistic regressions were performed to associate the binary outcomes (laryngeal paresthesia, throat clearing and cough triggers) with LCQ, VHI and DI as the predictors, and odds ratios were reported. Multivariate linear regression and logistic regression models were performed that adjusted for age, sex, body mass index (BMI), and ethnicity.

Analyses were performed using statistical software Stata (StataCorp., College Station, TX, US)

V. 15. All tests were two-sided, and p-values < 0.05 indicate statistically significant results.

RESULTS

Patient Characteristics

85 patients with chronic cough were seen at the University of Utah interdisciplinary Cough Clinic during the study timeline. Of these, 25 were excluded from further analysis due to comorbidities or other exclusion criteria (Figure 1). Mild restriction on PFTs was seen in 4/60 patients and 3/60 had radiologic abnormalities (lung nodules (2) and small cysts (1)) that were not felt to contribute to the chronic cough and therefore, these patients were included in the final data analysis.

Cough characteristics

Table 1 outlines patient demographics and concomitant diagnoses related to their chronic cough. Nearly 75% of patients were female, with 85% of individuals above 40 years of age. Patients above 70 years comprised 1/3rd of patients in the study and had the longest durations of cough (Tables 1 & 2). 40% of patients were obese (BMI > 30 kg/m²) and less than a third of patients (17/60) had normal BMI of ≤25 kg/m² with the rest of subjects in the overweight category. Cough duration of ten or more years was reported in 10/60 patients, but cough duration was not significantly associated with patients' age as a continuous variable (p=0.068). Mean duration of cough was not significantly longer in men than women (87.1 months vs. 71.5 months; p=0.61). Cough was rated as moderate to severe based on VAS scores (63.79 mm ±0.47) and total LCQ scores (9.91 ±2.75). Cough duration was not predictive of total LCQ or VAS scores (p=0.609 and 0.069). Paroxysms of cough were reported by the majority of patients,

although more frequently in those less than 70 years of age. There was no effect of age, sex, BMI, cough duration and LCQ on the occurrence of paroxysms of cough.

Of the 1/3rd of patients (20/60) experiencing cough onset with a presumed respiratory tract infection (RTI), females reported an RTI more frequently than men (OR 1.71; 95% CI: 0.47-6.21); however, this difference was not significantly different ($p=0.412$). Another sixth of patients reported events such as exposure to strong chemicals, acute pulmonary conditions (such as pulmonary embolism), periods of excessive voice use, and life stressors at the onset of chronic cough.

Even though the majority of patients reported previous active treatments for GERD, UACS and rhinosinusitis, only cough aetiologies ascertained at the time of initial clinic visit were included in Table 1. At the time of initial evaluation, 75% of patients did *not* have symptoms to suggest one or more of the three aetiologies of GERD, UACS and rhinosinusitis. Only 15/60 patients reported one or more of these three disorders to be an ongoing problem contributing to cough. Particularly for GERD, 42/60 patients were previously treated or were on treatment with anti-acid therapies (proton-pump inhibitors and/or histamine-2 receptor blockers) for months to years without improvement in cough.

Cough hypersensitivity

Cough hypersensitivity was determined based on symptoms of allotussia and hypertussia. Cough triggers ranged from talking to strong perfumes with talking being the most common trigger for cough (Figure 2). The amount of talking required to trigger cough varied between patients and was not quantified within clinical records. Patients experiencing coughing from multiple triggers were noted in 75% (45/60) of patients. No significant relationships occurred

between the number of cough triggers and LCQ scores, presence of laryngeal paresthesia, frequent throat clearing, VHI or DI scores.

Laryngeal dysfunction

The majority of individuals reported abnormal laryngeal sensations (57/60) described as tickle, globus (lump or sensation in the throat), irritation, dry throat, sore throat and presence of mucus in the throat. Multiple different descriptors were noted in 10/60 patients. Women reported a higher prevalence of abnormal throat sensations that were associated with increasing age. Nearly half of the patients reported frequent throat clearing. Presence of throat clearing was not related to LCQ or other signs or symptoms of laryngeal dysfunction.

Voice abnormalities were reported in more than half the patients that included hoarseness, loss of voice with talking or singing, change in pitch or a raspy voice quality. One fourth of patients reported symptoms of upper airway dyspnoea such as throat constriction and difficulty getting in air when dyspnoeic. Using the DI score index, 75% of patients reported abnormal scores and VHI total scores were abnormal in 43% of patients. A significant unadjusted-association (when not adjusting for age, gender and BMI) was shown between total scores on the LCQ and VAS, and between VHI and DI scores. Higher VHI total scores were associated with worse (i.e., lower) LCQ scores (coefficient= -3.4 change in LCQ) when adjusting for age, gender, ethnicity and BMI ($p=0.001$, $R^2 = 0.2536$). Unadjusted DI scores were statistically associated with LCQ scores such that one unit of decrease in LCQ was associated with an increase in DI scores by .89 points ($p=0.022$).

Laryngeal abnormalities – structural and functional

Laryngeal functional abnormalities were categorized based upon the findings of abnormal levels of hyperfunction (anteroposterior, mediolateral, or sphincteric compression i.e. simultaneous anteroposterior and mediolateral compression of vocal folds, aryepiglottic folds and ventricular folds) of the larynx at rest, during phonation, or during breathing tasks. Functional abnormalities were documented in the majority of patients (44/60) (Figure 3) although paradoxical vocal fold motion (PVFM) was only documented in 6/60 patients. Structural abnormalities were noted in nearly half of the patients with the majority of these being erythema and oedema (Figure 3). Neurological vocal fold abnormalities were indicated in 9/60 (sluggish or immobile vocal folds (n=4) and vocal tremor (n=5)). Only five patients had normal laryngeal appearance and function and 24/60 patients exhibited overlapping structural and functional abnormalities. The occurrence of neurologic or structural abnormalities (seen in 1/5th of patients in our study) did not influence the treatment for cough.

Treatment

Besides specific treatments for GERD, UACS and CVA, treatment recommendations for cough included behavioural speech therapy (BST) and management of comorbid obstructive sleep apnoea (OSA). BST was recommended for 56/60 patients for treatment of their cough, and any other upper airway or voice complaints (Figure 1). 34/56 patients completed their BST while OSA management was completed in 43% with 22 new diagnoses of OSA rendered. Combinations of different therapies included BST, continuous positive airway pressure therapy (CPAP) for OSA, and management of GERD, UACS, rhinosinusitis with improvement in cough noted in 34/40 (85.0%) patients that adhered to recommended therapies (Figure 1).

DISCUSSION

This observational study in patients with UCC and RCC determined that a high proportion of those with cough hypersensitivity (i.e. allotussia and/or hypertussia) exhibited signs or symptoms of laryngeal dysfunction. This study was unique for its interdisciplinary approach that combined the empiric clinical approach based upon the 2006 chronic cough guidelines with a comprehensive voice and airway evaluation. While an interdisciplinary approach to chronic cough has the potential for lower costs and healthcare utilization during assessment [15], there are no studies reporting clinical outcomes associated with such a combined team-based approach. The team-based collaboration between a pulmonologist and SLP specializing in voice and airway populations pools complimentary skills in the evaluation and management of cough. For physicians dealing with chronic cough, the assessment of the voice and upper airway by an SLP and subsequent BST has the potential to lead to faster resolution of chronic cough.

Cough is a vital protective reflex preventing aspiration and the abnormal triggering of cough with activities such as eating, talking, changes in position etc. is indicative of a sensitized cough reflex or cough hypersensitivity [16]. Cough hypersensitivity syndrome is defined as “a condition characterized by cough triggered by low levels of thermal, mechanical, or chemical exposure” [5]. Using this definition, we utilized the symptoms of allotussia and hypertussia as indicative of cough hypersensitivity and this was demonstrable in all our patients with RCC/UCC. Cough hypersensitivity can also be assessed objectively by measuring sensitivity to inhaled capsaicin given that capsaicin sensitivity is abnormal in patients with chronic cough irrespective of phenotype (UCC, GERD, CVA, rhinosinusitis) and improves with cough resolution [17-19]. However, the use of inhalational capsaicin challenge to assess cough hypersensitivity has wide diagnostic cut-offs and is not widely available. Capsaicin activates the

“nociceptor” cough pathway by binding to TRPV1 airway receptors [17]. Recent insights into mechanistic heterogeneity of cough point to the possibility of different endotypes of chronic cough [20] in which TRPV1 activation may not be a consistent abnormality. The persistence of cough in studies of novel TRPV1 receptor blockers highlights the need to recognize cough hypersensitivity beyond abnormal capsaicin sensitivity [20]. All UCC/RCC patients in this study reported one or more cough triggers with multiple triggers occurring in 75% of patients. The most common trigger for cough in UCC/RCC patients in this study was phonation (i.e. talking and laughing) as was seen in a previous study [6]. Therefore, given the hallmark finding of abnormal cough triggering in this and other studies of UCC/RCC of coughing to multiple stimuli [6, 21-28] (Table 4) the history of allotussia and hypertussia can be considered a valuable clinical correlate for cough hypersensitivity.

Besides allotussia and hypertussia, there are other phenotypic attributes of cough hypersensitivity that are common across studies of UCC/RCC (Table 4) [6, 21-28]. Complaints most commonly seen with cough hypersensitivity include irritation in the throat or chest, throat clearing, dysphonia/hoarseness, upper airway dyspnoea, and globus – symptoms that may indicate a significant contribution from laryngeal structures toward cough hypersensitivity [29]. Amongst these sensations, the urge to cough or somatic sensations that precede the occurrence of cough have received the most attention [30]. In most studies evaluating this symptom, these sensations are localized to the neck and ascribed by clinicians as “laryngeal paresthesias” or laryngeal sensations [6, 22, 26, 28]. These laryngeal paresthesias are often present without any evident cough stimulus in patients with UCC/RCC [25]. 95% of our UCC/RCC patients reported symptoms consistent with laryngeal paresthesias compatible with the high prevalence noted in other studies (Table 4). In addition to laryngeal paresthesias, other signs or symptoms of

laryngeal dysfunction included frequent throat clearing, voice abnormalities and dyspnoea, and observation of functional abnormalities during nasoendoscopy such as paradoxical vocal fold motion (PVFM) or muscle-tension dysphonia (Table 4).

Cough hypersensitivity in RCC/UCC is attributed to neuropathic changes in cough pathways within the central and peripheral limbs of cough reflex pathways. Both peripheral sensitization and alteration in the modulating influences of central descending pathways are implicated in the causation of cough [20, 31]. Spatial and functional separation of peripheral cough afferents into chemosensory proximal airway neurons projecting to jugular vagal ganglia vs. distal mechanosensitive neurons projecting to nodose vagal ganglia offer a possible mechanistic explanation for the diversity seen in the cough reflex pathway. However, much remains to be understood in terms of the neuro-structural or neuro-functional abnormalities underlying chronic cough [31]. Despite this lack of information of the spatial localization of pathways driving cough hypersensitivity, there is increasing clinical recognition of the role of the larynx and surrounding structures in chronic cough.

The cough reflex involves the coordinated action of a number of different muscles of the chest and airway with the larynx playing a key role in the compression and expiratory phases of cough [32]. The rich innervation of the larynx enables response to tactile, chemical, and mechanical stimuli, either with cough or forceful vocal fold adduction (expiratory reflex) [33]. Laryngeal dysfunction manifesting as cough can be triggered by viral infection and perpetuated by further trauma due to throat clearing and coughing [5, 34]. Terms to describe such laryngeal dysfunction driving cough include “laryngeal hypersensitivity” and “laryngeal hyperresponsiveness” in the context of abnormal respiratory sensation [23] and/or abnormal vocal fold adduction during inspiration [27]. Whether laryngeal hypersensitivity is the only

clinical correlate of cough hypersensitivity or whether other symptoms such as chest irritation represent different expressions of cough hypersensitivity is unclear [29].

The role of laryngeal dysfunction in RCC is poorly appreciated despite reports of a wide spectrum of laryngeal abnormalities in patients with RCC (Table 4). Conversely, a higher prevalence of cough is noted in patients with PVFM, muscle tension dysphonia (MTD), and vocal cord dysfunction (VCD) [6, 22, 27]. Despite categorization of patients into those with predominant dyspnoea (VCD), voice (MTD) and cough problems, the considerable overlap in abnormalities indicates the laryngeal airway as a major source for these syndromes [25, 27]. Additionally, correlations between abnormalities of laryngeal and cough indices such as the LCQ highlight the potential for the vagal laryngeal afferents as the salient neuropathic substrate of cough hypersensitivity [27, 28]. In our study, the relationship between DI and VHI with LCQ was significant and indicated that the occurrence of cough due to somatic stimuli that affect the larynx such as talking, eating, changes in position, deep breathing with position can track with upper airway dyspnoea and voice problems. However correlation does not infer causality. A recent study by Vertigan et al. that systematically assessed patients with RCC, MTD and VCD demonstrated both a relationship between the laryngeal hypersensitivity questionnaire (LHQ) score and cough frequency along with similar impairments in voice scores and laryngoscopic abnormalities [27]. While PVFM was only noted in a small number of patients in our study due to inconsistent use of provocation techniques, a significant prevalence of PVFM in RCC has been reported in other studies [21, 27]. Another recent study on 416 chronic cough and 62 RCC patients demonstrated significant correlations between LCQ and VAS to the number of positive laryngeal sensations, number of cough triggers with LCQ and cough hypersensitivity as assessed by the cough hypersensitivity questionnaire [28].

The improvement of cough with BST further supports a role for “laryngeal hypersensitivity” in driving chronic cough. In our study, it was unclear whether laryngeal hypersensitivity was an independent phenomenon or related to GERD for which 75% of patients received anti-acid therapy. Therapy for GERD is recommended in chronic cough patients that are symptomatic for GERD [35] and despite elimination of the symptoms of GERD in a significant proportion of our patients, their cough tended to persist especially in those with features of cough hypersensitivity and laryngeal abnormalities. Even though this study did not purport to assess efficacy of a particular management approach to cough, we saw improvements in cough with BST, therapy for OSA with CPAP, or the combination of both for those with GERD, UACS and CVA. OSA, a condition characterized by recurrent trauma to the upper airway during sleep, is associated with both chronic cough [36] and laryngeal abnormalities [37, 38]. A recent longitudinal assessment showed preservation of sensory palatal thresholds after 5 years of CPAP therapy as compared to worsening in untreated snorers [39]. Further research is necessary to unravel the association between GERD and OSA and laryngeal hypersensitivity and the role of BST along other therapies typically recommended.

The 2006 ACCP cough guidelines paved the way for systematic analysis and treatment of chronic cough [2]. An anatomic diagnostic protocol as outlined in the cough guidelines identifies disease aetiologies for cough from almost all regions with cough afferents. However, disorders of the larynx were omitted from the list of aetiologies for chronic cough. Instead, laryngeal symptoms associated with chronic cough were attributed to GERD [40] and UACS [41] in chronic cough patients.

Limitations

Despite being a retrospective, single-centre study, this report offers important insights into the value of an interdisciplinary assessment for UCC/RCC that includes systematic laryngeal evaluation. Currently, there is no agreement on the protocol for laryngeal evaluation for chronic cough due to the lack of consensus regarding functional laryngeal anomalies and laryngeal hypersensitivity [42]. The operator dependency of results, sporadic nature of laryngeal complaints, lack of validated cutoffs for laryngeal closure and lack of understanding of how mechanical or other stimuli uncover laryngeal dysfunction can further confound the diagnosis of laryngeal hypersensitivity [42]. To address reliability issues inherent with clinical interpretation of laryngeal structure and function in nasoendoscopy recordings, an SLP independently documented impressions blind to the original finding for purposes of this study. However, future studies need to systematically evaluate nasoendoscopy findings across multiple clinic sites to further examine this study's finding of laryngeal functional abnormalities in UCC/RCC. In this study, validated questionnaires to assess complaints of hypertussia, allotussia, and abnormal laryngeal sensation were not used. The VHI and DI were used to assess voice and upper airway dyspnoea respectively although the validated laryngeal hypersensitivity questionnaire (LHQ) was not used to assess laryngeal hypersensitivity [43]. One of the goals of this study was to determine whether laryngeal dysfunction as assessed by SLPs during their routine clinical evaluation could elucidate the potential contribution of laryngeal hypersensitivity to chronic cough. This aspect is important given that other practitioners need to be able evaluate for laryngeal dysfunction based on symptom assessment (as is done for other etiologies of cough)

and referred for appropriate management of laryngeal dysfunction early on in disease course before prolonged and frequently ineffective empiric therapeutic trials are done.

Conclusion

This report highlights the prevalence of a constellation of symptoms typifying laryngeal dysfunction and cough hypersensitivity with a call to designate laryngeal hypersensitivity as a specific phenotype of cough. The larynx has been termed as the “pinnacle of cough” given the integration of motor processes at this location that are required for the execution of the cough reflex [41]. Although further work is required to validate the proposed criteria for laryngeal hypersensitivity, its high prevalence [44] and response to BST [13] makes it a necessary to evaluate for laryngeal hypersensitivity during the management of the patient with chronic cough. The use of targeted questions to identify both laryngeal dysfunction and cough hypersensitivity can delineate the syndrome of laryngeal hypersensitivity and facilitate development of therapies directed at the underlying neuropathic basis of cough.

Acknowledgments – Authors thank the speech language pathologists at the Voice Disorders Center (Tiffany Lyon SLP, Julia Ellerston SLP and Elizabeth Hary SLP) and Dr. Marshall Smith for the multidisciplinary evaluations of patients studied in this cough clinic. We also thank all the referring providers within Pulmonary, ENT, Internal Medicine, Family Practice, Allergy clinics within and outside the University of Utah Health Care system.

REFERENCES

1. Chung KF, Pavord ID. Chronic cough 1 Prevalence, pathogenesis and causes of chronic cough. *Lancet* 2008;371:1364-1374.
2. Pratter MR, Brightling CE, Boulet LP, Irwin RS (2006) An empiric integrative approach to the management of cough: ACCP evidence-based clinical practice guidelines. *Chest* 129 (1suppl):222S–231S.
3. Gibson PG, Vertigan AE. Management of chronic refractory cough. *BMJ* 2015;351:h5590 doi: 10.1136/bmj.h5590.
4. Morice AH. The cough hypersensitivity syndrome: a novel paradigm for understanding cough. *Lung* 2010;188 (suppl 1):S87-S90.
5. Morice AH, Millqvist E, Belvisi MG, Bieksiene K, Birring SS, Chung KF, Dal RWD Negro, Dicipinigaitis P, Kantar A, McGarvey LP, Pacheco A, Sakalauskas R, Smith JA. Expert opinion on the cough hypersensitivity syndrome in respiratory medicine. *Eur Respir J* 2014;44:1132.
6. Vertigan A, Theodoros D, Gibson P, Winkworth A. Voice and upper airway symptoms in people with chronic cough and paradoxical vocal fold movement. *J Voice* 2007;21:361-383.
7. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP, STROBE Initiative. Strengthening the reporting of observational studies in epidemiology (STROBE): guidelines for reporting of observational studies. *BMJ* 2007;335:806-808.
8. Birring SS., Prudon B, Carr AJ, Singh SJ, Morgan MDL, Pavord ID. Development of a symptom specific health status measure for patients with chronic cough: Leicester Cough Questionnaire (LCQ). *Thorax* 2003;58:339-343.

9. Jacobson BH, Johnson A, Grywalski C, Silbergleit A, Jacobson G, Benninger MS, Newman CW. The voice handicap index (VHI) development and validation. *American Journal of Speech-Language Pathology* 1997;6:66-70.
10. Gartner-Schmidt JL, Shembel AC, Zullo TG, Rosen CA. Development and validation of the Dyspnea Index (DI): a severity index for upper airway-related dyspnea. *J Voice* 2014;28:775-782.
11. Patel RR, Awan SN, Barkmeier-Kraemer J, Courey M, Deliyski D, Eadie T, Paul D, Svec JG, Hillman R. Recommended protocols for instrumental assessment of voice: American Speech-Language-Hearing Association expert panel to develop a protocol for instrumental assessment of vocal function. *American Journal of Speech- Language- Pathology* 2018;27:887-905.
12. McGarvey L, McKeagney P, Polley L, MacMahon J, Costello RW. Are there clinical features of a sensitized cough reflex? *Pulm Pharmacol Ther* 2011;2009;22:59-64.
13. Chamberlain Mitchell SA, Garrod R, Clark L, Douiri A, Parker SM, Ellis J, Fowler SJ, Ludlow S, Hull JH, Chung KF, Lee KK, Bellas H, Pandyan A, Birring SS. Physiotherapy, and speech language therapy intervention for patients with refractory chronic cough: a multicenter randomized control trial. *Thorax* 2017;72:129-136.
14. Epstein LJ, Kristo D, Strollo Jr PJ, Friedman N, Malhotra A, Patil SP, Ramar K, Rogers R, Schwab RJ, Weaver EM, Weinstein MD, Adult obstructive sleep apnea task force of the American Academy of Sleep Medicine. *J Clin Sleep Med* 2009;15:5:263-276.
15. Patton CM, Lim KG, Ramlow LW, White KM. Increasing efficiency in evaluation of chronic cough: a multidisciplinary, collaborative approach. *Q Manage Health Care* 2015;24:177-82.

16. Chung KF, Widdicombe JG. Cough: Setting the scene. In Handbook of Experimental Pharmacology. Pharmacology and Therapeutics of Cough. Eds. Chung and Widdicombe. Pages 1-21.
17. Birring SS. The search for hypersensitivity in chronic cough. Eur Respir J 2017;49:1700082 DOI: 10.1183/13993003.00082-2017
18. Song W-J, Morice AH. Cough hypersensitivity syndrome: a few more steps forward. Allergy Asthma Immunol Res 2017;9:394-402.
19. O'Connell F, Thomas VE, Pride NB, Fuller RW. Capsaicin cough sensitivity decreases with successful treatment of chronic cough. Am J Respir Crit Care Med 1994;150:374-80.
20. Mazzone SB, Chung KF, McGarvey L. The heterogeneity of chronic cough: a case for endotypes of cough hypersensitivity. Lancet Respir Med 2018;6:636-646.
21. Ryan NM, Gibson PG. Characterization of laryngeal dysfunction in chronic persistent cough. Laryngoscope 2009;119:640-45.
22. Vertigan AE, Gibson PG. Chronic refractory cough as a sensory neuropathy: evidence from a reinterpretation of cough triggers. J Voice 2011;5:596-601.
23. Bucca C, Bugiani M, Culla B, et al. Chronic cough and irritable larynx. J Allergy Clin Immunol 2011;127:412-420.
24. Ryan NM, Birring SS, Gibson PG. Gabapentin for refractory chronic cough: a randomized, double-blind, placebo-controlled trial. Lancet 2012;380:1583-89.
25. Vertigan AE, Bone SL, Gibson PG. Laryngeal sensory dysfunction in laryngeal hypersensitivity syndrome. Respirology 2013;18:943-56.

26. Hilton E, Marsden P, Thurston A, Kennedy S, Decalmer S, Smith JA. Clinical features of the urge-to-cough sensation in patients with chronic cough. *Respir Med* 2015;109:701-707.
27. Vertigan AE, Kapela SM, Kearney EK, Gibson PG. Laryngeal hypersensitivity in cough hypersensitivity syndrome: a cross-sectional observational study. *J Allergy Clin Immunol Pract* 2018;6:2087-95.
28. Won H-K, Kang S-Y, Kang Y, et al. Cough-related laryngeal sensations and triggers in adults with chronic cough: Symptom profile and impact. *Allergy Asthma Immunol Res* 2019;11:622-31.
29. Chung KF. Approach to chronic cough: the neuropathic basis for cough hypersensitivity syndrome *J Thoracic Dis* 2014;6(suppl 7):S699-S707.
30. Davenport PW. Urge-to-cough: What can it teach us about cough? *Lung* 2008;186 (suppl 1):S107-S111.
31. Mazzone SB, Undem BJ. Vagal afferent innervation of the airways in health and disease. *Physiol Rev* 2016;96:975-10.
32. Polverino M, Polverino F, Fasolino M, Ando F, Alfieri A, De Blasio F. Anatomy and neuro-pathophysiology of the cough reflex arc. *Multidiscip Respir Med* 2012;7:5
33. Ludlow C. Laryngeal reflexes: Physiology, technique and clinical use. *J Clin Neurophysiol* 2015;32:284-93
34. Irwin R.S., Ownbey R., Cagle P.T., Baker S., and Fraire A.E. Interpreting the histopathology of chronic cough: a prospective, controlled, comparative study. *Chest* 2006; 130: pp. 362-370

35. Kahrilas PJ, Altman KW, Chang AB, et al. Chronic cough due to gastroesophageal reflux in adults CHEST guideline and expert panel report. CHEST 2016;150:1341-1360.
36. Sundar KM, Willis AM, Smith S, Hu N, Kitt JP, Birring SS. A randomized, controlled, pilot study of CPAP for patients with chronic cough and obstructive sleep apnea. Lung 2020; 198:
37. Roy N, Merrill RM, Pierce J, Sundar KM. Evidence of possible irritable larynx syndrome in obstructive sleep apnea – an epidemiologic approach. J Voice 2020 online ahead of print March 11.
38. Nguyen AT, Jobin V, Payne R, Beauregard J, Naor N, Kimoff RJ. Laryngeal and velopharyngeal sensory impairment in obstructive sleep apnea. Sleep 2019;28:575-593.
39. Svanborg E, Ulander M, Broström A, Sunnergren O. Palatal sensory function worsens in untreated snorers but not in CPAP-treated patients with sleep apnea, indicating vibration-induced nervous lesions. CHEST 2020;157:1296-1303.
40. Phua SY, McGarvey LP, Ngu MC, Ing AJ. Patients with gastroesophageal reflux disease and cough have impaired laryngopharyngeal mechanosensitivity. Thorax. 2005;60:488---91.
41. Altman KW, Lane AP, Irwin RS. Otolaryngology aspects of chronic cough. J Allergy Clin Immunol 2019;7(6):1750-55.
42. Famokunwa B, Walsted ES, Hull JH. Assessing laryngeal function and hypersensitivity. Pulm Pharmacol Ther 2019;56:108-115.
43. Vertigan AE, Bone SL, Gibson PG. Development and validation of the Newcastle laryngeal hypersensitivity questionnaire. Cough 2014;10:1.

44. Roy N, Stemple J, Merrill RM, Thomas L. Epidemiology of voice disorders in the elderly: preliminary findings. *Laryngoscope* 2007;117:628-633.

	Overall	Age-distribution			Sex-distribution	
	n=60	18-40 years n=9	41-70 years n= 33	>71 years n= 18	Male n=16	Female n=44
Age [mean]	59.5±SD 16.4	30.5 3± SD 7.9	58.50 ± SD 8.8	76.08 ± SD 4.3	55.06 ± SD 16.2	61.2 2± SD 16.3
Sex	M=16 F=44	M=2 F=7	M=10 F=23	M=4 F=14		
BMI	29.47 ± SD 6.4	28.87 ± SD 7.13	30.11 ± SD 6.7	28.65 ± SD 5.8	30.49 ± SD 3.8	29.07 ± SD 7.2
Race/Ethnicity						
- White	52	5	29	18	15	37
- Hispanic	2	1	1	0	0	2
- Asian	2	1	1	0	1	1
- Pacific	1	1	0	0	0	1
- Islander	1	0	1	0	0	1
- African	4	2	2	0	0	4
- American						
- Other						
Comorbidities						
- GERD	9/60	0	7	2	2	7
- UACS	2	0	2	0	1	2
- CVA	1	0	0	1	0	0
- GERD and	2	0	0	2	2	0
- UACS	46	9/9	24	13	11	35
- None						

Table 1: Demographics of patients and cormorbidities diagnosed. Abbreviations; N – Number; SD – Standard deviation; GERD – Gastroesophageal reflux disease; UACS – Upper airway cough syndrome; CVA – Cough variant asthma.

	Overall	Age-distribution			Sex-distribution	
	Overall N=60	18-40 years [n=9]	41-70 years [n=33]	>71 years [n= 18]	Male [n=16]	Female [n=44]
Duration [in months]	75.62± SD 105	39± SD 75.8	78.06±SD 106.6	89.44± SD 117.6	87.06± SD 105.7	71.45± SD 106.8
LCQ	9.91 ±2.75 n=56	9.56 ±SD 2.6 n=9	10.08 ±SD 2.699 n=31	9.77 ±SD 3.027 n=16	8.89 ±SD 1.89 n=16	10.25±SD 2.9 n=42
VAS	63.79 ±SD.47 n=56	70.25 ±SD 17.49 n=8	64.16 ±SD 27.355 n=31	63.79 ±SD 17.49 n=56	64.44 ±SD 28.844 n=16	63.52 ±SD 25.86 n=40
Paroxysms of coughing	44/60	7/9	27/33	10/18	11/16	33/44
- Present	8	2	3	3	1	7
- Absent	8	0	3	5	4	4
- Not reported						

Table 2: Cough Characteristics of patients. Abbreviations: N – number; SD – Standard deviation; LCQ – Leicester Cough Questionnaire; VAS – Visual analog scale.

	Overall	Age-distribution			Sex-distribution	
	Overall n=60	18-40 years n=9	41-70 years n= 33	>70 years n=18	Male n=16	Female n=44
<u>Laryngeal</u>						
<u>paresthesia</u>	13/60 [22%]	3	6	4	4	9
Globus	24 [40%]	5	15	4	8	16
Tickle	14 [23%]	3	6	5	4	10
Irritation	8 [13%]	1	6	1	2	6
Dry Throat	4 [6%]	0	3	1	2	2
Sore Throat	4 [6%]	0	1	3	2	2
Mucus						
Throat clearing						
- Present	27 [45%]	3	16	8	8	19
- Absent	6 [10%]	1	3	2	1	5
- Not specified	27 [45%]	5	14	8	7	20
Voice Change	32/60 [53%]	4	16	12	7	25
Voice Handicap Index	21.50 ±SD 21.9 n=58	24 ±SD 6.3 n= 9	20.69 ±SD 8 n=31	21.71 ±SD 7.3 n=16	19.07 ±SD 8.3 n=14	22.27 ±SD 7.9 n=44
Upper airway dyspnoea	13/60 [22%]	3	9	1	2	11
Dyspnea index	14.12 ±SD 8.04 n=56	20.56 ±SD 18.13 n=9	13.71±SD 8.0 n=31	11.31 ±SD 7.3 n=17	15.57 ±SD 8.3 n=14	13.64 ±SD 7.9 n=42

Table 3: Laryngeal abnormalities in patients with chronic cough.

Study	Subjects evaluated and number studied	Abnormal cough triggers In RCC/UCC	Laryngeal paresthesia	Throat clearing in UCC/RCC	Voice abnormalities and upper airway dyspnoea in UCC/RCC	Laryngeal abnormalities on endoscopy	Other findings
Vertigan (2007) (ref 6)	111 chronic cough patients (55/111 with PVFM also)	Average of 8 triggers noted for each participant	41% reported warning sensations with most identifying throat as the site of trigger	Noted but percentage of patients with this symptom not specified	Abnormal voice and dyspnoea scores in chronic cough and cough plus PVFM patients	55 chronic cough with PVFM included in analysis	Talking, cigarette smoke, cold air and irritability in throat identified as most frequent triggers.
Ryan (2009) (ref 22)	24 subjects with chronic persistent cough	Not reported	Not reported	Not reported	Not reported	14/24 noted with PVFM	Abnormal extrathoracic airway hyper-reponsiveness using hypertonic saline in cough + PVFM group
Vertigan (2011) (ref 23)	53 RCC patients	100% with 1 or more cough triggers	Seen in 94%	Not reported	Not reported	Shortness of breath reported as a cough trigger	Categorized tussive and non-tussive triggers to capture hypertussia and allotussia
Bucca (2011) (ref 24)	61 UCC	Not reported	Not reported	Not reported	Not reported	66% of UCC patients had abnormal LHR	LHR seen not only in UCC but also in GERD (62) asthma (41), rhinosinusitis (208)
Ryan (2012) (ref 25)	62 patients	50% with laryngeal hypersensitivity	63% with central reflex sensitisation	Not reported	76% with voice changes	Not reported	This was a placebo-controlled study of gabapentin in chronic cough that showed higher cough improvement in those with central sensitization.
Vertigan (2013) (ref 26)	33 RCC	Not reported	RCC patients had worse laryngeal paresthesia scores	Not reported	Worse VHI scores and DSI as compared to controls	Not reported	57% of RCC had abnormal fall in FIF50 following hypertonic

			compared to controls				saline and worse compared to control groups
Hilton (2015) (ref 27)	100 consecutive chronic cough patients	79% with hypertussia 72% with allotussia	75% with neck sensations	~50% with throat clearing	Not reported	Not reported	Correlation between urge-to-cough sensation intensity and total LCQ score.
Vertigan (2018) (ref 28)	20 patients with RCC	Not reported	Abnormal LHQ with correlation with cough frequency	Not reported	Abnormal mean VHI values in RCC group compared to controls	Laryngeal motor dysfunction with breathing in 47% RCC (67% with odor challenge). Mediolateral constriction of false vocal cords in 45% with incomplete or abnormal vocal cord closure in 75% of RCC group.	Specialized voice testing and timed swallow test different in RCC group as compared to controls. 69% with of 27/69 patients that had abnormal cough had PVFM
Won (2020) (ref 29)	62 patients with RCC	100% of patients with 2 or more cough triggers	100% of patients with at least one abnormal sensation	Not reported	Not reported	Not assessed	Correlations between LCQ and number of laryngeal sensations, number of cough triggers, CHQ score
Sundar (current study)	60 patients RCC/UCC	100% with 1 or more cough triggers	Seen in 97%	Seen in 45% of patients	VHI abnormal in 43% and DI in 75% of patients	Functional laryngeal abnormalities in 88% patients	Correlations between LCQ and VHI and DI noted

Table 4. Studies of UCC/RCC with abnormalities of cough hypersensitivity and laryngeal findings. Abbreviations: ref – reference; RCC – Chronic refractory cough; UCC – Unexplained chronic cough; PVFM – Paradoxical vocal fold motion; LHR – Laryngeal hyper-responsiveness; LHQ – Laryngeal hypersensitivity questionnaire; LCQ – Leicester Cough Questionnaire; CHQ – Cough Hypersensitivity Questionnaire; DSI – Dyspnea Symptom Index; VHI – Voice Handicap Index; DI – Dyspnea Index.






Figure 1. Flow diagram of patients screened and enrolled into observational cohort of patients with chronic refractory cough (CRC) and unexplained chronic cough (UCC). Abbreviations – COPD – Chronic obstructive pulmonary disease; GERD – Gastroesophageal reflux disease; UACS – Upper airway cough syndrome; CVA – Cough variant asthma; BST – Behavioral speech therapy; CPAP – Continuous positive airway pressure.



2a.



2b.

Fig 2: Frequency of cough- related triggers, overall (2a) and based upon different age groups (2b).

Nasoendoscopy with Stroboscopy Findings

