



Early View

Review

ERS International Congress 2021 virtual: highlights from the Paediatric Assembly

Cristina Ardura-Garcia, Alicia Abellan, Sara Cuevas-Ocaña, Nadine Freitag, Yin Ting Lam, Heidi Makrinioti, Monique Slaats, Matteo Storti, Emma E Williams, Theodore Dassios, Liesbeth Duijts, Refika H Ersu, Stojka Fustik, Rory E. Morty, Marijke Proesmans, Dirk Schramm, Sejal Saglani, Alexander , Marielle W. Moeller and Pijnenburg

Please cite this article as: Ardura-Garcia C, Abellan A, Cuevas-Ocaña S, *et al.* ERS International Congress 2021 virtual: highlights from the Paediatric Assembly. *ERJ Open Res* 2022; in press (<https://doi.org/10.1183/23120541.00643-2021>).

This manuscript has recently been accepted for publication in the *ERJ Open Research*. It is published here in its accepted form prior to copyediting and typesetting by our production team. After these production processes are complete and the authors have approved the resulting proofs, the article will move to the latest issue of the ERJOR online.

Copyright ©The authors 2022. This version is distributed under the terms of the Creative Commons Attribution Non-Commercial Licence 4.0. For commercial reproduction rights and permissions contact permissions@ersnet.org

ERS International Congress 2021 virtual: highlights from the Paediatric Assembly

Cristina Ardura-Garcia¹, Alicia Abellan²⁻⁵, Sara Cuevas-Ocaña⁶, Nadine Freitag⁷, Yin Ting Lam¹, Heidi Makrinioti⁸, Monique Slaats⁹, Matteo Storti¹⁰, Emma E Williams¹¹, Theodore Dassios^{11,12}, Liesbeth Duijts^{9,13}, Refika H Ersu¹⁴, Stojka Fustik¹⁵, Rory E. Morty^{16,17}, Marijke Proesmans¹⁸, Dirk Schramm⁷, Sejal Saglani¹⁹, Alexander Moeller²⁰ and Marielle W. Pijnenburg⁹

Affiliations:

1. Institute of Social and Preventive Medicine, University of Bern, Bern, Switzerland
2. ISGlobal, Barcelona, Spain.
3. Universitat Pompeu Fabra (UPF), Barcelona, Spain.
4. Fundació Institut Universitari per a la recerca a l'Atenció Primària de Salut Jordi Gol i Gurina (IDIAPJGol), Barcelona, Spain.
5. Spanish Consortium for Research on Epidemiology and Public Health (CIBERESP), Madrid, Spain.
6. Nottingham Biodiscovery Institute, School of Medicine, University of Nottingham, University Park, Nottingham, UK.
7. Department of General Pediatrics, Neonatology and Pediatric Cardiology, Medical Faculty, University Hospital Düsseldorf, Heinrich-Heine-University Düsseldorf, Düsseldorf, Germany.
8. West Middlesex University Hospital, Chelsea, and Westminster Foundation Trust, UK.
9. Department of Paediatrics, Division of Paediatric Respiratory Medicine and Allergology, Erasmus MC – Sophia Children's Hospital, University Medical Centre, Rotterdam, The Netherlands
10. Department of Chemical & Life Sciences and Environmental Sustainability, University of Parma, Parma, Italy
11. Department of Women and Children's Health, School of Life Course Sciences, Faculty of Life Sciences and Medicine, King's College London, London, United Kingdom
12. Neonatal Intensive Care Centre, King's College Hospital NHS Foundation Trust, London, United Kingdom
13. Division of Neonatology, Department of Pediatrics, Erasmus MC, University Medical Center Rotterdam, Rotterdam, the Netherlands.

14. Division of Respiriology, University of Ottawa Children's Hospital of Eastern Ontario, Ottawa, Canada.
15. University Children's Hospital, Skopje, North Macedonia.
16. Department of Lung Development and Remodelling, Max Planck Institute for Heart and Lung Research, Bad Nauheim, Germany
17. Department of Translational Pulmonology and the Translational Lung Research Center Heidelberg, University Hospital Heidelberg, member of the German Center for Lung Research (DZL), Heidelberg, Germany
18. Division Woman and Child, Department of Paediatrics, University Hospitals Leuven, Leuven, Belgium
19. National Heart & Lung Institute, Imperial College London, London, UK
20. Division of Paediatric Pulmonology, University Children's Hospital Zurich, University of Zurich, Zurich, Switzerland

Correspondence: Cristina Ardura-Garcia. Institute of Social and Preventive Medicine, University of Bern, Mittelstrasse 43, 3012 Bern, Switzerland. E-mail: cristina.ardura@ispm.unibe.ch

Take home message: Highlights from the Paediatrics Assembly at the #ERSCongress 2021

Manuscript word count: 5130 words

Abstract

In this review, Early Career Members of the European Respiratory Society (ERS) and the Chairs of the Assembly 7 – Paediatrics Groups, present the highlights in paediatric respiratory medicine from the ERS International Congress 2021. The Groups cover respiratory physiology and sleep, asthma and allergy, cystic fibrosis (CF), respiratory infection and immunology, neonatology and intensive care, epidemiology, bronchology, and lung and airway development. We here describe new developments in lung function testing and sleep disordered breathing diagnosis, early life exposures affecting pulmonary function in children and effect of COVID-19 on sleep and lung function. In paediatric asthma, we present the important role of the exposome in asthma development, and how biologics can provide better outcomes. We discuss new methods to assess distal airways in children with CF, as some details remain blind when using Lung Clearance Index. Moreover, we summarise the new ERS guidelines for bronchiectasis management in children and adolescents. We present interventions to reduce morbidity and monitor pulmonary function in newborns at risk of bronchopulmonary dysplasia and long term chronic respiratory morbidity of this disease. In respiratory epidemiology, we characterise Primary Ciliary Dyskinesia, identify early life determinants of respiratory health and describe the effect of COVID-19 preventive measures on respiratory symptoms. Also, we describe the epidemiology of interstitial lung diseases, possible consequences of tracheomalacia and a classification of diffuse alveolar haemorrhage in children. Finally, we highlight that the characterisation of genes and pathways involved in the development of a disease is essential to identify new biomarkers and therapeutic targets.

250 words

The Paediatric Assembly (Assembly 7) reviewed recent research advances in the paediatric respiratory field presented by leading experts and early career members at the European Respiratory Society (ERS) virtual International Congress 2021. Assembly 7 members presented 249 abstracts in oral and e-poster sessions and organised several symposia, including two clinical guidelines sessions. The Paediatrics Assembly content is very diverse as it covers the whole field of paediatric respiratory diseases: from very rare diseases such as childhood interstitial disease, to extremely common diseases such as asthma. Additionally, many diseases in adulthood have their origin in childhood, and environmental factors such as air pollution and climate change have a disproportionate effect on children. We are proud that the paediatric assembly was able to compose a high-quality program for paediatricians and other health care professionals.

We here review the paediatric highlights from the 2021 virtual congress, summarised by Assembly 7 Early Career Members and Chairs of the 8 Groups that form the Assembly. For each Group, the Chair selected the most relevant session. The sessions are presented in the order of the Group's numbers.

Group 7.1: Paediatric respiratory physiology and sleep

Several themes have emerged from this year's session on paediatric physiology and sleep research, including new developments in lung function testing and sleep disordered breathing diagnosis, and the effect of early life exposures and COVID-19 on lung function.

Earlier studies showed that prematurity, low birthweight and maternal abuse of nicotine have an adverse effect on children's lung function[1]. Salem Mahmoud Y. *et al.* examined the association of low birthweight and impaired lung function in monochorionic- diamniotic twins with selective fetal growth restriction (FGR). Nineteen twin pairs (age between 8-29 years) were included with mean birthweight difference of 29% and a mean gestational age of 33.5 weeks. Mean z-score for FEV1 was 0.61 and for FVC 0.56, and both were lower in twins with FGR compared to those without[2].

Gudmundsdóttir *et al.* investigated the influence of mid-pregnancy physical activity (PA) on lung function of 3-month-olds. Tidal flow volume measurements of 814 infants were compared with self-reported PA. The ratio time to Peak Tidal Expiratory Flow to Expiratory time (tPTEF/tE) as a continuum was not significantly associated with maternal PA. However, infants of inactive mothers compared with active mothers had higher odds of having a lower lung function by tPTEF/tE <0.25[3].

Färdig *et al.* evaluated the tidal flow-volume (TFV) loops of 93 three-month-olds to investigate differences in awake and sleeping state. A higher ratio of tPTEF/tE, the corresponding volume ratio and respiratory rate, but similar tidal volume were observed during awake compared to the sleeping state suggesting separate normative values may be required[4].

Fouzas *et al.* examined the characteristics of FEV1 fluctuations in children with controlled asthma who were treated with inhaled corticosteroids (ICS). They compared FEV1 recordings (for 3 months) of 31 participants with controlled asthma with recordings of 68 healthy controls. Children with controlled asthma had normal FEV1 values with correlated long-range fluctuations unrelated to age[5]. A retrospective study determined the relationship between forced expiratory time (FET) and airway obstruction in 2623 children. FET was highly related to central airway obstruction (FEV1/FVC)[6].

Impulse oscillometry (IOS), a non-invasive modality of testing lung function, measures airway impedance (resistance and reactance). However, no standards have been set for IOS[7]. Park *et al.* compared IOS to conventional pulmonary function test (PFT) in 75 patients with neuromuscular disorders with a mean age of 20 years. PFT showed restrictive lung function in all participants. Lung volumes by PFT and resistance by IOS decreased by age. Higher lung volumes were correlated with higher resistances[8].

Two studies investigated the effect of COVID-19 and lockdown in children. Schlegtendal *et al.* compared lung functions of 73 children after an average of 2.6 months following COVID-19 infection to 45 controls. Respiratory symptoms following COVID-19 were reported in 11%. There were no significant differences in the lung function. Only two patients with persistent respiratory symptoms showed abnormal lung function[9]. Another study conducted a global survey to assess the sleep habits of children before and during the COVID-19 lockdown and analysed 845 questionnaires (ages 3-17 years). Lockdown was associated with later bedtime and wake time in children but this shift did not change the sleep duration in half of the children, with the 14-17 year-olds group having the lowest rate of no change (28%)[10].

Nocturnal pulse oximetry (NOx) can be used to screen for obstructive sleep apnoea (OSA)[11]. Fouzas *et al.* analysed advanced signals and indices that may accurately diagnose OSA for 45 children (aged 2-10 years). They suggested that cumulative desaturation area, variability of SpO₂, and heart rate could improve the diagnostic ability of NOx[12].

In conclusion, mid-pregnancy physical inactivity and low birthweight may be associated with impaired lung function in children. Impulse oscillometry may be a useful technique to evaluate lung function in children with neuromuscular disease. COVID-19 seems to have no significant impact in lung function nor lockdown on sleep duration in children. Nocturnal pulse oximetry could be improved with the use of cumulative desaturation area, variability of SpO₂, and heart rate to diagnose OSA.

Group 7.2: Paediatric asthma and allergy

The session on "Advances in childhood asthma: biologics, biomarkers and infections" included eight oral presentations and focussed on recent advances in the use of biologics and biomarkers in childhood asthma research.

The session started with two presentations around asthma in adolescents. Oyenuga *et al.* presented data about asthma diagnosis and asthma control in children aged 12-14 years old in Sub-Saharan Africa[13]. Indeed, there is only scarce available data around asthma control and impact in quality of life for Sub-Saharan adolescents [14]. Interestingly, they showed that 62% of the adolescent participants had severe symptoms and 81% of them had no doctor-confirmed diagnosis of asthma, while 37% of them had objective markers of asthma. A predictive approach was attempted by Amaral *et al.* who performed latent class analysis on clinical data from 162 adolescents with asthma. The aim was to identify distinct clusters of disease progression. What they showed was that the cluster of adolescents who presented the higher number of exacerbations and associated hospital visits were the ones who had poorly controlled asthma at recruitment stage[15].

The session continued with data around use of biologics in children with asthma. Ribas *et al.* presented data from the Phase III Voyage study showing that the daily use of dupilumab [16], compared to placebo, was associated with a reduced number of asthma exacerbations and improved lung function, but not with improved asthma control, in children aged 6-11 years with allergic asthma ($p < 0.05$ and $p < 0.001$)[17]. Carraro *et al.* focused on identifying urinary markers of response to omalizumab treatment and showed that histidine, the precursor of histamine, emerged as a sensitive marker that helped to identify response to omalizumab[18].

Beaufils *et al.* received the award for the best abstract of the Paediatric Assembly. This study aimed to identify clusters of preschool children with recurrent wheeze who are at higher risk to develop asthma based on bronchial remodelling features [19]. They performed fiberoptic bronchoscopy in 56 preschool children with persistent wheeze and identified histological parameters that were stronger predictors of asthma development. These parameters included the reticular basement membrane (RBM) thickness and epithelial cell shedding[20]. Validation of these predictors in a larger sample is anticipated.

The last part of the session focussed on data around asthma prevention. Hose *et al.* applied latent class analysis to systematically assess feeding patterns and to relate these to asthma risk at school age. This data-driven approach identified a pattern of increased meat consumption conferring a risk for asthma diagnosis at 6 years. The possible underlying mechanism could involve gut microbial imbalance triggering immune dysfunction but is yet to be determined [21,22]. Da Silva *et al.* aimed

to investigate the associations between rhinovirus (RV)-induced bronchiolitis in infancy and lung inhomogeneities at preschool age using multiple breath nitrogen washout (MBW). Their cohort of 103 infants showed that hospitalisation with RV-positive bronchiolitis was associated with the highest risk for abnormal Lung Clearance Index (LCI) at preschool age[23]. Following on this association, Makrinioti *et al.* presented data from a meta-analysis demonstrating that the associations between RV-bronchiolitis and preschool wheeze and asthma development are stronger than those for RSV-induced bronchiolitis[24]. Future studies with further data around allergic sensitization could help identify whether RV-bronchiolitis infants may be a specific group for intervention studies for asthma prevention.

In summary, data presented showed that the exposome (viruses, diet) play an important role in asthma development and biologics can provide better outcomes in childhood asthma management. Assessment of control is important, as poor asthma control is associated with higher number of exacerbations.

Group 7.3.: Monitoring of lung disease and CFTR function in paediatric cystic fibrosis

This session focused on current and new methods to assess cystic fibrosis transmembrane conductance regulator (CFTR) function and lung disease in children with cystic fibrosis (CF).

Sandvik *et al.* examined MBW outcomes using EXHALYZER-D®-SPIROWARE® recordings of resident nitrogen (N₂) or sulphur hexafluoride (SF₆) in a cohort of 45 infants and toddlers with CF and 57 healthy controls. The use of SF₆MBW, regarded as gold standard for MBW[25], was recommended, as it provided less pronounced inter-test and between occasion differences for LCI than N₂MBW[26,27]. Dumas *et al.* used a cohort of 17 preschool children with CF[28] and 13 healthy controls to analyse Scond and Sacin indices of MBW[29] demonstrating 79% met acceptability criteria, similar to previously reported, in preschool children[30].

Longitudinal SF₆MBW assessments by Sandvik *et al.* in 2–45 month-olds with CF and healthy controls, detected a higher LCI in the CF than the control group, with 64% of children with CF showing intermittent lung function impairment and a LCI increase of 0.22 units/year associated with an overall worsening[31]. This highlights the importance of early therapeutic intervention and regular SF₆MBW assessments during the early stages of CF. Oestreich *et al.* also examined the evolution of LCI in healthy and CF cohorts by measuring SF₆MBW[32] at 6 weeks and 1 year of age. Despite similar lung function at 6 weeks of age, LCI significantly deteriorated over the first year of life in infants with CF[33]. The association found between LCI at 6 weeks and LCI at 1 year highlights the importance of targeting prevention strategies for patients at risk from their first months of life.

Perrem *et al.* presented another LCI-focused prospective observational study in a cohort of 98 school-age children with CF and 48 healthy controls[34] over 2 years. 19% and 37% of the CF visits showed an LCI change relative to the last visit in the range of $\pm 15\%$ and $\pm 10\%$ threshold, respectively. Changes associated with additional clinical factors, e.g. FEV₁, were much higher in the CF group, 78% at $\pm 15\%$ and 72% at $\pm 10\%$ threshold[35]. Thus, a change in LCI in the range of 10%–15% can be considered clinically meaningful and could be used to guide clinical decision-making. Frauchiger *et al.* also longitudinally assessed LCI in 74 children with CF (4–18 years old) and 19 healthy controls over a 7-year period, reporting 9% coefficient variation at population level, but 15% relative change in healthy and 23% in CF at individual level[36]. Thus, applying individual limits might be an appropriate approach to define minimal clinically important difference for routine clinical surveillance.

Willers *et al.* evaluated lung functionality, using label-free functional Matrix-Pencil Magnetic Resonance Imaging (MP-MRI) that allows for simultaneous ventilation and perfusion measurements, in 37 children with CF (4–17 years old) over 1 year. Although changes in LCI moderately correlated with changes in ventilation and perfusion defect, they did not correlate with change in structural pathology, highlighting the potential of MP-MRI to assess lung areas that are blind to LCI measurements[37]. Interestingly, Lv *et al.* presented the Thirona-developed Automated AA-LungQ method to analyse changes in small airways in a completely automated way. AA-LungQ method showed high sensitivity in detection of progressive widening and thickening of small airways in 111 chest-CT scans of 57 children with CF at 2-year intervals[38]. Finally, Grosse-Onnebrink *et al.* proposed using air-liquid interface cultures of respiratory epithelia derived from patient nasal brushings as *in vitro* model to assess personalised treatment responses based on transepithelial conductance measurements by the Ussing chamber system[39].

Overall, we need to define clinically relevant thresholds for LCI changes. Despite the high sensitivity of LCI, some details remain LCI blind, especially in more distal airways. Further studies on Scnd and Sacin indices might help uncover silent disease by LCI, but currently CT scans are highly recommended to gain information on distal lung structures and any structural changes.

Group 7.4: ERS Clinical practice guidelines for managing children and adolescents with bronchiectasis

Early diagnosis of bronchiectasis is crucial not just for therapeutic management but also for the family finally receiving this diagnosis after years of uncertainty. The task force for the management of paediatric bronchiectasis aimed to identify evidence-based management strategies for

investigation and treatment. This session summarises the main findings and recommendations of the task force's resulting report[40].

Zena Powell shared the story of her 14-year-old son, who was described as a "wheezy baby" with years of recurring exacerbation cycles of the lower airways before he was diagnosed with bronchiectasis at the age of eight. In 2018, she joined the task force to form the management guidelines that she describes as a starting point, not just the end. Parents can take these guidelines to their consultant, who receives a basis to start the parents on the journey of bronchiectasis management[40].

Andrew Bush presented the overall clinical approach. Protracted bacterial bronchitis (PBB) leads to reversible followed by irreversible airway dilatations. The diagnosis and adequate treatment of PBB are crucial before irreversible airway dilatations develop. For an improved bronchiectasis diagnosis, the guidelines recommend performing a chest CT scan and apply ≥ 0.8 broncho-arterial ratio instead of the 1.0 cut-off for adults. Investigations such as sweat test and immune function testing are necessary in every child with bronchiectasis to diagnose possible underlying causes. Exacerbations in bronchiectasis belong to the top three factors affecting a child's quality of life (QoL) according to a European Lung Foundation survey[41].

For treatment options, the guidelines recommend regular airway clearance techniques (ACT), as an individualised therapy performed by a paediatric trained respiratory physiotherapist, although no data are available on optimal ACT nor frequency[42]. Keith Grimwood further described the need to identify those who would benefit from hyperosmolar therapies while rhDNAse is advised against. Antibiotics are considered the standard of care for treating exacerbations, although there has only been one randomised controlled trial (RCT)[43]. The choice and route of antibiotics are based on airway cultures, history of hypersensitivity reactions to the antibiotic and the severity of the exacerbation. Recommendations for long-term macrolides to reduce exacerbations are based on three RCTs[44]. Findings of halving exacerbation frequency are consistent with RCTs in adults and children with primary ciliary dyskinesia. The guidelines formulate a conditional recommendation to eradicate *Pseudomonas aeruginosa* upon detection, based on the association between *P aeruginosa* and deteriorating clinical status. Since no data are available, no preferred regimen is formulated. There is only low-quality evidence and lack of efficacy for the use of asthma-based medications; they should not be used unless clear signs of underlying asthma. The guidelines recommend surgical resection of the affected lung lobe only if maximal therapies have failed and QoL is substantially impaired. Lobectomy should only be conducted by an expert surgeon at a specialised centre after assessment by a skilled interdisciplinary team[40].

Finally, Angela Zacharasiewicz described the practical approach with case presentations. She highlighted the need to ask important questions about cough quality, clinical signs and other symptoms, because they might uncover red flags. These include timing, tendency, triggers and sound. Other symptoms to inquire are failure to thrive, weight loss, fever, night sweat, hypoxia, chest pain, dyspnoea at rest or with exercise or tachypnoea. On clinical exam signs of chronic illness, neurodevelopmental abnormalities and pathological auscultation should be noted.

This session on the management guidelines for bronchiectasis in children and adolescents highlighted the involvement of parents and caretakers early on, from listening carefully to their symptom description to practical implementation of the guidelines they helped to develop. With the treatment plan, parents should receive an explanation of the disease pattern in lay terms.

Group 7.5: Respiratory disorders in neonatal and paediatric intensive care

This session included presentations on interventions to reduce morbidity and monitor pulmonary function in newborn infants who are at risk of developing bronchopulmonary dysplasia (BPD) and long term chronic respiratory morbidity in childhood and beyond [45].

Volumetric capnography is a novel non-invasive technique which can inform on ventilation inhomogeneity, though technically difficult in preterm infants due to anatomical characteristics and technical limitations [46]. Williams *et al.* described this technique to be feasible during the resuscitation of mechanically ventilated prematurely born infants. Yet immediately following surfactant administration ventilation inhomogeneity increased ($25.9 \text{ mmHg.ml}^{-1}$ versus $17.2 \text{ mmHg.ml}^{-1}$), likely due to partial small airway obstruction caused by the presence of liquid surfactant in the small airways [47]. Novel techniques of surfactant delivery in preterm infants are being developed. Use of high-flow nasal cannula (HFNC) oxygen during less invasive surfactant administration to preterm infants is one such method. This practice was deemed effective in reducing the rates of mechanical ventilation, with 56.6% of infants not requiring the use of premedication[48].

Upper airway compliance was described by Bizzotto *et al.* to contribute to carbon dioxide removal during non-invasive high frequency oscillatory ventilation (nHFOV) and be maximal at 10Hz. This dependence upon oscillatory frequency suggests tailoring of nHFOV could help optimise effectiveness of nHFOV [49]. Delivery of effective fraction of inspired oxygen (FiO_2) is also important for oxygen dependent infants supported by HFNC. Fouzas *et al.* showed that when minute ventilation is high at 1200ml/min (compared to 400ml/min) and the flow rate set at less than 5L/min, the FiO_2 that is effectively delivered diverges from the target oxygen level that is set [50].

Such observations may be useful when considering weaning of HFNC in those preterm infants who remain oxygen dependent.

In prematurely born infants who require initial invasive respiratory support, accurate prediction of extubation success is a challenge. Surface diaphragmatic electromyography is a novel technique to measure electrical activity of the diaphragm. Arattu-Thodika *et al.* utilised this method during a spontaneous breathing trial (SBT) in mechanically ventilated infants and found that an increase in electrical activity of the diaphragm during the SBT was 71.4% sensitive and 85.7% specific in predicting extubation failure [51]. Neurally adjusted ventilatory assist (NAVA) also monitors diaphragmatic activity and can be used as both an invasive and a non-invasive mode of ventilation for preterm infants with evolving/established BPD [52]. Shetty *et al.* described how NAVA effectively reduced the duration of respiratory support and the length of hospital stay when compared to conventional modes of ventilation (112 versus 140 days).

Lung function trajectories, as measured by respiratory reactance (Xrs) in preterm infants born earlier than 32 weeks gestation, may also aid in guiding ventilatory support with positive end-expiratory pressure (PEEP). Infants with BPD supported with high PEEP at one week of life had reduced Xrs suggestive of pulmonary overdistension. At 36 weeks postmenstrual age however, an increase in Xrs in response to high PEEP was thought to be suggestive of peripheral small airway collapse [53]. Airway function post puberty was reported to be better in infants born earlier than 29 weeks of gestation to multiple pregnancies rather than singleton pregnancies (FEV1 -1.11 versus -0.50) [54]. Moreover, Luchnikova *et al.* described how infants born to mothers with severe bronchial asthma who required systemic glucocorticoids during pregnancy were more likely to be exposed to intrauterine infection and to be born prematurely (38.5% versus 11.7%) [55]. When considering exposure to antenatal corticosteroids and the effect by gender, female infants showed a greater beneficial response when considering the outcome of death before discharge compared to male infants (OR 1.81 versus OR 1.36) [56].

The novel techniques and respiratory modalities presented hold promise for improvement in the care of prematurely born infants. Greater understanding of the pathophysiology underlying specific diseases, combined with newer modes of non-invasive ventilatory support, may allow for reducing long-term pulmonary morbidity.

Group 7.6: Paediatric respiratory epidemiology

Current research highlighted the importance of characterisation, causes and consequences of various paediatric respiratory diseases. Goutaki *et al.* characterised the prevalence and severity of respiratory

symptoms among patients in the Swiss primary ciliary dyskinesia (PCD) Registry through a FOLLOW-PCD [57] based questionnaire, and concluded that repeated standardised clinical data will allow improved characterisation of PCD phenotypes, disease course and prognosis[58]. Rumman *et al.* showed how collaborations can facilitate PCD diagnosis in resource-limited settings among children participating in a large Palestinian PCD cohort, and identified nasal nitric oxide and targeted genetic testing to be useful diagnostic tools in local and resource-limited settings[59].

Early life is a critical period of airway and lung development and especially relevant in terms of long-term effects on respiratory health. Decrue *et al.* observed among 254 preterm and 517 term infants of the Basel-Bern Infant Lung Development cohort[60] that higher particulate matter of less than 10 μm during the second trimester of pregnancy was associated with lower lung function and higher oxidative stress shortly after birth. Participants exposed to low-to-moderate air pollution levels and the effects observed were stronger among preterm infants[61]. Abellan *et al.* applied unique exposome approaches among 5624 mother-child pairs from the Generation R birth cohort[62]. They showed that living during pregnancy in more polluted, noisier, with less green and blue spaces, and more urbanised neighbourhoods may contribute to lower lung function at 10 years of age and increased odds of wheezing during childhood[63]. Talaei *et al.* used principal component analysis on food frequency questionnaire data, and identified processed, traditional, and health-conscious dietary patterns at age 7 years. The health-conscious pattern was associated with higher lung function, and the processed pattern was with lower lung function at 15 years of age[64]. Koch *et al.* [65] identified four lung function growth trajectories (low, average, high, and catch-up) using repeated lung function data from the INfancia y Medio Ambiente birth cohort[66]. They described a catch-up trajectory[67] that identified children with low lung function early in childhood that catch up to average levels by age 10 years. Maternal age and paediatric allergic diseases were associated with lower odds of belonging to the high lung function growth trajectory, and maternal allergic diseases with lower odds of the catch-up trajectory. Children's higher physical activity and greater changes in BMI between 4 and 7 years of age were associated with the catch-up lung function growth trajectory.

This year, emphasis was on the effects of COVID-19 preventive measures on children's respiratory health. Ardura-Garcia *et al.* [68] showed that among 253 children seen in a respiratory outpatient clinic from the Swiss Paediatric Airway cohort[69], preventive measures reduced the reported respiratory infections, symptoms and medication use, but did not find differences on asthma exacerbation rates that required unscheduled medical care. School closure restrictions did not further reduce any respiratory outcome occurrence. Ghirardo *et al.* found that COVID-19 preventive measures in Italy did not delay the onset of bronchiolitis season in infants, but were related with lower number of hospitalisations, shorter hospitalization stays, lower oxygen supplementation and lower admissions

to ICU, which is suggestive of a less severe disease course[70]. Mallet *et al.* measured oxygen saturation and capillary blood gas after exercise testing in children with exercise induced symptoms while wearing face masks, and concluded that there was no evidence of abnormal gas exchange[71].

In conclusion, characterisation of PCD, identifying early life determinants and COVID-19 preventive measures on respiratory health are crucial for understanding and optimizing respiratory health in children.

Group 7.7: Paediatric bronchology

Several topics arose from this year's presentations in the field of paediatric bronchology; among others the incidence and prevalence of interstitial lung diseases, possible consequences of tracheomalacia and a classification of diffuse alveolar haemorrhage (DAH) in children.

Torrent *et al.*[72] analysed the incidence and prevalence of paediatric interstitial lung disease (chILD) in Spain during 2018-19. They found an average incidence of 8.18 cases/million/year and an average prevalence of 46.39 cases/million, which was higher compared to UK, Ireland, Germany, and Australasia. The authors believe these differences are due to greater understanding and increased awareness of chILD. Regarding age groups, the researchers found that children younger than 1 year of age have the highest prevalence of interstitial lung diseases while there is a heterogenous distribution of age groups according to different types of disorders: some were more prevalent in children from 0-2 years while others peaked in children from 2-18 years.

Ring *et al.*[73] collected 93 cases of DAH in children with a mean age of 4.2 years at debut of symptoms and 5.5 years at diagnosis, in an international, cross-sectional study. The most frequent underlying causes for DAH were idiopathic pulmonary hemosiderosis (44%), systemic disease (21%), immune-allergic disease (2%), autoinflammatory disease (4%) unspecified (13%) or other condition (15%). Children most frequently presented with anaemia (59%) or haemoptysis (42%), cough and dyspnoea were found in 37% and 35% respectively, and 19% suffered from fatigue.

Immunosuppressants were most administered, above all a combination of glucocorticoids with hydroxychloroquine, azathioprine, or both. Nine patients (10%) died. The authors intend to form a large, prospective multicentre study to standardize clinical care and test new treatment strategies based on their results.

Bloise *et al.*[74] proposed tidal breath (TB) analysis as a new diagnostic method in children with stridor. They evaluated the shape of the airflow signal using TB analysis in five children with stridor at first evaluation (T0) and in 3 infants after 3 months (T1). As demonstrated in Figure 1, they

discovered an abnormal morphological pattern of TB flow volume loops at T0. Additionally, they observed lower peak inspiratory flow and longer inspiratory time compared to healthy children. At T1, tidal volume had increased and respiratory rate and ratio of inspiratory time to expiratory time had decreased. The authors conclude that TB analysis could be an important non-invasive tool in the management of laryngomalacia.

A case-control study performed by Thomas *et al.*[75] assessed the risk of bronchiectasis in children with tracheomalacia. The research team retrospectively evaluated high resolution computed tomography (c-HRCT) for bronchiectasis and flexible bronchoscopies for tracheomalacia. They conducted both examinations in 45 cases with bronchiectasis and 90 controls without bronchiectasis under the age of 18 years within 4 weeks of time interval. After adjustment for age differences, they revealed an increased risk of developing bronchiectasis in children with tracheomalacia as defined by the ERS (adjusted Odds ratio 20.2 [CI 2.3-174.2]) compared to controls. They concluded that children with tracheomalacia should be monitored for chronic wet cough and, if present, investigated for bronchiectasis.

In conclusion, we learned of a higher incidence and prevalence of chiLD in Spain than previously reported; we introduced a large case collection of children with DAH, describing clinical characteristics and underlying pathologies; TB analysis technique was proposed for non-invasive management of laryngomalacia; and we identified tracheomalacia as a risk factor for bronchiectasis.

Group 7.8: Lung and airway developmental biology

The session “Mechanistic pathways in chronic and new lung diseases” overviewed current studies that aim to unravel novel mechanisms of chronic obstructive pulmonary disease (COPD), idiopathic pulmonary fibrosis (IPF), pulmonary hypertension (PH), primary ciliary dyskinesia (PCD), asthma, and the pulmonary sequelae of preterm birth.

Starting with the mesenchyme in COPD, Bekker *et al.* identified 774 proteins differentially regulated in lung fibroblasts from COPD patients, using proteomic analyses, where HNRNPA2B1, FHL1, TUBA1C, and HNRNPA1 represented novel candidate disease mediators or biomarkers that warranted functional follow-up studies[76]. Turning to infections in COPD patients, Brake *et al.* documented increased abundance of the SARS-CoV-2 entry- and co-receptors ACE2, TMPRSS2, and FURIN in type II pneumocytes, small airway epithelium, and alveolar macrophages in smokers and in COPD patients, raising the possibility of increased susceptibility of those individuals to severe SARS-CoV-2 infection[77]. Remaining with smokers, Paschalaki *et al.* revealed that endothelial-colony-forming-cells (ECFCs) from smokers and COPD patients exhibited increased expression of both senescence and

DNA damage response markers, and that inhalation of corticosteroids reduced the senescence and interferon γ -inducible-protein 10 in ECFC from COPD patients, thus suggesting a novel protective mechanism of corticosteroids[78]. The vasculature in smokers was not neglected, and Bhattarai *et al.* reported decreased arterial density and increased arterial wall thickness of vessels in smokers and mild-to-moderate COPD patients, indicative of early-onset of PH. The increased arterial wall thickness in the COPD smoker group affected FEV₁/FVC, which was reversed by smoking cessation[79]. Finally, with pulmonary malignancies in mind, Cerón Pisa *et al.* documented that microRNA (miR)-34c-5p expression could differentiate COPD patients with and without lung cancer, and that miR-200c-3p and miR-449c-5p expression was increased in the COPD patients[80]. These preliminary data highlight the potential utility of miRs as biomarkers in COPD.

Considering diffuse parenchymal lung disease, Gaikwad *et al.* performed a comprehensive quantitative analysis of structural changes to the pulmonary vasculature in IPF patients, where intimal thickness was highest in the arterial diameter range of 200-399 μm and $>600<1000$ μm in IPF patients, whilst medial and adventitial thickness was significantly increased across all arterial diameter ranges in IPF patients[81].

Turning to the airways, Pérez García *et al.* documented lower salivary microbiome diversity and more abundant *Rothia* spp. in asthma patients[82]. Moving on to viral infections in the airways, Gay *et al.* presented a transcriptional response to respiratory syncytial virus (RSV) in primary bronchial epithelial cells from asthmatic patients, where increased numbers of ciliated cells and decreased numbers of basal cells were noted in asthmatic patients after RSV infection[83]. Thus, the response to RSV infection is altered in asthma, with reduced inflammatory response and a shift in cell-type composition after infection. Remaining with the airways, Hjejij *et al.* revealed that DNAH5, DNAH11, CCDC40, CCNO, DNAI1 were the most frequent mutated genes in PCD patients[84].

Finally, the impact of prematurity on rabbit lung development was studied by Storti *et al.*, using histological and transcriptomic analyses in a preterm rabbit model, where radial alveolar count (RAC) progressively increased, whilst septal thickness and medial thickness of pulmonary arteries progressively decreased over the course of lung development. Transcriptomic analysis suggested that cell cycle, epithelial morphogenesis, and vascular endothelial growth factor signalling were more prominent in foetal samples, whilst immune system-related pathways were activated after birth[85].

In conclusion, this session highlighted that the characterization of genes and pathways involved in the development of a disease is essential to identify new biomarkers and therapeutic targets.

Concluding remarks

This comprehensive summary is only a glimpse of the high-quality research presented during the 2021 virtual ERS Congress Paediatric sessions. We highlighted the key advances in the paediatric respiratory field, with the aim of helping researchers and clinicians keep up to date with the latest and most relevant research. These research advances are also of vital importance for patients and their families, as they may reduce the time to diagnosis and improve their quality of life with more effective and targeted treatments. We encourage the interested readers to participate in the ERS International Congress 2022 that will take place in Barcelona, Spain and online, to meet the experts in the field and be exposed to the latest paediatric respiratory research first-hand.

Conflict of interests

CAG is supported by the Swiss National Science Foundation (grant number 32003B_162820), support for attending meetings from the ERS, and is the Early Career Member Representative of the Paediatrics Assembly of the European Respiratory Society (ERS). SCO's has received support for attending meetings from the ERS. MS received funding from the Chiesi Pharmaceutical S.p.a. for his PhD studies. RM has received grants or contracts from Deutsche Forschungsgemeinschaft, Max Planck Society, Universitätsklinikum Heidelberg, and Bundesministerium für Bildung und Forschung; honoraria as the Editor-in-Chief, of the American Journal of Physiology-Lung Cellular and Molecular Physiology; has a role as director of the International Graduate Programme, Molecular Biology and Medicine of the Lung, coordinator of the German Center for Lung Research Academy, Vice-Chair of the Gordon Conference of Lung Injury, Development, and Repair 2021, Chair of the Group 07.08 of the ERS, Chair of the Department of Translational Pulmonology at the University Hospital Heidelberg. AM has received consulting fees from Vertex Inc, lecture fees from Vertex Inc and Vifor Inc, has participated in the advisory board for Vertex Inc, and is the Secretary of the ERS Paediatric Assembly. MPijnenburg has received grants or contracts from Lung Foundation Netherlands and the Netherlands Organisation for Health Research and Development., consultation fees from Sanofi, honoraria as a speaker from Abbvie and Novartis, and support for attending meetings from the ERS, AA, LD, HD, NF, YTL, MS, EW, TD, RE, SF, MProesmans, SS, DS have nothing to disclose.

Acknowledgements:

Cristina Ardura-Garcia is supported by the Swiss National Science Foundation (grant number 32003B_162820). Matteo Storti is supported by Chiesi Pharmaceutical S.p.a.

References

1. Bush A. Mini-symposium: Impact of early life exposures on respiratory disease. *Paediatr Respir Rev* 2021.
2. Salem Mahmoud Y, Willers C, Amylidi-Mohr S, et al. Later lung function in birthweight discordant monozygotic twins. *Eur Respir J* 2021;58:1300.
3. Gudmundsdóttir H, Carlsen O, Stensby Bains K, et al. Infant lung function and physical activity in pregnancy. *Eur Respir J* 2021;58:1294.
4. Färdig M, Bains K, Gudmundsdóttir H, et al. Infant tidal flow-volume parameters and arousal state. *Eur Respir J* 2021;58:1297.
5. Fouzas S ESF, Dimos Gidarís, Nikolaos Karantaglis, Grigorios Chatziparasidis, Ilias, Theodorakopoulos MBA, Urs Frey. Fluctuation analysis of FEV1 in children and adolescents with controlled asthma. *Eur Respir J* 2021;58:1296.
6. Beydon N CF, Koskas M. Spirometry forced expiratory time is driven by central airway obstruction in asthmatic children. *Eur Respir J* 2021;58:1301.
7. Bednarek M, Grabicki M, Piorunek T, et al. Current place of impulse oscillometry in the assessment of pulmonary diseases. *Respir Med* 2020;170:105952.
8. Park J, Kim J, Kim M, et al. Impulse oscillometry findings compared to spirometry in patients with neuromuscular disorders. *Eur Respir J* 2021;58:1295.
9. Schlegtehdal A, Knoke L, Maier C, et al. Pulmonary function in children and adolescents after COVID-19. *Eur Respir J* 2021;58:1303.
10. Kaditis AG, Ohler A, Gileles-Hillel A, et al. Effects of the COVID-19 lockdown on sleep duration in children and adolescents: A survey across different continents. *Pediatr Pulmonol* 2021.
11. Kaditis A, Kheirandish-Gozal L, Gozal D. Algorithm for the diagnosis and treatment of pediatric OSA: a proposal of two pediatric sleep centers. *Sleep medicine* 2012;13:217-27.
12. Fouzas S, Pappa P, Goulioumis G, et al. Novel indices to improve the diagnostic ability of nocturnal oximetry in children with OSA. *Eur Respir J* 2021;58:1302.
13. Oyenuga V, Mosler G, Addo-Yobo E, et al. Asthma symptom severity and diagnosis in adolescents in sub-Saharan Africa. *Eur Respir J* 2021;58:2565.
14. Adeloye D, Chan KY, Rudan I, et al. An estimate of asthma prevalence in Africa: a systematic analysis. *Croat Med J* 2013;54:519-31.
15. Amaral R, Jácome C, Almeida R, et al. Phenotypes of persistent asthma in adolescents revealed different patterns in longitudinal asthma-related outcomes. *Eur Respir J* 2021;58:2566.
16. Manson ML, Safholm J, James A, et al. IL-13 and IL-4, but not IL-5 nor IL-17A, induce hyperresponsiveness in isolated human small airways. *J Allergy Clin Immunol* 2020;145:808-17 e2.
17. Ribas C, Papadopoulos N, Szeffler S, et al. Dupilumab efficacy in children with uncontrolled, moderate-to-severe asthma with and without an allergic phenotype. *Eur Respir J* 2021;58:2568.
18. Carraro S, Di Palma E, Barni S, et al. Metabolomic approach in the characterization of omalizumab responders and non-responders among children with severe asthma: a multicenter prospective study. *Eur Respir J* 2021;58:2567.

19. Andersson CK, Iwasaki J, Cook J, et al. Impaired airway epithelial cell wound-healing capacity is associated with airway remodelling following RSV infection in severe preschool wheeze. *Allergy* 2020;75:3195-207.
20. Beaufils F, Fayon M, Michelet M, et al. Bronchial remodeling-based latent class analysis predicts exacerbations in severe preschool wheezers. *Eur Respir J* 2021;58:2569.
21. Hose AJ, Pagani G, Karvonen AM, et al. Excessive Unbalanced Meat Consumption in the First Year of Life Increases Asthma Risk in the PASTURE and LUKAS2 Birth Cohorts. *Front Immunol* 2021;12:651709.
22. Hose A, Pagani G, Karvonen A, et al. Excessive meat consumption in the first year increases asthma risk in the PASTURE and LUKAS2 birth cohorts. *Eur Respir J* 2021;58:2570.
23. Da Silva Sena C, Shaar A, Morten M, et al. Rhinovirus bronchiolitis during infancy and later pre-school lung inhomogeneity. *Eur Respir J* 2021;58:2571.
24. Makrinioti H, Lakoumentas J, Xepapadaki P, et al. The role of respiratory syncytial virus and rhinovirus in early recurrent wheeze and asthma inception – a systematic review. *Eur Respir J* 2021;58:2572.
25. Gustafsson PM, Robinson PD, Lindblad A, et al. Novel methodology to perform sulfur hexafluoride (SF6)-based multiple-breath wash-in and washout in infants using current commercially available equipment. *J Appl Physiol (1985)* 2016;121:1087-97.
26. Sandvik RM, Gustafsson PM, Lindblad A, et al. Improved agreement between N2 and SF6 multiple-breath washout in healthy infants and toddlers with improved EXHALYZER D sensor performance. *J Appl Physiol (1985)* 2021;131:107-18.
27. Sandvik RM MNS, Jørgen Hovland Olsen, Sune Rubak, Anders Lindblad, Hanne Vebert Olesen, Frederik Buchvald, Tacjana Pressler, Marianne Skov, Per Gustafsson, Kim Gjerum Nielsen, K. G. Cross-sectional and longitudinal comparison of N2 and SF6 multiple breath washout in infants and toddlers with CF and healthy controls. *Eur Respir J* 2021;58:2670.
28. Stanojevic S, Davis SD, Retsch-Bogart G, et al. Progression of Lung Disease in Preschool Patients with Cystic Fibrosis. *Am J Respir Crit Care Med* 2017;195:1216-25.
29. Robinson PD, Latzin P, Ramsey KA, et al. Preschool Multiple-Breath Washout Testing. An Official American Thoracic Society Technical Statement. *Am J Respir Crit Care Med* 2018;197:e1-e19.
30. Dumas M, Karsli N, Jensen R, et al. Evaluation of SnIII parameters as outcome measures of multiple breath washout (MBW) in preschool children with cystic fibrosis. *Eur Respir J* 2021;58:2677.
31. Sandvik R, Schmidt M, Kragh M, et al. Longitudinal SF6 multiple breath washout in infants and toddlers with cystic fibrosis. *Eur Respir J* 2021;58:2671.
32. Oestreich MA, Wyler F, Latzin P, et al. Shedding light into the black box of infant multiple-breath washout. *Pediatr Pulmonol* 2021;56:2642-53.
33. Oestreich M, Manogaran T, Frauchiger B, et al. Evolution of lung clearance index during the first year of life in cystic fibrosis. *Eur Respir J* 2021;58:2673.
34. Stanojevic S, Davis SD, Perrem L, et al. Determinants of lung disease progression measured by lung clearance index in children with cystic fibrosis. *Eur Respir J* 2021;58.
35. Perrem L, Stanojevic S, Davis S, et al. Changes in the Lung Clearance Index outside 15% are clinically relevant in children with cystic fibrosis. *Eur Respir J* 2021;58:2672.
36. Frauchiger B, Ramsey K, Usemann J, et al. Natural Variability of Clinically Measured Lung Clearance Index in Children with Cystic Fibrosis. *Eur Respir J* 2021;58:2674.

37. Willers C, Frauchiger B, Stranzinger E, et al. Change in functional lung MRI and multiple breath washout outcomes over one year in children with cystic fibrosis. *Eur Respir J* 2021;58:2675.
38. Lv Q, Sandvik R, Nielsen K, et al. Sensitive automated airway-artery method to monitor progression of CF airway disease. *Eur Respir J* 2021;58:2676.
39. Grosse-Onnebrink J, Mete V, Kolonko K, et al. Measuring CFTR function in respiratory epithelial cell cultures: step to individualize treatments in cystic fibrosis? *Eur Respir J* 2021;58:2669.
40. Chang AB, Fortescue R, Grimwood K, et al. Task Force report: European Respiratory Society guidelines for the management of children and adolescents with bronchiectasis. *Eur Respir J* 2021:2002990.
41. Chang AB, Boyd J, Bell L, et al. Clinical and research priorities for children and young people with bronchiectasis: an international roadmap. *ERJ Open Res* 2021;7.
42. Indinnimeo L, Tancredi G, Barreto M, et al. Effects of a program of hospital-supervised chest physical therapy on lung function tests in children with chronic respiratory disease: 1-year follow-up. *Int J Immunopathol Pharmacol* 2007;20:841-5.
43. Goyal V, Grimwood K, Ware RS, et al. Efficacy of oral amoxicillin-clavulanate or azithromycin for non-severe respiratory exacerbations in children with bronchiectasis (BEST-1): a multicentre, three-arm, double-blind, randomised placebo-controlled trial. *Lancet Respir Med* 2019;7:791-801.
44. Valery PC, Morris PS, Byrnes CA, et al. Long-term azithromycin for Indigenous children with non-cystic-fibrosis bronchiectasis or chronic suppurative lung disease (Bronchiectasis Intervention Study): a multicentre, double-blind, randomised controlled trial. *Lancet Respir Med* 2013;1:610-20.
45. Greenough A. Long-Term Pulmonary Outcome in the Preterm Infant. *Neonatology* 2008;93:324-7.
46. Schmalisch G. Current methodological and technical limitations of time and volumetric capnography in newborns. *Biomed Eng Online* 2016;15:104.
47. Williams E, Dassios T, Greenough A. Feasibility of assessing ventilation inhomogeneity by volumetric capnography during resuscitation of preterm infants. *Eur Respir J* 2021;58:3954.
48. Nasim A, Zivanovic S, Abubakar A, et al. Less invasive surfactant administration (LISA) to preterm infants on high flow nasal cannula (nHFT). *Eur Respir J* 2021;58:3963.
49. Bizzotto DV, C. Bani, D. Dellaca, R. Effect of upper airways during non-invasive high frequency oscillatory ventilation (nHFOV). *Eur Respir J* 2021;58:3958.
50. Fouzas SB, A. Dimitriou, G. Influence of breathing dynamics on the effective concentration of oxygen delivered to infants via high-flow nasal cannula. *Eur Respir J* 2021;58:3957.
51. Arattu Thodika FW, E. Chappelow, I. Dassios, T. Greenough, A. Prediction of infant extubation success using the diaphragmatic electromyography (EMG) signal during a spontaneous breathing test (SBT). *Eur Respir J* 2021;58:3955.
52. Shetty S, Evans K, Kulkani A, et al. Neurally adjusted ventilatory assist (NAVA) in very preterm infants: A single tertiary neonatal unit's experience. *Eur Respir J* 2021;58:3960.
53. E. Z, C. R, A. A, et al. Lung function trajectories in very preterm infant. *Eur Respir J* 2021;58:3962.
54. Harris CL, A. Peacock, J. Greenough, A. Long term lung function in those born of multiple versus singleton pregnancies. *Eur Respir J* 2021;58:3956.

55. Luchnikova TP, O. Voitsekhovskiy, V. Romantsova, E. Pavlenko, V. . Health status of newborns from mothers with severe uncontrolled bronchial asthma. *Eur Respir J* 2021;58:3961.
56. Lee RW, E. Dassios, T. Greenough, A. Influence of antenatal corticosteroids and sex on the mortality and morbidity of extremely prematurely born infants. *Eur Respir J* 2021;58:3959.
57. Goutaki M, Papon J-F, Boon M, et al. Standardised clinical data from patients with primary ciliary dyskinesia: FOLLOW-PCD. *ERJ Open Res* 2020;6.
58. Goutaki M HL, Lam Y, Collaud E, Koppe H, Pedersen E, Kuehni C. Respiratory symptoms of Swiss people with Primary Ciliary Dyskinesia. *Eur Respir J* 2021;58:2956.
59. Rumman N, Driessens C, Goggin P, et al. The Palestinian primary ciliary dyskinesia (PCD) cohort: clinical, diagnostic and genetic spectrum. . *Eur Respir J* 2021;58:2957.
60. Fuchs O, Latzin P, Kuehni CE, et al. Cohort profile: the Bern infant lung development cohort. *Int J Epidemiol* 2012;41:366-76.
61. Decrue F, Gorlanova O, Salem Y, et al. Increased impact of air pollution on lung function in preterm vs. term infants: the BILD study. . *Eur Respir J* 2021;58:2958.
62. Kooijman MN, Kruithof CJ, van Duijn CM, et al. The Generation R Study: design and cohort update 2017. *Eur J Epidemiol* 2016;31:1243-64.
63. Abellan A WC, Warembourg C, Mensink-Bout S, Ambros A, de Castro M, Fossati S, Guxens M, Jaddoe V, Nieuwenhuijsen M, Vrijheid M, Santos S, Casas M, Duijts L. Urban environment during pregnancy and lung function, wheeze, and asthma in school-age children. The Generation R Study. . *Eur Respir J* 2021;58:2959.
64. Talaei M, Emmett P, Northstone K, et al. Dietary patterns and lung function in childhood: A longitudinal study. *Eur Respir J* 2021;58:2960.
65. Koch S PG, Carsin A, Cirugeda L, Sunyer J, Vrijheid M, Garcia-Esteban R, Abellan A, Guxens M, Torrent M, Ballester F, Ferrero A, Iñiguez C, Zabaleta C, Basterrechea M, Casas M, Garcia-Aymerich J. . Lung function trajectories and their determinants in 4-18 year-old children from the INMA birth cohort. . *Eur Respir J* 2021;58:2961.
66. Guxens M, Ballester F, Espada M, et al. Cohort profile: the INMA—INfancia y Medio Ambiente—(environment and childhood) project. *Int J Epidemiol* 2012;41:930-40.
67. Bui DS, Lodge CJ, Burgess JA, et al. Childhood predictors of lung function trajectories and future COPD risk: a prospective cohort study from the first to the sixth decade of life. *Lancet Respir Med* 2018;6:535-44.
68. Ardura-Garcia C MM, Pedersen E, Berger D, Barben J, Blanchon S, Casaulta C, Hoyler K, Jochmann A, Moeller A, Mueller-Suter D, Regamey N, Latzin P, Kuehni C. . Effect of COVID19 preventive measures on respiratory infections and symptoms in children. . *Eur Respir J* 2021;58:2962.
69. Pedersen ES, de Jong CC, Ardura-Garcia C, et al. The Swiss Paediatric Airway Cohort (SPAC). *ERJ Open Res* 2018;4.
70. Ghirardo S UN, Ciofi Degli Atti M, Raponi M, Cutrera R. . Delayed bronchiolitis season's onset during COVID-19 pandemic. *Eur Respir J* 2021;58:2963.
71. Mallet M HM, Lurà M, Kuehni C, Regamey N. . Do face masks cause abnormal gas exchange during exercise testing in children?. *Eur Respir J* 2021;58:2964.
72. Torrent Vernetta A, Rovira-Amigo S, Iglesias Serrano I, et al. Incidence and prevalence of children's diffuse lung disease in Spain. *Eur Respir J* 2021;58:4073.
73. Ring A, Nicolaus Schwerk, Nural Kiper, et al. Diffuse alveolar haemorrhage in children: an international multicentre study. *Eur Respir J* 2021;58:4069.

74. Bloise S, Nenna R, Petrarca L, et al. Respiratory function testing in infants with laryngeal stridor: diagnosis at a glance and follow up. *Eur Respir J* 2021;58:4072.
75. Thomas R, Chang A, Masters I, et al. Is there an association between tracheomalacia and bronchiectasis in children? *Eur Respir J* 2021;58:4068.
76. Bekke rN, Van Pijkeren A, Wolters J, et al. Proteomics approach to identify COPD-related changes in pulmonary fibroblasts. *Eur Respir J* 2021;58:1215.
77. Brake S, Wenying Lu W, Markos J, et al. Increased expression of SARS-CoV-2 (Covid-19) adhesion sites on type II pneumocytes, small airway epithelium, and alveolar macrophages in smokers and patients with COPD. *Eur Respir J* 2021;58:1217.
78. Paschalaki K, Rossios C, Gorgoulis V, et al. Inhaled corticosteroids reduce senescence in endothelial progenitor cells from COPD patients. *Eur Respir J* 2021;58:1214.
79. Bhattarai P, Lu W, Markos J, et al. Increased arterial remodelling in smokers and mild-moderate COPD patients are indicative of potential early onset of pulmonary hypertension. *Eur Respir J* 2021;58:1216.
80. Cerón Pisa N, Iglesias Coma A, Verdú Rivera F, et al. Usefulness of miRNAs as predictors of lung cancer in patients with COPD. A pilot study. *Eur Respir J* 2021;58:1219.
81. Gaikwad A, Lu W, Dey S, et al. Arterial remodelling in patients with idiopathic pulmonary fibrosis (IPF) and the possible role of endothelial to mesenchymal transition (EndMT). *Eur Respir J* 2021;58:1222.
82. Perez-Garcia J, Espuela-Ortiz A, Hernandez- Perez J, et al. Association of the salivary microbiome with asthma exacerbations. *Eur Respir J* 2021;58:1218.
83. Gay G, Berg B, Carpaij O, et al. Transcriptional response to RSV in asthma primary bronchial epithelial cells: are the basal cells responsible for the reduced NF-κB response? *Eur Respir J* 2021;58:1223.
84. Hjejij R, Cindric S, Olbrich H, et al. Diagnosis of Primary Ciliary Dyskinesia: a multi-center experience. *Eur Respir J* 2021;58:1221.
85. Storti M, Ricci F, Casiraghi C, et al. Impact of prematurity on rabbit lung development: a key factor for mimicking chronic neonatal lung disease. *Eur Respir J* 2021;58:1220.

Figure 1: (A) Typical tidal breath flow volume (TB-FV) loop in a healthy child and (B) TBFV loop in children with inspiratory stridor. *Figure modified from Bloise et al. [74]*

